

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


DH

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/10, C12Q 1/68	A2	(11) International Publication Number: WO 00/53744 (43) International Publication Date: 14 September 2000 (14.09.00)								
<p>(21) International Application Number: PCT/US00/06497</p> <p>(22) International Filing Date: 9 March 2000 (09.03.00)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>09/267,118</td> <td>9 March 1999 (09.03.99)</td> <td>US</td> </tr> <tr> <td>09/276,860</td> <td>26 March 1999 (26.03.99)</td> <td>US</td> </tr> <tr> <td>09/332,835</td> <td>14 June 1999 (14.06.99)</td> <td>US</td> </tr> </table> <p>(71) Applicant: DIVERSA CORPORATION [US/US]; 10665 Sorrento Valley Road, San Diego, CA 92121 (US).</p> <p>(72) Inventors: SHORT, Jay, M.; 6801 Paseo Delicias, Rancho Santa Fe, CA 92067 (US). FREY, Gerhard, Johann; Apartment 5214, 9952 Kika Court, San Diego, CA 92129 (US).</p> <p>(74) Agent: HAILE, Lisa, A.; Gray Cary Ware & Friedenrich LLP, Suite 1600, 4365 Executive Drive, San Diego, CA 92121-2189 (US).</p>	09/267,118	9 March 1999 (09.03.99)	US	09/276,860	26 March 1999 (26.03.99)	US	09/332,835	14 June 1999 (14.06.99)	US	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
09/267,118	9 March 1999 (09.03.99)	US								
09/276,860	26 March 1999 (26.03.99)	US								
09/332,835	14 June 1999 (14.06.99)	US								

(54) Title: END SELECTION IN DIRECTED EVOLUTION

End Selection Gene Assembly



(57) Abstract

This invention provides methods of obtaining novel polynucleotides and encoded polypeptides by the use of non-stochastic methods of directed evolution (DirectEvolution™). A particular advantage of end-selection-based methods is the ability to recover full-length polynucleotides from a library of progeny molecules generated by mutagenesis methods. These methods include non-stochastic polynucleotide site-saturation mutagenesis (Gene Site Saturation Mutagenesis™) and non-stochastic polynucleotide reassembly (GeneReassembly™). This invention provides methods of obtaining novel enzymes that have optimized physical and/or biological properties. Through use of the claimed methods, genetic vaccines, enzymes, small molecules, and other desirable molecules can be evolved towards desirable properties. For example, vaccine vectors can be obtained that exhibit increased efficacy for use as genetic vaccines. Vectors obtained by using the methods can have, for example, enhanced antigen expression, increased uptake into a cell, increased stability in a cell, ability to tailor an immune response, and the like. Furthermore, this invention provides methods of obtaining a variety of novel biologically active molecules, in the fields of antibiotics, pharmacotherapeutics, and transgenic traits.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

END SELECTION IN DIRECTED EVOLUTION

FIELD OF THE INVENTION

5

This invention relates to the field of protein engineering. More specifically, this relates to a directed evolution method for preparing a polynucleotides encoding polypeptide, which method comprises the step of generating site-directed mutagenesis optionally in combination with the step of polynucleotide chimerization, the step of
10 selecting for potentially desirable progeny molecules, including by a process termed end-selection (which may then be screened further), and the step of screening the polynucleotides for the production of polypeptide(s) having a useful property.

In a particular aspect, the present invention is relevant to enzymes, particularly
15 to thermostable enzymes, and to their generation by directed evolution. More particularly, the present invention relates to thermostable enzymes which are stable at high temperature and which have improved activity at lower temperatures.

20

BACKGROUND

Harvesting the full potential of nature's diversity can include both the step of discovery and the step of optimizing what is discovered. For example, the step of discovery allows one to mine biological molecules that have industrial utility. However,
25 for certain industrial needs, it is advantageous to further modify these enzymes experimentally to achieve properties beyond what natural evolution has provided and is likely to provide in the near future.

The process, termed directed evolution, of experimentally modifying a biological
30 molecule towards a desirable property, can be achieved by mutagenizing one or more parental molecular templates and identifying any desirable molecules among the progeny

molecules. However, currently available technologies used in directed evolution have several shortfalls. Among these shortfalls are:

- 5 1) Site-directed mutagenesis technologies, such as sloppy or low-fidelity PCR, are ineffective for systematically achieving at each position (site) along a polypeptide sequence the full (saturated) range of possible mutations (i.e. all possible amino acid substitutions).
- 10 2) There is no relatively easy systematic means for rapidly analyzing the large amount of information that can be contained in a molecular sequence and in the potentially colossal number or progeny molecules that could be conceivably obtained by the directed evolution of one or more molecular templates.
- 3) There is no relatively easy systematic means for providing comprehensive empirical information relating structure to function for molecular positions.
- 15 4) There is no easy systematic means for incorporating internal controls in certain mutagenesis (e.g. chimerization) procedures.
- 5) There is no easy systematic means to select for specific progeny molecules, such as full-length chimeras, from among smaller partial sequences.

Molecular mutagenesis occurs in nature and has resulted in the generation of a
20 wealth of biological compounds that have shown utility in certain industrial applications. However, evolution in nature often selects for molecular properties that are discordant with many unmet industrial needs. Additionally, it is often the case that when an industrially useful mutations would otherwise be favored at the molecular level, natural evolution often overrides the positive selection of such mutations when there is a
25 concurrent detriment to an organism as a whole (such as when a favorable mutation is accompanied by a detrimental mutation). Additionally still, natural evolution is slow, and places high emphasis on fidelity in replication. Finally, natural evolution prefers a path paved mainly by beneficial mutations while tending to avoid a plurality of successive negative mutations, even though such negative mutations may prove beneficial when
30 combined, or may lead - through a circuitous route - to final state that is beneficial.

Directed evolution, on the other hand, can be performed much more rapidly and aimed directly at evolving a molecular property that is industrially desirable where nature does not provide one.

5 An exceedingly large number of possibilities exist for purposeful and random combinations of amino acids within a protein to produce useful hybrid proteins and their corresponding biological molecules encoding for these hybrid proteins, i.e., DNA, RNA. Accordingly, there is a need to produce and screen a wide variety of such hybrid proteins for a desirable utility, particularly widely varying random proteins.

10

 The complexity of an active sequence of a biological macromolecule (e.g., polynucleotides, polypeptides, and molecules that are comprised of both polynucleotide and polypeptide sequences) has been called its information content ("IC"), which has been defined as the resistance of the active protein to amino acid sequence variation (calculated
15 from the minimum number of invariable amino acids (bits) required to describe a family of related sequences with the same function). Proteins that are more sensitive to random mutagenesis have a high information content.

 Molecular biology developments, such as molecular libraries, have allowed the
20 identification of quite a large number of variable bases, and even provide ways to select functional sequences from random libraries. In such libraries, most residues can be varied (although typically not all at the same time) depending on compensating changes in the context. Thus, while a 100 amino acid protein can contain only 2,000 different mutations, 20^{100} sequence combinations are possible.

25

 Information density is the IC per unit length of a sequence. Active sites of enzymes tend to have a high information density. By contrast, flexible linkers of information in enzymes have a low information density.

30 Current methods in widespread use for creating alternative proteins in a library format are error-prone polymerase chain reactions and cassette mutagenesis, in which

the specific region to be optimized is replaced with a synthetically mutagenized oligonucleotide. In both cases, a substantial number of mutant sites are generated around certain sites in the original sequence.

5 Error-prone PCR uses low-fidelity polymerization conditions to introduce a low level of point mutations randomly over a long sequence. In a mixture of fragments of unknown sequence, error-prone PCR can be used to mutagenize the mixture. The published error-prone PCR protocols suffer from a low processivity of the polymerase. Therefore, the protocol is unable to result in the random mutagenesis of an average-
10 sized gene. This inability limits the practical application of error-prone PCR. Some computer simulations have suggested that point mutagenesis alone may often be too gradual to allow the large-scale block changes that are required for continued and dramatic sequence evolution. Further, the published error-prone PCR protocols do not allow for amplification of DNA fragments greater than 0.5 to 1.0 kb, limiting their
15 practical application. In addition, repeated cycles of error-prone PCR can lead to an accumulation of neutral mutations with undesired results, such as affecting a protein's immunogenicity but not its binding affinity.

In oligonucleotide-directed mutagenesis, a short sequence is replaced with a
20 synthetically mutagenized oligonucleotide. This approach does not generate combinations of distant mutations and is thus not combinatorial. The limited library size relative to the vast sequence length means that many rounds of selection are unavoidable for protein optimization. Mutagenesis with synthetic oligonucleotides requires sequencing of individual clones after each selection round followed by
25 grouping them into families, arbitrarily choosing a single family, and reducing it to a consensus motif. Such motif is resynthesized and reinserted into a single gene followed by additional selection. This step process constitutes a statistical bottleneck, is labor intensive, and is not practical for many rounds of mutagenesis.

Error-prone PCR and oligonucleotide-directed mutagenesis are thus useful for single cycles of sequence fine tuning, but rapidly become too limiting when they are applied for multiple cycles.

5 Another limitation of error-prone PCR is that the rate of down-mutations grows with the information content of the sequence. As the information content, library size, and mutagenesis rate increase, the balance of down-mutations to up-mutations will statistically prevent the selection of further improvements (statistical ceiling).

10 In cassette mutagenesis, a sequence block of a single template is typically replaced by a (partially) randomized sequence. Therefore, the maximum information content that can be obtained is statistically limited by the number of random sequences (i.e., library size). This eliminates other sequence families which are not currently best, but which may have greater long term potential.

15

 Also, mutagenesis with synthetic oligonucleotides requires sequencing of individual clones after each selection round. Thus, such an approach is tedious and impractical for many rounds of mutagenesis.

20 Thus, error-prone PCR and cassette mutagenesis are best suited, and have been widely used, for fine-tuning areas of comparatively low information content. One apparent exception is the selection of an RNA ligase ribozyme from a random library using many rounds of amplification by error-prone PCR and selection.

25 In nature, the evolution of most organisms occurs by natural selection and sexual reproduction. Sexual reproduction ensures mixing and combining of the genes in the offspring of the selected individuals. During meiosis, homologous chromosomes from the parents line up with one another and cross-over part way along their length, thus randomly swapping genetic material. Such swapping or shuffling of the DNA allows
30 organisms to evolve more rapidly.

In recombination, because the inserted sequences were of proven utility in a homologous environment, the inserted sequences are likely to still have substantial information content once they are inserted into the new sequence.

5 The term Applied Molecular Evolution ("AME") means the application of an evolutionary design algorithm to a specific, useful goal. While many different library formats for AME have been reported for polynucleotides, peptides and proteins (phage, lacI and polysomes), none of these formats have provided for recombination by random cross-overs to deliberately create a combinatorial library.

10

 Theoretically there are 2,000 different single mutants of a 100 amino acid protein. However, a protein of 100 amino acids has 20^{100} possible sequence combinations, a number which is too large to exhaustively explore by conventional methods. It would be advantageous to develop a system which would allow generation
15 and screening of all of these possible combination mutations.

 Some workers in the art have utilized an *in vivo* site specific recombination system to generate hybrids of combine light chain antibody genes with heavy chain antibody genes for expression in a phage system. However, their system relies on
20 specific sites of recombination and is limited accordingly. Simultaneous mutagenesis of antibody CDR regions in single chain antibodies (scFv) by overlapping extension and PCR have been reported.

 Others have described a method for generating a large population of multiple
25 hybrids using random *in vivo* recombination. This method requires the recombination of two different libraries of plasmids, each library having a different selectable marker. The method is limited to a finite number of recombinations equal to the number of selectable markers existing, and produces a concomitant linear increase in the number of marker genes linked to the selected sequence(s).

30

In vivo recombination between two homologous, but truncated, insect-toxin genes on a plasmid has been reported as a method of producing a hybrid gene. The *in vivo* recombination of substantially mismatched DNA sequences in a host cell having defective mismatch repair enzymes, resulting in hybrid molecule formation has been
5 reported.

SUMMARY OF THE INVENTION

This invention relates generally to the field of nucleic acid engineering and
10 correspondingly encoded recombinant protein engineering. More particularly, the invention relates to the directed evolution of nucleic acids and screening of clones containing the evolved nucleic acids for resultant activity(ies) of interest, such nucleic acid activity(ies) &/or specified protein, particularly enzyme, activity(ies) of interest.

15 This invention relates generally to a method of: 1) preparing a progeny generation molecule (including a molecule that is comprised of a polynucleotide sequence, a molecules that is comprised of a polypeptide sequence, and a molecules that is comprised in part of a polynucleotide sequence and in part of a polypeptide sequence), that is mutagenized to achieve at least one point mutation, addition, deletion, &/or chimerization, from one or more
20 ancestral or parental generation template(s); 2) screening the progeny generation molecule - preferably using a high throughput method - for at least one property of interest (such as an improvement in an enzyme activity or an increase in stability or a novel chemotherapeutic effect); 3) optionally obtaining &/or cataloguing structural &/or and functional information regarding the parental &/or progeny generation molecules; and 4) optionally repeating any
25 of steps 1) to 3).

In a preferred embodiment, there is generated (e.g. from a parent polynucleotide template) - in what is termed "codon site-saturation mutagenesis" - a progeny
generation of polynucleotides, each having at least one set of up to three contiguous
30 point mutations (i.e. different bases comprising a new codon), such that every codon (or every family of degenerate codons encoding the same amino acid) is represented at each

codon position. Corresponding to - and encoded by - this progeny generation of polynucleotides, there is also generated a set of progeny polypeptides, each having at least one single amino acid point mutation. In a preferred aspect, there is generated - in what is termed "amino acid site-saturation mutagenesis" - one such mutant polypeptide
5 for each of the 19 naturally encoded polypeptide-forming alpha-amino acid substitutions at each and every amino acid position along the polypeptide. This yields - for each and every amino acid position along the parental polypeptide - a total of 20 distinct progeny polypeptides including the original amino acid, or potentially more than 21 distinct progeny polypeptides if additional amino acids are used either instead
10 of or in addition to the 20 naturally encoded amino acids

Thus, in another aspect, this approach is also serviceable for generating mutants containing - in addition to &/or in combination with the 20 naturally encoded polypeptide-forming alpha-amino acids - other rare &/or not naturally-encoded amino
15 acids and amino acid derivatives. In yet another aspect, this approach is also serviceable for generating mutants by the use of - in addition to &/or in combination with natural or unaltered codon recognition systems of suitable hosts - altered, mutagenized, &/or designer codon recognition systems (such as in a host cell with one or more altered tRNA molecules).

20

In yet another aspect, this invention relates to recombination and more specifically to a method for preparing polynucleotides encoding a polypeptide by a method of *in vivo* re-assortment of polynucleotide sequences containing regions of partial homology, assembling the polynucleotides to form at least one polynucleotide and screening the polynucleotides for
25 the production of polypeptide(s) having a useful property.

In yet another preferred embodiment, this invention is serviceable for analyzing and cataloguing - with respect to any molecular property (e.g. an enzymatic activity) or combination of properties allowed by current technology - the effects of any mutational
30 change achieved (including particularly saturation mutagenesis). Thus, a comprehensive method is provided for determining the effect of changing each amino acid in a parental

polypeptide into each of at least 19 possible substitutions. This allows each amino acid in a parental polypeptide to be characterized and catalogued according to its spectrum of potential effects on a measurable property of the polypeptide.

5 In another aspect, the method of the present invention utilizes the natural property of cells to recombine molecules and/or to mediate reductive processes that reduce the complexity of sequences and extent of repeated or consecutive sequences possessing regions of homology.

10 It is an object of the present invention to provide a method for generating hybrid polynucleotides encoding biologically active hybrid polypeptides with enhanced activities. In accomplishing these and other objects, there has been provided, in accordance with one aspect of the invention, a method for introducing polynucleotides into a suitable host cell and growing the host cell under conditions that produce a hybrid
15 polynucleotide.

 In another aspect of the invention, the invention provides a method for screening for biologically active hybrid polypeptides encoded by hybrid polynucleotides. The present method allows for the identification of biologically active hybrid polypeptides
20 with enhanced biological activities.

 Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred
25 embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

In a specific embodiment, this invention provides method for producing and isolating a library of progeny polynucleotides having at least one desirable property comprised of the steps of:

- 5 (a) subjecting a starting or parental polynucleotide set to a mutagenesis process so as to produce a progeny polynucleotide set; and
 - (b) subjecting the progeny polynucleotide set to an end selection-based screening and enrichment process, so as to select for a desirable subset of the progeny polynucleotide set;
- 10 whereby the above steps can be performed iteratively and in any order and in combination,
- whereby the end selection-based process creates ligation-compatible ends, whereby the creation of ligation-compatible ends is optionally used to facilitate one or more intermolecular ligations, that are preferably directional
- 15 ligations, within members of the progeny polynucleotide set so as to achieve assembly &/or reassembly mutagenesis,
- whereby the creation of ligation-compatible ends serves to facilitate ligation of the progeny polynucleotide set into an expression vector system and expression cloning,
- 20 whereby the end selection-based screening and enrichment process allows one to produce a library of progeny polynucleotides generated by a mutagenesis process, include non-stochastic polynucleotide site-saturation mutagenesis (Gene Site Saturation Mutagenesis™) and non-stochastic polynucleotide reassembly (GeneReassembly™),
- 25 whereby the expression cloning of the progeny polynucleotide set serves to generate a full-length polypeptide set,
- whereby the generated polypeptide set can be subjected to an expression screening process, and
- whereby expression screening of the progeny polypeptide set provides a means
- 30 to identify a desirable species, e.g. a mutant polypeptide or alternatively

a polypeptide fragment, that has a desirable property, such as a specific enzymatic activity.

5 In another specific embodiment, this invention provides a method for producing and isolating a polypeptide having at least one desirable property comprised of the steps of:

- (a) subjecting a starting or parental polynucleotide set to a mutagenesis process so as to produce a progeny polynucleotide set; and
- 10 (b) subjecting the progeny polynucleotide set to an end selection-based screening and enrichment process, so as to select for a desirable subset of the progeny polynucleotide set;

whereby the above steps can be performed iteratively and in any order and in
15 combination,

whereby the end selection-based process creates ligation-compatible ends,
whereby the creation of ligation-compatible ends is optionally used to facilitate
one or more intermolecular ligations, that are preferably directional
ligations, within members of the progeny polynucleotide set so as to
20 achieve assembly &/or reassembly mutagenesis,

whereby the end selection-based screening and enrichment process allows one to
produce a library of progeny polynucleotides generated by a mutagenesis
process, include non-stochastic polynucleotide site-saturation
mutagenesis (Gene Site Saturation Mutagenesis™) and non-stochastic
25 polynucleotide reassembly (GeneReassembly™),

whereby the expression cloning of the progeny polynucleotide set serves to
generate a full-length polypeptide set,

whereby the creation of ligation-compatible ends serves to facilitate ligation of
the progeny polynucleotide set into an expression vector system and
30 expression cloning,

whereby the generated polypeptide set can be subjected to an expression screening process, and

whereby expression screening of the progeny polypeptide set provides a means to identify a desirable species, e.g. a mutant polypeptide or alternatively a polypeptide fragment, that has a desirable property, such as a specific enzymatic activity.

In a specific aspect of this embodiment, this invention provides the immediately preceding methods, wherein the mutagenesis process of step (a) is comprised of a process, termed saturation mutagenesis, for generating, from a codon-containing parental polypeptide template, a progeny polypeptide set in which a full range of single amino acid substitutions is represented at each amino acid position, comprising the steps of:

(a) subjecting a working codon-containing template polynucleotide to polymerase-based amplification using a degenerate oligonucleotide for each codon to mutagenized, where each of said degenerate oligonucleotides is comprised of a first homologous sequence and a degenerate triplet sequence, so as to generate a set of progeny polynucleotides; wherein said degenerate triplet sequence is selected from the group consisting of i) N,N,N; ii) N,N,G/T; iii) N,N,G/C; iv) N,N,C/G/T; v) N,N,A/G/T; vi) N,N,A/C/T; vii) N,N,A/C/G; and viii) any degenerate codon that encodes all 20 amino acids; and

(b) subjecting said set of progeny polynucleotides to recombinant expression such that polypeptides encoded by the progeny polynucleotides are produced;

whereby the above steps can be performed iteratively and in any order and in combination, and

whereby, said method provides a means for generating all 20 amino acid changes at each amino acid site along a parental polypeptide template, because the degeneracy of the triplet sequence includes codons for all 20 amino acids.

In a specific aspect of this embodiment, this invention further provides the immediately preceding methods, wherein the mutagenesis process of step (a) is comprised of a
5 process, termed synthetic ligation gene reassembly or simply synthetic ligation gene reassembly.

BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1-9 correspond to Example 7.

5

Figure 1 and Figure 2 show the determination of the half-life of enzymes upon
subjection to an elevated temperature according to Example 7. Dhla20F12 has a $T_{1/2}$ of
about 11 minutes and longer incubation times at elevated temperature revealed an
10 approximate $T_{1/2}$ of 29 000 minutes for Dhla5 (Figure 2). Furthermore, 20F12
precipitates when heated while Dhla-5 does not. Therefore, Dhla-5 is at least 3000
times more stable than 20F12.

15 Figures 3-is a diagram illustrating the specific activity of Dhla5 at 30°C and 55°C in
comparison to that of Dhla20F12 according to Example 7. These data show that there is
no loss in activity when caused by the mutations and only thermal stability is affected.
The rate of Dhla5 is enhanced at 55°C to the same degree as Dhla20F12.

20

Figures 4-6 show data similar to those described in Figs 1-3, except using 80°C
incubations. The thermal stability of Dhla20F12 was not tested at this temperature but
it is assumed to be very short (on the order of seconds). Dhla5 has a $T_{1/2}$ of
approximately 13 minutes at 80°C. When Dhla8 is treated in the same way it has an
25 approximate $T_{1/2}$ of 138 minutes or about 10X Dhla5 (therefore 30 000 times
Dhla20F12).

Figures 7-9 is a diagram showing the use of differential calorimetry. Differential
30 scanning calorimetry (DSC) is a method used to determine melting temperatures (T_m)
and the enthalpy of thermal denaturation. Dhla20F12, Dhla5 and Dhla8 were all

- analyzed using DSC and the data are shown. The data show that T_m of 20F12 is about 68°C and is irreversible, Dhla5 has a T_m of 73°C and is partially reversible while Dhla8 also has a T_m of 73°C and is fully reversible. The mechanism of this reversibility is understood to involve a structural refolding after subjection of the molecule to
- 5 denaturation. Thus, this invention provides for the generation and for the selection of molecules to attain &/or to improve the ability to refold so as to regain activity after subjection to denaturation (eg. denaturation induced by temperature, salt, pH, or pressure changes).
- 10 Fig 10 is a diagram of the application of end-selection to select for desirable polynucleotides (e.g. the selection of full length molecules) generated by polynucleotide reassembly.

DEFINITIONS OF TERMS

In order to facilitate understanding of the examples provided herein, certain frequently occurring methods and/or terms will be described.

5

The term "**agent**" is used herein to denote a chemical compound, a mixture of chemical compounds, an array of spatially localized compounds (e.g., a VLSIPS peptide array, polynucleotide array, and/or combinatorial small molecule array), biological macromolecule, a bacteriophage peptide display library, a bacteriophage antibody (e.g., scFv) display library, a polysome peptide display library, or an extract made from biological materials such as bacteria, plants, fungi, or animal (particular mammalian) cells or tissues. Agents are evaluated for potential activity as anti-neoplastics, anti-inflammatories or apoptosis modulators by inclusion in screening assays described hereinbelow. Agents are evaluated for potential activity as specific protein interaction inhibitors (i.e., an agent which selectively inhibits a binding interaction between two predetermined polypeptides but which does not substantially interfere with cell viability) by inclusion in screening assays described hereinbelow.

An "**ambiguous base requirement**" in a restriction site refers to a nucleotide base requirement that is not specified to the fullest extent, i.e. that is not a specific base (such as, in a non-limiting exemplification, a specific base selected from A, C, G, and T), but rather may be any one of at least two or more bases. Commonly accepted abbreviations that are used in the art as well as herein to represent ambiguity in bases include the following: **R** = G or A; **Y** = C or T; **M** = A or C; **K** = G or T; **S** = G or C; **W** = A or T; **H** = A or C or T; **B** = G or T or C; **V** = G or C or A; **D** = G or A or T; **N** = A or C or G or T.

The term "**amino acid**" as used herein refers to any organic compound that contains an amino group (-NH₂) and a carboxyl group (-COOH); preferably either as free groups or alternatively after condensation as part of peptide bonds. The "**twenty naturally encoded polypeptide-forming alpha-amino acids**" are understood in the art

and refer to: alanine (ala or A), arginine (arg or R), asparagine (asn or N), aspartic acid (asp or D), cysteine (cys or C), glutamic acid (glu or E), glutamine (gln or Q), glycine (gly or G), histidine (his or H), isoleucine (ile or I), leucine (leu or L), lysine (lys or K), methionine (met or M), phenylalanine (phe or F), proline (pro or P), serine (ser or S),
5 threonine (thr or T), tryptophan (trp or W), tyrosine (tyr or Y), and valine (val or V).

The term "**amplification**" means that the number of copies of a polynucleotide is increased.

10 The term "**antibody**", as used herein, refers to intact immunoglobulin molecules, as well as fragments of immunoglobulin molecules, such as Fab, Fab', (Fab')₂, Fv, and SCA fragments, that are capable of binding to an epitope of an antigen. These antibody fragments, which retain some ability to selectively bind to an antigen (*e.g.*, a polypeptide antigen) of the antibody from which they are derived, can be made using
15 well known methods in the art (see, *e.g.*, **Harlow and Lane**, *supra*), and are described further, as follows.

(1) An Fab fragment consists of a monovalent antigen-binding fragment of an antibody molecule, and can be produced by digestion of a whole antibody
20 molecule with the enzyme papain, to yield a fragment consisting of an intact light chain and a portion of a heavy chain.

(2) An Fab' fragment of an antibody molecule can be obtained by treating a whole antibody molecule with pepsin, followed by reduction, to yield a
25 molecule consisting of an intact light chain and a portion of a heavy chain. Two Fab' fragments are obtained per antibody molecule treated in this manner.

(3) An (Fab')₂ fragment of an antibody can be obtained by treating a whole
30 antibody molecule with the enzyme pepsin, without subsequent reduction.

A (Fab')₂ fragment is a dimer of two Fab' fragments, held together by two disulfide bonds.

5 (4) An Fv fragment is defined as a genetically engineered fragment containing the variable region of a light chain and the variable region of a heavy chain expressed as two chains.

10 (5) An single chain antibody ("SCA") is a genetically engineered single chain molecule containing the variable region of a light chain and the variable region of a heavy chain, linked by a suitable, flexible polypeptide linker.

A molecule that has a "**chimeric property**" is a molecule that is: 1) in part homologous and in part heterologous to a first reference molecule; while 2) at the same time being in part homologous and in part heterologous to a second reference molecule; 15 without 3) precluding the possibility of being at the same time in part homologous and in part heterologous to still one or more additional reference molecules. In a non-limiting embodiment, a chimeric molecule may be prepared by assembling a reassortment of partial molecular sequences. In a non-limiting aspect, a chimeric polynucleotide molecule may be prepared by synthesizing the chimeric polynucleotide 20 using plurality of molecular templates, such that the resultant chimeric polynucleotide has properties of a plurality of templates.

The term "**cognate**" as used herein refers to a gene sequence that is evolutionarily and functionally related between species. For example, but not 25 limitation, in the human genome the human CD4 gene is the cognate gene to the mouse 3d4 gene, since the sequences and structures of these two genes indicate that they are highly homologous and both genes encode a protein which functions in signaling T cell activation through MHC class II-restricted antigen recognition.

30 A "**comparison window**," as used herein, refers to a conceptual segment of at least 20 contiguous nucleotide positions wherein a polynucleotide sequence may be

compared to a reference sequence of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith (**Smith and Waterman**, *Adv Appl Math*, 1981; **Smith and Waterman**, *J Teor Biol*, 1981; **Smith and Waterman**, *J Mol Biol*, 1981; **Smith et al**, *J Mol Evol*, 1981), by the homology alignment algorithm of Needleman (**Needleman and Wuncsch**, 1970), by the search of similarity method of Pearson (Pearson and Lipman, 1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, WI), or by inspection, and the best alignment (i.e., resulting in the highest percentage of homology over the comparison window) generated by the various methods is selected.

As used herein, the term "**complementarity-determining region**" and "**CDR**" refer to the art-recognized term as exemplified by the Kabat and Chothia CDR definitions also generally known as supervariable regions or hypervariable loops (**Chothia and Lesk**, 1987; **Chothia et al**, 1989; **Kabat et al**, 1987; and **Tramontano et al**, 1990). Variable region domains typically comprise the amino-terminal approximately 105-115 amino acids of a naturally-occurring immunoglobulin chain (e.g., amino acids 1-110), although variable domains somewhat shorter or longer are also suitable for forming single-chain antibodies.

25

"**Conservative amino acid substitutions**" refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group

30

of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are : valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine.

The term "**corresponds to**" is used herein to mean that a polynucleotide sequence is homologous (i.e., is identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide sequence, or that a polypeptide sequence is identical to a reference polypeptide sequence. In contradistinction, the term "complementary to" is used herein to mean that the complementary sequence is homologous to all or a portion of a reference polynucleotide sequence. For illustration, the nucleotide sequence "TATAC" corresponds to a reference "TATAC" and is complementary to a reference sequence "GTATA."

15

The term "**degrading effective**" amount refers to the amount of enzyme which is required to process at least 50% of the substrate, as compared to substrate not contacted with the enzyme. Preferably, at least 80% of the substrate is degraded.

As used herein, the term "**defined sequence framework**" refers to a set of defined sequences that are selected on a non-random basis, generally on the basis of experimental data or structural data; for example, a defined sequence framework may comprise a set of amino acid sequences that are predicted to form a β -sheet structure or may comprise a leucine zipper heptad repeat motif, a zinc-finger domain, among other variations. A "defined sequence kernal" is a set of sequences which encompass a limited scope of variability. Whereas (1) a completely random 10-mer sequence of the 20 conventional amino acids can be any of $(20)^{10}$ sequences, and (2) a pseudorandom 10-mer sequence of the 20 conventional amino acids can be any of $(20)^{10}$ sequences but will exhibit a bias for certain residues at certain positions and/or overall, (3) a defined sequence kernal is a subset of sequences if each residue position was allowed to be any of the allowable 20 conventional amino acids (and/or allowable unconventional

amino/imino acids). A defined sequence kernel generally comprises variant and invariant residue positions and/or comprises variant residue positions which can comprise a residue selected from a defined subset of amino acid residues), and the like, either segmentally or over the entire length of the individual selected library member
5 sequence. Defined sequence kernels can refer to either amino acid sequences or polynucleotide sequences. Of illustration and not limitation, the sequences (NNK)₁₀ and (NNM)₁₀, wherein N represents A, T, G, or C; K represents G or T; and M represents A or C, are defined sequence kernels.

10 **"Digestion"** of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 µg of plasmid or DNA fragment is used with about
15 2 units of enzyme in about 20 µl of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 µg of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's
20 instructions. After digestion the reaction is electrophoresed directly on a gel to isolate the desired fragment.

"Directional ligation" refers to a ligation in which a 5' end and a 3' end of a polynucleotide are different enough to specify a preferred ligation orientation. For
25 example, an otherwise untreated and undigested PCR product that has two blunt ends will typically not have a preferred ligation orientation when ligated into a cloning vector digested to produce blunt ends in its multiple cloning site; thus, directional ligation will typically not be displayed under these circumstances. In contrast, directional ligation will typically displayed when a digested PCR product having a 5' *EcoR* I-treated end
30 and a 3' *BamH* I-is ligated into a cloning vector that has a multiple cloning site digested with *EcoR* I and *BamH* I.

The term "**DNA shuffling**" is used herein to indicate recombination between substantially homologous but non-identical sequences, in some embodiments DNA shuffling may involve crossover via non-homologous recombination, such as via cer/lox and/or flp/frt systems and the like.

As used in this invention, the term "**epitope**" refers to an antigenic determinant on an antigen, such as a phytase polypeptide, to which the paratope of an antibody, such as an phytase-specific antibody, binds. Antigenic determinants usually consist of chemically active surface groupings of molecules, such as amino acids or sugar side chains, and can have specific three-dimensional structural characteristics, as well as specific charge characteristics. As used herein "**epitope**" refers to that portion of an antigen or other macromolecule capable of forming a binding interaction that interacts with the variable region binding body of an antibody. Typically, such binding interaction is manifested as an intermolecular contact with one or more amino acid residues of a CDR.

The terms "**fragment**", "**derivative**" and "**analog**" when referring to a reference polypeptide comprise a polypeptide which retains at least one biological function or activity that is at least essentially same as that of the reference polypeptide. Furthermore, the terms "**fragment**", "**derivative**" or "**analog**" are exemplified by a "pro-form" molecule, such as a low activity proprotein that can be modified by cleavage to produce a mature enzyme with significantly higher activity.

A method is provided herein for producing from a template polypeptide a set of progeny polypeptides in which a "**full range of single amino acid substitutions**" is represented at each amino acid position. As used herein, "**full range of single amino acid substitutions**" is in reference to the naturally encoded 20 naturally encoded polypeptide-forming alpha-amino acids, as described herein.

The term "**gene**" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons).

5

"**Genetic instability**", as used herein, refers to the natural tendency of highly repetitive sequences to be lost through a process of reductive events generally involving sequence simplification through the loss of repeated sequences. Deletions tend to involve the loss of one copy of a repeat and everything between the repeats.

10

The term "**heterologous**" means that one single-stranded nucleic acid sequence is unable to hybridize to another single-stranded nucleic acid sequence or its complement. Thus areas of heterology means that areas of polynucleotides or polynucleotides have areas or regions within their sequence which are unable to hybridize to another nucleic acid or polynucleotide. Such regions or areas are for example areas of mutations.

15

The term "**homologous**" or "**homeologous**" means that one single-stranded nucleic acid sequence may hybridize to a complementary single-stranded nucleic acid sequence. The degree of hybridization may depend on a number of factors including the amount of identity between the sequences and the hybridization conditions such as temperature and salt concentrations as discussed later. Preferably the region of identity is greater than about 5 bp, more preferably the region of identity is greater than 10 bp.

20
25

An immunoglobulin light or heavy chain variable region consists of a "framework" region interrupted by three hypervariable regions, also called CDR's. The extent of the framework region and CDR's have been precisely defined; see "Sequences of Proteins of Immunological Interest" (Kabat et al, 1987). The sequences of the framework regions of different light or heavy chains are relatively conserved within a specie. As used herein, a "**human framework region**" is a framework region that is

30

substantially identical (about 85 or more, usually 90-95 or more) to the framework region of a naturally occurring human immunoglobulin. the framework region of an antibody, that is the combined framework regions of the constituent light and heavy chains, serves to position and align the CDR's. The CDR's are primarily responsible for
5 binding to an epitope of an antigen.

The benefits of this invention extend to "**industrial applications**" (or industrial processes), which term is used to include applications in commercial industry proper (or simply industry) as well as non-commercial industrial applications (e.g. biomedical
10 research at a non-profit institution). Relevant applications include those in areas of diagnosis, medicine, agriculture, manufacturing, and academia.

The term "**identical**" or "**identity**" means that two nucleic acid sequences have the same sequence or a complementary sequence. Thus, "areas of identity" means that
15 regions or areas of a polynucleotide or the overall polynucleotide are identical or complementary to areas of another polynucleotide or the polynucleotide.

The term "**isolated**" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a
20 naturally-occurring polynucleotide or enzyme present in a living animal is not isolated, but the same polynucleotide or enzyme, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or enzymes could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

25

By "**isolated nucleic acid**" is meant a nucleic acid, e.g., a DNA or RNA molecule, that is not immediately contiguous with the 5' and 3' flanking sequences with which it normally is immediately contiguous when present in the naturally occurring genome of the organism from which it is derived. The term thus describes, for example,
30 a nucleic acid that is incorporated into a vector, such as a plasmid or viral vector; a nucleic acid that is incorporated into the genome of a heterologous cell (or the genome

of a homologous cell, but at a site different from that at which it naturally occurs); and a nucleic acid that exists as a separate molecule, e.g., a DNA fragment produced by PCR amplification or restriction enzyme digestion, or an RNA molecule produced by *in vitro* transcription. The term also describes a recombinant nucleic acid that forms part of a hybrid gene encoding additional polypeptide sequences that can be used, for example, in the production of a fusion protein.

As used herein "**ligand**" refers to a molecule, such as a random peptide or variable segment sequence, that is recognized by a particular receptor. As one of skill in the art will recognize, a molecule (or macromolecular complex) can be both a receptor and a ligand. In general, the binding partner having a smaller molecular weight is referred to as the ligand and the binding partner having a greater molecular weight is referred to as a receptor.

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments (Sambrook et al, 1982, p. 146; Sambrook, 1989). Unless otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units of T4 DNA ligase ("ligase") per 0.5 µg of approximately equimolar amounts of the DNA fragments to be ligated.

As used herein, "**linker**" or "**spacer**" refers to a molecule or group of molecules that connects two molecules, such as a DNA binding protein and a random peptide, and serves to place the two molecules in a preferred configuration, e.g., so that the random peptide can bind to a receptor with minimal steric hindrance from the DNA binding protein.

As used herein, a "**molecular property to be evolved**" includes reference to molecules comprised of a polynucleotide sequence, molecules comprised of a polypeptide sequence, and molecules comprised in part of a polynucleotide sequence and in part of a polypeptide sequence. Particularly relevant - but by no means limiting - examples of molecular properties to be evolved include enzymatic activities at specified

conditions, such as related to temperature; salinity; pressure; pH; and concentration of glycerol, DMSO, detergent, &/or any other molecular species with which contact is made in a reaction environment. Additional particularly relevant - but by no means limiting - examples of molecular properties to be evolved include stabilities - e.g. the
5 amount of a residual molecular property that is present after a specified exposure time to a specified environment, such as may be encountered during storage.

The term "**mutations**" means changes in the sequence of a wild-type nucleic acid sequence or changes in the sequence of a peptide. Such mutations may be point
10 mutations such as transitions or transversions. The mutations may be deletions, insertions or duplications.

As used herein, the degenerate "**N,N,G/T**" nucleotide sequence represents 32 possible triplets, where "**N**" can be A, C, G or T.
15

The term "**naturally-occurring**" as used herein as applied to the object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man
20 in the laboratory is naturally occurring. Generally, the term naturally occurring refers to an object as present in a non-pathological (un-diseased) individual, such as would be typical for the species.

As used herein, a "**nucleic acid molecule**" is comprised of at least one base or
25 one base pair, depending on whether it is single-stranded or double-stranded, respectively. Furthermore, a nucleic acid molecule may belong exclusively or chimerically to any group of nucleotide-containing molecules, as exemplified by, but not limited to, the following groups of nucleic acid molecules: RNA, DNA, genomic nucleic acids, non-genomic nucleic acids, naturally occurring and not naturally
30 occurring nucleic acids, and synthetic nucleic acids. This includes, by way of non-limiting example, nucleic acids associated with any organelle, such as the mitochondria,

ribosomal RNA, and nucleic acid molecules comprised chimerically of one or more components that are not naturally occurring along with naturally occurring components.

Additionally, a **"nucleic acid molecule"** may contain in part one or more non-
5 nucleotide-based components as exemplified by, but not limited to, amino acids and sugars. Thus, by way of example, but not limitation, a ribozyme that is in part nucleotide-based and in part protein-based is considered a **"nucleic acid molecule"**.

In addition, by way of example, but not limitation, a nucleic acid molecule that
10 is labeled with a detectable moiety, such as a radioactive or alternatively a non-radioactive label, is likewise considered a **"nucleic acid molecule"**.

The terms **"nucleic acid sequence coding for"** or a **"DNA coding sequence of"**
or a **"nucleotide sequence encoding"** a particular enzyme – as well as other
15 synonymous terms – refer to a DNA sequence which is transcribed and translated into an enzyme when placed under the control of appropriate regulatory sequences. A "promotor sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. The promoter is part of the DNA sequence. This sequence region has a start codon at
20 its 3' terminus. The promoter sequence does include the minimum number of bases where elements necessary to initiate transcription at levels detectable above background. However, after the RNA polymerase binds the sequence and transcription is initiated at the start codon (3' terminus with a promoter), transcription proceeds downstream in the 3' direction. Within the promotor sequence will be found a
25 transcription initiation site (conveniently defined by mapping with nuclease S1) as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

The terms **"nucleic acid encoding an enzyme (protein)"** or **"DNA encoding an
30 enzyme (protein)"** or **"polynucleotide encoding an enzyme (protein)"** and other synonymous terms encompasses a polynucleotide which includes only coding sequence

for the enzyme as well as a polynucleotide which includes additional coding and/or non-coding sequence.

In one preferred embodiment, a **“specific nucleic acid molecule species”** is defined by its chemical structure, as exemplified by, but not limited to, its primary sequence. In another preferred embodiment, a specific **“nucleic acid molecule species”** is defined by a function of the nucleic acid species or by a function of a product derived from the nucleic acid species. Thus, by way of non-limiting example, a **“specific nucleic acid molecule species”** may be defined by one or more activities or properties attributable to it, including activities or properties attributable its expressed product.

The instant definition of **“assembling a working nucleic acid sample into a nucleic acid library”** includes the process of incorporating a nucleic acid sample into a vector-based collection, such as by ligation into a vector and transformation of a host. A description of relevant vectors, hosts, and other reagents as well as specific non-limiting examples thereof are provided hereinafter. The instant definition of **“assembling a working nucleic acid sample into a nucleic acid library”** also includes the process of incorporating a nucleic acid sample into a non-vector-based collection, such as by ligation to adaptors. Preferably the adaptors can anneal to PCR primers to facilitate amplification by PCR.

Accordingly, in a non-limiting embodiment, a **“nucleic acid library”** is comprised of a vector-based collection of one or more nucleic acid molecules. In another preferred embodiment a **“nucleic acid library”** is comprised of a non-vector-based collection of nucleic acid molecules. In yet another preferred embodiment a **“nucleic acid library”** is comprised of a combined collection of nucleic acid molecules that is in part vector-based and in part non-vector-based. Preferably, the collection of molecules comprising a library is searchable and separable according to individual nucleic acid molecule species.

The present invention provides a **“nucleic acid construct”** or alternatively a **“nucleotide construct”** or alternatively a **“DNA construct”**. The term "construct" is used herein to describe a molecule, such as a polynucleotide (*e.g.*, a phytase polynucleotide) may optionally be chemically bonded to one or more additional
5 molecular moieties, such as a vector, or parts of a vector. In a specific - but by no means limiting - aspect, a nucleotide construct is exemplified by a DNA expression DNA expression constructs suitable for the transformation of a host cell.

An **“oligonucleotide”** (or synonymously an **“oligo”**) refers to either a single
10 stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands which may be chemically synthesized. Such synthetic oligonucleotides may or may not have a 5' phosphate. Those that do not will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide will ligate to a fragment that has not been dephosphorylated. To
15 achieve polymerase-based amplification (such as with PCR), a **“32-fold degenerate oligonucleotide that is comprised of, in series, at least a first homologous sequence, a degenerate N,N,G/T sequence, and a second homologous sequence”** is mentioned. As used in this context, “homologous” is in reference to homology between the oligo and the parental polynucleotide that is subjected to the polymerase-based amplification.

20

As used herein, the term **“operably linked”** refers to a linkage of polynucleotide elements in a functional relationship. A nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the
25 transcription of the coding sequence. Operably linked means that the DNA sequences being linked are typically contiguous and, where necessary to join two protein coding regions, contiguous and in reading frame.

A coding sequence is **“operably linked to”** another coding sequence when RNA
30 polymerase will transcribe the two coding sequences into a single mRNA, which is then translated into a single polypeptide having amino acids derived from both coding

sequences. The coding sequences need not be contiguous to one another so long as the expressed sequences are ultimately processed to produce the desired protein.

As used herein the term "**parental polynucleotide set**" is a set comprised of one or more distinct polynucleotide species. Usually this term is used in reference to a progeny polynucleotide set which is preferably obtained by mutagenization of the parental set, in which case the terms "**parental**", "**starting**" and "**template**" are used interchangeably.

As used herein the term "**physiological conditions**" refers to temperature, pH, ionic strength, viscosity, and like biochemical parameters which are compatible with a viable organism, and/or which typically exist intracellularly in a viable cultured yeast cell or mammalian cell. For example, the intracellular conditions in a yeast cell grown under typical laboratory culture conditions are physiological conditions. Suitable *in vitro* reaction conditions for *in vitro* transcription cocktails are generally physiological conditions. In general, *in vitro* physiological conditions comprise 50-200 mM NaCl or KCl, pH 6.5-8.5, 20-45°C and 0.001-10 mM divalent cation (e.g., Mg^{++} , Ca^{++}); preferably about 150 mM NaCl or KCl, pH 7.2-7.6, 5 mM divalent cation, and often include 0.01-1.0 percent nonspecific protein (e.g., BSA). A non-ionic detergent (Tween, NP-40, Triton X-100) can often be present, usually at about 0.001 to 2%, typically 0.05-0.2% (v/v). Particular aqueous conditions may be selected by the practitioner according to conventional methods. For general guidance, the following buffered aqueous conditions may be applicable: 10-250 mM NaCl, 5-50 mM Tris HCl, pH 5-8, with optional addition of divalent cation(s) and/or metal chelators and/or non-ionic detergents and/or membrane fractions and/or anti-foam agents and/or scintillants.

Standard convention (5' to 3') is used herein to describe the sequence of double stranded polynucleotides.

The term "**population**" as used herein means a collection of components such as polynucleotides, portions or polynucleotides or proteins. A "mixed population" means a

collection of components which belong to the same family of nucleic acids or proteins (i.e., are related) but which differ in their sequence (i.e., are not identical) and hence in their biological activity.

5 A molecule having a “**pro-form**” refers to a molecule that undergoes any combination of one or more covalent and noncovalent chemical modifications (e.g. glycosylation, proteolytic cleavage, dimerization or oligomerization, temperature-induced or pH-induced conformational change, association with a co-factor, etc.) en route to attain a more mature molecular form having a property difference (e.g. an
10 increase in activity) in comparison with the reference pro-form molecule. When two or more chemical modification (e.g. two proteolytic cleavages, or a proteolytic cleavage and a deglycosylation) can be distinguished en route to the production of a mature molecule, the reference precursor molecule may be termed a “**pre-pro-form**” molecule.

15 As used herein, the term “**pseudorandom**” refers to a set of sequences that have limited variability, such that, for example, the degree of residue variability at another position, but any pseudorandom position is allowed some degree of residue variation, however circumscribed.

20 “**Quasi-repeated units**”, as used herein, refers to the repeats to be re-assorted and are by definition not identical. Indeed the method is proposed not only for practically identical encoding units produced by mutagenesis of the identical starting sequence, but also the reassortment of similar or related sequences which may diverge
25 significantly in some regions. Nevertheless, if the sequences contain sufficient homologies to be reassorted by this approach, they can be referred to as “quasi-repeated” units.

 As used herein “**random peptide library**” refers to a set of polynucleotide
30 sequences that encodes a set of random peptides, and to the set of random peptides

encoded by those polynucleotide sequences, as well as the fusion proteins contain those random peptides.

As used herein, "**random peptide sequence**" refers to an amino acid sequence
5 composed of two or more amino acid monomers and constructed by a stochastic or random process. A random peptide can include framework or scaffolding motifs, which may comprise invariant sequences.

As used herein, "**receptor**" refers to a molecule that has an affinity for a given
10 ligand. Receptors can be naturally occurring or synthetic molecules. Receptors can be employed in an unaltered state or as aggregates with other species. Receptors can be attached, covalently or non-covalently, to a binding member, either directly or via a specific binding substance. Examples of receptors include, but are not limited to, antibodies, including monoclonal antibodies and antisera reactive with specific
15 antigenic determinants (such as on viruses, cells, or other materials), cell membrane receptors, complex carbohydrates and glycoproteins, enzymes, and hormone receptors.

"**Recombinant**" enzymes refer to enzymes produced by recombinant DNA techniques, i.e., produced from cells transformed by an exogenous DNA construct
20 encoding the desired enzyme. "**Synthetic**" enzymes are those prepared by chemical synthesis.

The term "**related polynucleotides**" means that regions or areas of the polynucleotides are identical and regions or areas of the polynucleotides are
25 heterologous.

"**Reductive reassortment**", as used herein, refers to the increase in molecular diversity that is accrued through deletion (and/or insertion) events that are mediated by repeated sequences.
30

The following terms are used to describe the sequence relationships between two or more polynucleotides: "**reference sequence**," "**comparison window**," "**sequence identity**," "**percentage of sequence identity**," and "**substantial identity**."

5 A "**reference sequence**" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing, or may comprise a complete cDNA or gene sequence. Generally, a reference sequence is at least 20 nucleotides in length, frequently at least 25 nucleotides in length, and often at
10 least 50 nucleotides in length. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides and (2) may further comprise a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two
15 polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity.

"**Repetitive Index (RI)**", as used herein, is the average number of copies of the quasi-repeated units contained in the cloning vector.

20

The term "**restriction site**" refers to a recognition sequence that is necessary for the manifestation of the action of a restriction enzyme, and includes a site of catalytic cleavage. It is appreciated that a site of cleavage may or may not be contained within a portion of a restriction site that comprises a low ambiguity sequence (i.e. a sequence
25 containing the principal determinant of the frequency of occurrence of the restriction site). Thus, in many cases, relevant restriction sites contain only a low ambiguity sequence with an internal cleavage site (e.g. G/AATTC in the EcoR I site) or an immediately adjacent cleavage site (e.g. /CCWGG in the EcoR II site). In other cases, relevant restriction enzymes [e.g. the Eco57 I site or CTGAAG(16/14)] contain a low
30 ambiguity sequence (e.g. the CTGAAG sequence in the Eco57 I site) with an external cleavage site (e.g. in the N₁₆ portion of the Eco57 I site). When an enzyme (e.g. a

restriction enzyme) is said to “cleave” a polynucleotide, it is understood to mean that the restriction enzyme catalyzes or facilitates a cleavage of a polynucleotide.

In a non-limiting aspect, a “selectable polynucleotide” is comprised of a 5’ terminal region (or end region), an intermediate region (i.e. an internal or central region), and a 3’ terminal region (or end region). As used in this aspect, a 5’ terminal region is a region that is located towards a 5’ polynucleotide terminus (or a 5’ polynucleotide end); thus it is either partially or entirely in a 5’ half of a polynucleotide. Likewise, a 3’ terminal region is a region that is located towards a 3’ polynucleotide terminus (or a 3’ polynucleotide end); thus it is either partially or entirely in a 3’ half of a polynucleotide. As used in this non-limiting exemplification, there may be sequence overlap between any two regions or even among all three regions.

The term “sequence identity” means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term “percentage of sequence identity” is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. This “substantial identity”, as used herein, denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence having at least 80 percent sequence identity, preferably at least 85 percent identity, often 90 to 95 percent sequence identity, and most commonly at least 99 percent sequence identity as compared to a reference sequence of a comparison window of at least 25-50 nucleotides, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the window of comparison.

As known in the art "**similarity**" between two enzymes is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one enzyme to the sequence of a second enzyme. Similarity may be determined by procedures which are well-known in the art, for example, a BLAST program (Basic
5 Local Alignment Search Tool at the National Center for Biological Information).

As used herein, the term "**single-chain antibody**" refers to a polypeptide comprising a V_H domain and a V_L domain in polypeptide linkage, generally linked via a spacer peptide (e.g., [Gly-Gly-Gly-Gly-Ser]_x), and which may comprise additional
10 amino acid sequences at the amino- and/or carboxy- termini. For example, a single-chain antibody may comprise a tether segment for linking to the encoding polynucleotide. As an example, a scFv is a single-chain antibody. Single-chain antibodies are generally proteins consisting of one or more polypeptide segments of at least 10 contiguous amino substantially encoded by genes of the immunoglobulin
15 superfamily (e.g., see **Williams and Barclay**, 1989, pp. 361-368, which is incorporated herein by reference), most frequently encoded by a rodent, non-human primate, avian, porcine bovine, ovine, goat, or human heavy chain or light chain gene sequence. A functional single-chain antibody generally contains a sufficient portion of an immunoglobulin superfamily gene product so as to retain the property of binding to a
20 specific target molecule, typically a receptor or antigen (epitope).

The members of a pair of molecules (e.g., an antibody-antigen pair or a nucleic acid pair) are said to "**specifically bind**" to each other if they bind to each other with greater affinity than to other, non-specific molecules. For example, an antibody raised
25 against an antigen to which it binds more efficiently than to a non-specific protein can be described as specifically binding to the antigen. (Similarly, a nucleic acid probe can be described as specifically binding to a nucleic acid target if it forms a specific duplex with the target by base pairing interactions (see above).)

30 "**Specific hybridization**" is defined herein as the formation of hybrids between a first polynucleotide and a second polynucleotide (e.g., a polynucleotide having a

distinct but substantially identical sequence to the first polynucleotide), wherein substantially unrelated polynucleotide sequences do not form hybrids in the mixture.

5 The term "**specific polynucleotide**" means a polynucleotide having certain end points and having a certain nucleic acid sequence. Two polynucleotides wherein one polynucleotide has the identical sequence as a portion of the second polynucleotide but different ends comprises two different specific polynucleotides.

10 "**Stringent hybridization conditions**" means hybridization will occur only if there is at least 90% identity, preferably at least 95% identity and most preferably at least 97% identity between the sequences. See Sambrook et al, 1989, which is hereby incorporated by reference in its entirety.

Also included in the invention are polypeptides having sequences that are
15 "**substantially identical**" to the sequence of a phytase polypeptide, such as one of SEQ ID 1. A "**substantially identical**" amino acid sequence is a sequence that differs from a reference sequence only by conservative amino acid substitutions, for example, substitutions of one amino acid for another of the same class (e.g., substitution of one hydrophobic amino acid, such as isoleucine, valine, leucine, or methionine, for another,
20 or substitution of one polar amino acid for another, such as substitution of arginine for lysine, glutamic acid for aspartic acid, or glutamine for asparagine).

Additionally a "**substantially identical**" amino acid sequence is a sequence that differs from a reference sequence or by one or more non-conservative substitutions,
25 deletions, or insertions, particularly when such a substitution occurs at a site that is not the active site the molecule, and provided that the polypeptide essentially retains its behavioural properties. For example, one or more amino acids can be deleted from a phytase polypeptide, resulting in modification of the structure of the polypeptide, without significantly altering its biological activity. For example, amino- or carboxyl-
30 terminal amino acids that are not required for phytase biological activity can be

removed. Such modifications can result in the development of smaller active phytase polypeptides.

The present invention provides a "**substantially pure enzyme**". The term
5 "**substantially pure enzyme**" is used herein to describe a molecule, such as a polypeptide (*e.g.*, a phytase polypeptide, or a fragment thereof) that is substantially free of other proteins, lipids, carbohydrates, nucleic acids, and other biological materials with which it is naturally associated. For example, a substantially pure molecule, such as a polypeptide, can be at least 60%, by dry weight, the molecule of interest. The
10 purity of the polypeptides can be determined using standard methods including, *e.g.*, polyacrylamide gel electrophoresis (*e.g.*, SDS-PAGE), column chromatography (*e.g.*, high performance liquid chromatography (HPLC)), and amino-terminal amino acid sequence analysis.

15 As used herein, "**substantially pure**" means an object species is the predominant species present (*i.e.*, on a molar basis it is more abundant than any other individual macromolecular species in the composition), and preferably substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present. Generally, a
20 substantially pure composition will comprise more than about 80 to 90 percent of all macromolecular species present in the composition. Most preferably, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species. Solvent species, small molecules (<500
25 Daltons), and elemental ion species are not considered macromolecular species.

As used herein, the term "**variable segment**" refers to a portion of a nascent peptide which comprises a random, pseudorandom, or defined kernal sequence. A
30 "variable segment" refers to a portion of a nascent peptide which comprises a random pseudorandom, or defined kernal sequence. A variable segment can comprise both variant and invariant residue positions, and the degree of residue variation at a variant

residue position may be limited: both options are selected at the discretion of the practitioner. Typically, variable segments are about 5 to 20 amino acid residues in length (e.g., 8 to 10), although variable segments may be longer and may comprise antibody portions or receptor proteins, such as an antibody fragment, a nucleic acid binding protein, a receptor protein, and the like.

The term "**wild-type**" means that the polynucleotide does not comprise any mutations. A "wild type" protein means that the protein will be active at a level of activity found in nature and will comprise the amino acid sequence found in nature.

The term "**working**", as in "**working sample**", for example, is simply a sample with which one is working. Likewise, a "**working molecule**", for example is a molecule with which one is working.

DETAILED DESCRIPTION OF THE INVENTION

The invention described herein is directed to the use of repeated cycles of reductive reassortment, recombination and selection which allow for the directed
5 molecular evolution of highly complex linear sequences, such as DNA, RNA or proteins thorough recombination.

In vivo shuffling of molecules can be performed utilizing the natural property of cells to recombine multimers. While recombination *in vivo* has provided the major
10 natural route to molecular diversity, genetic recombination remains a relatively complex process that involves 1) the recognition of homologies; 2) strand cleavage, strand invasion, and metabolic steps leading to the production of recombinant chiasma; and finally 3) the resolution of chiasma into discrete recombined molecules. The formation of the chiasma requires the recognition of homologous sequences.

15

In a preferred embodiment, the invention relates to a method for producing a hybrid polynucleotide from at least a first polynucleotide and a second polynucleotide. The present invention can be used to produce a hybrid polynucleotide by introducing at least a first polynucleotide and a second polynucleotide which share at least one region
20 of partial sequence homology into a suitable host cell. The regions of partial sequence homology promote processes which result in sequence reorganization producing a hybrid polynucleotide. The term "hybrid polynucleotide", as used herein, is any nucleotide sequence which results from the method of the present invention and contains sequence from at least two original polynucleotide sequences. Such hybrid
25 polynucleotides can result from intermolecular recombination events which promote sequence integration between DNA molecules. In addition, such hybrid polynucleotides can result from intramolecular reductive reassortment processes which utilize repeated sequences to alter a nucleotide sequence within a DNA molecule.

30 The invention provides a means for generating hybrid polynucleotides which may encode biologically active hybrid polypeptides. In one aspect, the original

polynucleotides encode biologically active polypeptides. The method of the invention produces new hybrid polypeptides by utilizing cellular processes which integrate the sequence of the original polynucleotides such that the resulting hybrid polynucleotide encodes a polypeptide demonstrating activities derived from the original biologically active polypeptides. For example, the original polynucleotides may encode a particular enzyme from different microorganisms. An enzyme encoded by a first polynucleotide from one organism may, for example, function effectively under a particular environmental condition, e.g. high salinity. An enzyme encoded by a second polynucleotide from a different organism may function effectively under a different environmental condition, such as extremely high temperatures. A hybrid polynucleotide containing sequences from the first and second original polynucleotides may encode an enzyme which exhibits characteristics of both enzymes encoded by the original polynucleotides. Thus, the enzyme encoded by the hybrid polynucleotide may function effectively under environmental conditions shared by each of the enzymes encoded by the first and second polynucleotides, e.g., high salinity and extreme temperatures.

Enzymes encoded by the original polynucleotides of the invention include, but are not limited to; oxidoreductases, transferases, hydrolases, lyases, isomerases and ligases. A hybrid polypeptide resulting from the method of the invention may exhibit specialized enzyme activity not displayed in the original enzymes. For example, following recombination and/or reductive reassortment of polynucleotides encoding hydrolase activities, the resulting hybrid polypeptide encoded by a hybrid polynucleotide can be screened for specialized hydrolase activities obtained from each of the original enzymes, i.e. the type of bond on which the hydrolase acts and the temperature at which the hydrolase functions. Thus, for example, the hydrolase may be screened to ascertain those chemical functionalities which distinguish the hybrid hydrolase from the original hydrolyases, such as: (a) amide (peptide bonds), i.e. proteases; (b) ester bonds, i.e. esterases and lipases; (c) acetals, i.e., glycosidases and, for example, the temperature, pH or salt concentration at which the hybrid polypeptide functions.

Sources of the original polynucleotides may be isolated from individual organisms ("isolates"), collections of organisms that have been grown in defined media ("enrichment cultures"), or, most preferably, uncultivated organisms ("environmental samples"). The use of a culture-independent approach to derive polynucleotides encoding novel bioactivities from environmental samples is most preferable since it allows one to access untapped resources of biodiversity.

"Environmental libraries" are generated from environmental samples and represent the collective genomes of naturally occurring organisms archived in cloning vectors that can be propagated in suitable prokaryotic hosts. Because the cloned DNA is initially extracted directly from environmental samples, the libraries are not limited to the small fraction of prokaryotes that can be grown in pure culture. Additionally, a normalization of the environmental DNA present in these samples could allow more equal representation of the DNA from all of the species present in the original sample. This can dramatically increase the efficiency of finding interesting genes from minor constituents of the sample which may be under-represented by several orders of magnitude compared to the dominant species.

For example, gene libraries generated from one or more uncultivated microorganisms are screened for an activity of interest. Potential pathways encoding bioactive molecules of interest are first captured in prokaryotic cells in the form of gene expression libraries. Polynucleotides encoding activities of interest are isolated from such libraries and introduced into a host cell. The host cell is grown under conditions which promote recombination and/or reductive reassortment creating potentially active biomolecules with novel or enhanced activities.

The microorganisms from which the polynucleotide may be prepared include prokaryotic microorganisms, such as Eubacteria and Archaeobacteria, and lower eukaryotic microorganisms such as fungi, some algae and protozoa. Polynucleotides may be isolated from environmental samples in which case the nucleic acid may be

recovered without the culturing of an organism or recovered from one or more cultured organisms. In one aspect, such microorganisms may be extremophiles, such as hyperthermophiles, psychrophiles, psychrotrophs, halophiles, barophiles and acidophiles. Polynucleotides encoding enzymes isolated from extremophilic microorganisms are particularly preferred. Such enzymes may function at temperatures above 100°C in terrestrial hot springs and deep sea thermal vents, at temperatures below 0°C in arctic waters, in the saturated salt environment of the Dead Sea, at pH values around 0 in coal deposits and geothermal sulfur-rich springs, or at pH values greater than 11 in sewage sludge. For example, several esterases and lipases cloned and expressed from extremophilic organisms show high activity throughout a wide range of temperatures and pHs.

Polynucleotides selected and isolated as hereinabove described are introduced into a suitable host cell. A suitable host cell is any cell which is capable of promoting recombination and/or reductive reassortment. The selected polynucleotides are preferably already in a vector which includes appropriate control sequences. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or preferably, the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis et al, 1986).

As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as *E. coli*, *Streptomyces*, *Salmonella typhimurium*; fungal cells, such as yeast; insect cells such as *Drosophila S2* and *Spodoptera Sf9*; animal cells such as CHO, COS or Bowes melanoma; adenoviruses; and plant cells. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

With particular references to various mammalian cell culture systems that can be employed to express recombinant protein, examples of mammalian expression systems

include the COS-7 lines of monkey kidney fibroblasts, described in "SV40-transformed simian cells support the replication of early SV40 mutants" (Gluzman, 1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

10

Host cells containing the polynucleotides of interest can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying genes. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan. The clones which are identified as having the specified enzyme activity may then be sequenced to identify the polynucleotide sequence encoding an enzyme having the enhanced activity.

In another aspect, it is envisioned the method of the present invention can be used to generate novel polynucleotides encoding biochemical pathways from one or more operons or gene clusters or portions thereof. For example, bacteria and many eukaryotes have a coordinated mechanism for regulating genes whose products are involved in related processes. The genes are clustered, in structures referred to as "gene clusters," on a single chromosome and are transcribed together under the control of a single regulatory sequence, including a single promoter which initiates transcription of the entire cluster. Thus, a gene cluster is a group of adjacent genes that are either identical or related, usually as to their function. An example of a biochemical pathway encoded by gene clusters are polyketides. Polyketides are molecules which are an extremely rich source of bioactivities, including antibiotics (such as tetracyclines and erythromycin), anti-cancer agents (daunomycin), immunosuppressants (FK506 and rapamycin), and veterinary products (monensin). Many polyketides (produced by

20
25
30

polyketide synthases) are valuable as therapeutic agents. Polyketide synthases are multifunctional enzymes that catalyze the biosynthesis of an enormous variety of carbon chains differing in length and patterns of functionality and cyclization.

Polyketide synthase genes fall into gene clusters and at least one type (designated type

- 5 I) of polyketide synthases have large size genes and enzymes, complicating genetic manipulation and *in vitro* studies of these genes/proteins.

- The ability to select and combine desired components from a library of polyketides, or fragments thereof, and postpolyketide biosynthesis genes for generation
10 of novel polyketides for study is appealing. The method of the present invention makes it possible to facilitate the production of novel polyketide synthases through intermolecular recombination.

- Preferably, gene cluster DNA can be isolated from different organisms and
15 ligated into vectors, particularly vectors containing expression regulatory sequences which can control and regulate the production of a detectable protein or protein-related array activity from the ligated gene clusters. Use of vectors which have an exceptionally large capacity for exogenous DNA introduction are particularly appropriate for use with such gene clusters and are described by way of example herein
20 to include the f-factor (or fertility factor) of *E. coli*. This f-factor of *E. coli* is a plasmid which affect high-frequency transfer of itself during conjugation and is ideal to achieve and stably propagate large DNA fragments, such as gene clusters from mixed microbial samples. Once ligated into an appropriate vector, two or more vectors containing different polyketide synthase gene clusters can be introduced into a suitable host cell.
25 Regions of partial sequence homology shared by the gene clusters will promote processes which result in sequence reorganization resulting in a hybrid gene cluster. The novel hybrid gene cluster can then be screened for enhanced activities not found in the original gene clusters.

Therefore, in a preferred embodiment, the present invention relates to a method for producing a biologically active hybrid polypeptide and screening such a polypeptide for enhanced activity by:

- 1) introducing at least a first polynucleotide in operable linkage and a second
5 polynucleotide in operable linkage, said at least first polynucleotide and second polynucleotide sharing at least one region of partial sequence homology, into a suitable host cell;
- 2) growing the host cell under conditions which promote sequence reorganization resulting in a hybrid polynucleotide in operable linkage;
- 10 3) expressing a hybrid polypeptide encoded by the hybrid polynucleotide;
- 4) screening the hybrid polypeptide under conditions which promote identification of enhanced biological activity; and
- 5) isolating the a polynucleotide encoding the hybrid polypeptide.

15 Methods for screening for various enzyme activities are known to those of skill in the art and discussed throughout the present specification. Such methods may be employed when isolating the polypeptides and polynucleotides of the present invention.

As representative examples of expression vectors which may be used there may
20 be mentioned viral particles, baculovirus, phage, plasmids, phagemids, cosmids, fosmids, bacterial artificial chromosomes, viral DNA (e.g. vaccinia, adenovirus, fowl pox virus, pseudorabies and derivatives of SV40), P1-based artificial chromosomes, yeast plasmids, yeast artificial chromosomes, and any other vectors specific for specific hosts of interest (such as bacillus, aspergillus and yeast). Thus, for example, the DNA
25 may be included in any one of a variety of expression vectors for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences. Large numbers of suitable vectors are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example; Bacterial: pQE vectors (Qiagen), pBluescript plasmids, pNH vectors, (lambda-ZAP
30 vectors (Stratagene); ptrc99a, pKK223-3, pDR540, pRIT2T (Pharmacia); Eukaryotic: pXT1, pSG5 (Stratagene), pSVK3, pBPV, pMSG, pSVLSV40 (Pharmacia). However,

any other plasmid or other vector may be used as long as they are replicable and viable in the host. Low copy number or high copy number vectors may be employed with the present invention.

- 5 A preferred type of vector for use in the present invention contains an f-factor origin replication. The f-factor (or fertility factor) in *E. coli* is a plasmid which effects high frequency transfer of itself during conjugation and less frequent transfer of the bacterial chromosome itself. A particularly preferred embodiment is to use cloning vectors, referred to as "fosmids" or bacterial artificial chromosome (BAC) vectors.
- 10 These are derived from *E. coli* f-factor which is able to stably integrate large segments of genomic DNA. When integrated with DNA from a mixed uncultured environmental sample, this makes it possible to achieve large genomic fragments in the form of a stable "environmental DNA library."
- 15 Another preferred type of vector for use in the present invention is a cosmid vector. Cosmid vectors were originally designed to clone and propagate large segments of genomic DNA. Cloning into cosmid vectors is described in detail in "Molecular Cloning: A laboratory Manual" (Sambrook et al, 1989).
- 20 The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct RNA synthesis. Particular named bacterial promoters include *lacI*, *lacZ*, T3, T7, *gpt*, λ P_R, P_L and *trp*. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the
- 25 appropriate vector and promoter is well within the level of ordinary skill in the art. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression. Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers.

30

In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

5

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence.

10 Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or
15 extracellular medium.

The cloning strategy permits expression via both vector driven and endogenous promoters; vector promotion may be important with expression of genes whose endogenous promoter will not function in *E. coli*.

20

The DNA isolated or derived from microorganisms can preferably be inserted into a vector or a plasmid prior to probing for selected DNA. Such vectors or plasmids are preferably those containing expression regulatory sequences, including promoters, enhancers and the like. Such polynucleotides can be part of a vector and/or a
25 composition and still be isolated, in that such vector or composition is not part of its natural environment. Particularly preferred phage or plasmid and methods for introduction and packaging into them are described in detail in the protocol set forth herein.

30 The selection of the cloning vector depends upon the approach taken, for example, the vector can be any cloning vector with an adequate capacity to multiply

repeated copies of a sequence, or multiple sequences that can be successfully transformed and selected in a host cell. One example of such a vector is described in "Polycos vectors: a system for packaging filamentous phage and phagemid vectors using lambda phage packaging extracts" (Alting-Mecs and Short, 1993).

- 5 Propagation/maintenance can be by an antibiotic resistance carried by the cloning vector. After a period of growth, the naturally abbreviated molecules are recovered and identified by size fractionation on a gel or column, or amplified directly. The cloning vector utilized may contain a selectable gene that is disrupted by the insertion of the lengthy construct. As reductive reassortment progresses, the number of repeated units
- 10 is reduced and the interrupted gene is again expressed and hence selection for the processed construct can be applied. The vector may be an expression/selection vector which will allow for the selection of an expressed product possessing desirable biologically properties. The insert may be positioned downstream of a functional promotor and the desirable property screened by appropriate means.

15

- In vivo* reassortment is focused on "inter-molecular" processes collectively referred to as "recombination" which in bacteria, is generally viewed as a "RecA-dependent" phenomenon. The present invention can rely on recombination processes of a host cell to recombine and re-assort sequences, or the cells' ability to mediate
- 20 reductive processes to decrease the complexity of quasi-repeated sequences in the cell by deletion. This process of "reductive reassortment" occurs by an "intra-molecular", RecA-independent process.

- Therefore, in another aspect of the present invention, novel polynucleotides can
- 25 be generated by the process of reductive reassortment. The method involves the generation of constructs containing consecutive sequences (original encoding sequences), their insertion into an appropriate vector, and their subsequent introduction into an appropriate host cell. The reassortment of the individual molecular identities occurs by combinatorial processes between the consecutive sequences in the construct
- 30 possessing regions of homology, or between quasi-repeated units. The reassortment process recombines and/or reduces the complexity and extent of the repeated sequences,

and results in the production of novel molecular species. Various treatments may be applied to enhance the rate of reassortment. These could include treatment with ultra-violet light, or DNA damaging chemicals, and/or the use of host cell lines displaying enhanced levels of "genetic instability". Thus the reassortment process may involve
5 homologous recombination or the natural property of quasi-repeated sequences to direct their own evolution.

Repeated or "quasi-repeated" sequences play a role in genetic instability. In the present invention, "quasi-repeats" are repeats that are not restricted to their original unit
10 structure. Quasi-repeated units can be presented as an array of sequences in a construct; consecutive units of similar sequences. Once ligated, the junctions between the consecutive sequences become essentially invisible and the quasi-repetitive nature of the resulting construct is now continuous at the molecular level. The deletion process the cell performs to reduce the complexity of the resulting construct operates between
15 the quasi-repeated sequences. The quasi-repeated units provide a practically limitless repertoire of templates upon which slippage events can occur. The constructs containing the quasi-repeats thus effectively provide sufficient molecular elasticity that deletion (and potentially insertion) events can occur virtually anywhere within the quasi-repetitive units.

20

When the quasi-repeated sequences are all ligated in the same orientation, for instance head to tail or vice versa, the cell cannot distinguish individual units. Consequently, the reductive process can occur throughout the sequences. In contrast, when for example, the units are presented head to head, rather than head to tail, the
25 inversion delineates the endpoints of the adjacent unit so that deletion formation will favor the loss of discrete units. Thus, it is preferable with the present method that the sequences are in the same orientation. Random orientation of quasi-repeated sequences will result in the loss of reassortment efficiency, while consistent orientation of the sequences will offer the highest efficiency. However, while having fewer of the
30 contiguous sequences in the same orientation decreases the efficiency, it may still provide sufficient elasticity for the effective recovery of novel molecules. Constructs

can be made with the quasi-repeated sequences in the same orientation to allow higher efficiency.

Sequences can be assembled in a head to tail orientation using any of a variety
5 of methods, including the following:

- a) Primers that include a poly-A head and poly-T tail which when made
single-stranded would provide orientation can be utilized. This is
accomplished by having the first few bases of the primers made from RNA
and hence easily removed RNaseH.
- 10 b) Primers that include unique restriction cleavage sites can be utilized.
Multiple sites, a battery of unique sequences, and repeated synthesis and
ligation steps would be required.
- c) The inner few bases of the primer could be thiolated and an exonuclease
used to produce properly tailed molecules.

15

The recovery of the re-assorted sequences relies on the identification of cloning
vectors with a reduced RI. The re-assorted encoding sequences can then be recovered
by amplification. The products are re-cloned and expressed. The recovery of cloning
vectors with reduced RI can be effected by:

- 20 1) The use of vectors only stably maintained when the construct is reduced in
complexity.
- 2) The physical recovery of shortened vectors by physical procedures. In this case,
the cloning vector would be recovered using standard plasmid isolation
procedures and size fractionated on either an agarose gel, or column with a low
25 molecular weight cut off utilizing standard procedures.
- 3) The recovery of vectors containing interrupted genes which can be selected
when insert size decreases.
- 4) The use of direct selection techniques with an expression vector and the
appropriate selection.

30

Encoding sequences (for example, genes) from related organisms may demonstrate a high degree of homology and encode quite diverse protein products. These types of sequences are particularly useful in the present invention as quasi-repeats. However, while the examples illustrated below demonstrate the reassortment
5 of nearly identical original encoding sequences (quasi-repeats), this process is not limited to such nearly identical repeats.

The following example demonstrates the method of the invention. Encoding nucleic acid sequences (quasi-repeats) derived from three (3) unique species are
10 depicted. Each sequence encodes a protein with a distinct set of properties. Each of the sequences differs by a single or a few base pairs at a unique position in the sequence which are designated "A", "B" and "C". The quasi-repeated sequences are separately or collectively amplified and ligated into random assemblies such that all possible permutations and combinations are available in the population of ligated molecules.
15 The number of quasi-repeat units can be controlled by the assembly conditions. The average number of quasi-repeated units in a construct is defined as the repetitive index (RI).

Once formed, the constructs may, or may not be size fractionated on an agarose
20 gel according to published protocols, inserted into a cloning vector, and transfected into an appropriate host cell. The cells are then propagated and "reductive reassortment" is effected. The rate of the reductive reassortment process may be stimulated by the introduction of DNA damage if desired. Whether the reduction in RI is mediated by deletion formation between repeated sequences by an "intra-molecular" mechanism, or
25 mediated by recombination-like events through "inter-molecular" mechanisms is immaterial. The end result is a reassortment of the molecules into all possible combinations.

Optionally, the method comprises the additional step of screening the library
30 members of the shuffled pool to identify individual shuffled library members having the ability to bind or otherwise interact (e.g., such as catalytic antibodies) with a

predetermined macromolecule, such as for example a proteinaceous receptor, peptide oligosaccharide, viron, or other predetermined compound or structure.

The displayed polypeptides, antibodies, peptidomimetic antibodies, and variable
5 region sequences that are identified from such libraries can be used for therapeutic,
diagnostic, research and related purposes (e.g., catalysts, solutes for increasing
osmolarity of an aqueous solution, and the like), and/or can be subjected to one or more
additional cycles of shuffling and/or affinity selection. The method can be modified
such that the step of selecting for a phenotypic characteristic can be other than of
10 binding affinity for a predetermined molecule (e.g., for catalytic activity, stability
oxidation resistance, drug resistance, or detectable phenotype conferred upon a host
cell).

The present invention provides a method for generating libraries of displayed
15 antibodies suitable for affinity interactions screening. The method comprises (1)
obtaining first a plurality of selected library members comprising a displayed antibody
and an associated polynucleotide encoding said displayed antibody, and obtaining said
associated polynucleotide encoding for said displayed antibody and obtaining said
associated polynucleotides or copies thereof, wherein said associated polynucleotides
20 comprise a region of substantially identical variable region framework sequence, and
(2) introducing said polynucleotides into a suitable host cell and growing the cells under
conditions which promote recombination and reductive reassortment resulting in
shuffled polynucleotides. CDR combinations comprised by the shuffled pool are not
present in the first plurality of selected library members, said shuffled pool composing a
25 library of displayed antibodies comprising CDR permutations and suitable for affinity
interaction screening. Optionally, the shuffled pool is subjected to affinity screening to
select shuffled library members which bind to a predetermined epitope (antigen) and
thereby selecting a plurality of selected shuffled library members. Further, the plurality
of selectively shuffled library members can be shuffled and screened iteratively, from 1
30 to about 1000 cycles or as desired until library members having a desired binding
affinity are obtained.

In another aspect of the invention, it is envisioned that prior to or during recombination or reassortment, polynucleotides generated by the method of the present invention can be subjected to agents or processes which promote the introduction of mutations into the original polynucleotides. The introduction of such mutations would increase the diversity of resulting hybrid polynucleotides and polypeptides encoded therefrom. The agents or processes which promote mutagenesis can include, but are not limited to: (+)-CC-1065, or a synthetic analog such as (+)-CC-1065-(N3-Adenine, see **Sun and Hurley**, 1992); an N-acetylated or deacetylated 4'-fluoro-4-aminobiphenyl adduct capable of inhibiting DNA synthesis (see, for example, **van de Poll et al**, 1992); or a N-acetylated or deacetylated 4-aminobiphenyl adduct capable of inhibiting DNA synthesis (see also, **van de Poll et al**, 1992, pp. 751-758); trivalent chromium, a trivalent chromium salt, a polycyclic aromatic hydrocarbon ("PAH") DNA adduct capable of inhibiting DNA replication, such as 7-bromomethyl-benz[*a*]anthracene ("BMA"), tris(2,3-dibromopropyl)phosphate ("Tris-BP"), 1,2-dibromo-3-chloropropane ("DBCP"), 2-bromoacrolein (2BA), benzo[*a*]pyrene-7,8-dihydrodiol-9-10-epoxide ("BPDE"), a platinum(II) halogen salt, N-hydroxy-2-amino-3-methylimidazo[4,5-*f*]quinoline ("N-hydroxy-IQ"), and N-hydroxy-2-amino-1-methyl-6-phenylimidazo[4,5-*f*]pyridine ("N-hydroxy-PhIP"). Especially preferred "means for slowing or halting PCR amplification consist of UV light (+)-CC-1065 and (+)-CC-1065-(N3-Adenine). Particularly encompassed means are DNA adducts or polynucleotides comprising the DNA adducts from the polynucleotides or polynucleotides pool, which can be released or removed by a process including heating the solution comprising the polynucleotides prior to further processing.

25

In another aspect the present invention is directed to a method of producing recombinant proteins having biological activity by treating a sample comprising double-stranded template polynucleotides encoding a wild-type protein under conditions according to the present invention which provide for the production of hybrid or reassorted polynucleotides.

30

The invention also provides the use of polynucleotide shuffling to shuffle a population of viral genes (e.g., capsid proteins, spike glycoproteins, polymerases, and proteases) or viral genomes (e.g., paramyxoviridae, orthomyxoviridae, herpesviruses, retroviruses, reoviruses and rhinoviruses). In an embodiment, the invention provides a method for shuffling sequences encoding all or portions of immunogenic viral proteins to generate novel combinations of epitopes as well as novel epitopes created by recombination; such shuffled viral proteins may comprise epitopes or combinations of epitopes as well as novel epitopes created by recombination; such shuffled viral proteins may comprise epitopes or combinations of epitopes which are likely to arise in the natural environment as a consequence of viral evolution; (e.g., such as recombination of influenza virus strains).

The invention also provides a method suitable for shuffling polynucleotide sequences for generating gene therapy vectors and replication-defective gene therapy constructs, such as may be used for human gene therapy, including but not limited to vaccination vectors for DNA-based vaccination, as well as anti-neoplastic gene therapy and other general therapy formats.

In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxy-terminal direction, in accordance with standard usage and convention. Similarly, unless specified otherwise, the left-hand end of single-stranded polynucleotide sequences is the 5' end; the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction; sequence regions on the DNA strand having the same sequence as the RNA and which are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences"; sequence regions on the DNA strand having the same sequence as the RNA and which are 3' to the 3' end of the coding RNA transcript are referred to as "downstream sequences".

Methodology

Nucleic acid shuffling is a method for *in vitro* or *in vivo* homologous recombination of pools of shorter or smaller polynucleotides to produce a polynucleotide or polynucleotides. Mixtures of related nucleic acid sequences or polynucleotides are subjected to sexual PCR to provide random polynucleotides, and
5 reassembled to yield a library or mixed population of recombinant hybrid nucleic acid molecules or polynucleotides.

In contrast to cassette mutagenesis, only shuffling and error-prone PCR allow one to mutate a pool of sequences blindly (without sequence information other than
10 primers).

The advantage of the mutagenic shuffling of this invention over error-prone PCR alone for repeated selection can best be explained with an example from antibody engineering. Consider DNA shuffling as compared with error-prone PCR (not sexual
15 PCR). The initial library of selected pooled sequences can consist of related sequences of diverse origin (i.e. antibodies from naive mRNA) or can be derived by any type of mutagenesis (including shuffling) of a single antibody gene. A collection of selected complementarity determining regions ("CDRs") is obtained after the first round of affinity selection. In the diagram the thick CDRs confer onto the antibody molecule
20 increased affinity for the antigen. Shuffling allows the free combinatorial association of all of the CDR1s with all of the CDR2s with all of the CDR3s, for example.

This method differs from error-prone PCR, in that it is an inverse chain reaction. In error-prone PCR, the number of polymerase start sites and the number of molecules
25 grows exponentially. However, the sequence of the polymerase start sites and the sequence of the molecules remains essentially the same. In contrast, in nucleic acid reassembly or shuffling of random polynucleotides the number of start sites and the number (but not size) of the random polynucleotides decreases over time. For polynucleotides derived from whole plasmids the theoretical endpoint is a single, large
30 concatemeric molecule.

Since cross-overs occur at regions of homology, recombination will primarily occur between members of the same sequence family. This discourages combinations of CDRs that are grossly incompatible (e.g., directed against different epitopes of the same antigen). It is contemplated that multiple families of sequences can be shuffled in the same reaction. Further, shuffling generally conserves the relative order, such that, for example, CDR1 will not be found in the position of CDR2.

Rare shufflants will contain a large number of the best (eg. highest affinity) CDRs and these rare shufflants may be selected based on their superior affinity.

CDRs from a pool of 100 different selected antibody sequences can be permuted in up to 1006 different ways. This large number of permutations cannot be represented in a single library of DNA sequences. Accordingly, it is contemplated that multiple cycles of DNA shuffling and selection may be required depending on the length of the sequence and the sequence diversity desired.

Error-prone PCR, in contrast, keeps all the selected CDRs in the same relative sequence, generating a much smaller mutant cloud.

The template polynucleotide which may be used in the methods of this invention may be DNA or RNA. It may be of various lengths depending on the size of the gene or shorter or smaller polynucleotide to be recombined or reassembled. Preferably, the template polynucleotide is from 50 bp to 50 kb. It is contemplated that entire vectors containing the nucleic acid encoding the protein of interest can be used in the methods of this invention, and in fact have been successfully used.

The template polynucleotide may be obtained by amplification using the PCR reaction (USPN 4,683,202 and USPN 4,683,195) or other amplification or cloning methods. However, the removal of free primers from the PCR products before subjecting them to pooling of the PCR products and sexual PCR may provide more

efficient results. Failure to adequately remove the primers from the original pool before sexual PCR can lead to a low frequency of crossover clones.

The template polynucleotide often should be double-stranded. A
5 double-stranded nucleic acid molecule is recommended to ensure that regions of the resulting single-stranded polynucleotides are complementary to each other and thus can hybridize to form a double-stranded molecule.

It is contemplated that single-stranded or double-stranded nucleic acid
10 polynucleotides having regions of identity to the template polynucleotide and regions of heterology to the template polynucleotide may be added to the template polynucleotide, at this step. It is also contemplated that two different but related polynucleotide templates can be mixed at this step.

15 The double-stranded polynucleotide template and any added double-or single-stranded polynucleotides are subjected to sexual PCR which includes slowing or halting to provide a mixture of from about 5 bp to 5 kb or more. Preferably the size of the random polynucleotides is from about 10 bp to 1000 bp, more preferably the size of the polynucleotides is from about 20 bp to 500 bp.

20 Alternatively, it is also contemplated that double-stranded nucleic acid having multiple nicks may be used in the methods of this invention. A nick is a break in one strand of the double-stranded nucleic acid. The distance between such nicks is preferably 5 bp to 5 kb, more preferably between 10 bp to 1000 bp. This can provide
25 areas of self-priming to produce shorter or smaller polynucleotides to be included with the polynucleotides resulting from random primers, for example.

The concentration of any one specific polynucleotide will not be greater than 1% by weight of the total polynucleotides, more preferably the concentration of any one
30 specific nucleic acid sequence will not be greater than 0.1% by weight of the total nucleic acid.

The number of different specific polynucleotides in the mixture will be at least about 100, preferably at least about 500, and more preferably at least about 1000.

- 5 At this step single-stranded or double-stranded polynucleotides, either synthetic or natural, may be added to the random double-stranded shorter or smaller polynucleotides in order to increase the heterogeneity of the mixture of polynucleotides.

- 10 It is also contemplated that populations of double-stranded randomly broken polynucleotides may be mixed or combined at this step with the polynucleotides from the sexual PCR process and optionally subjected to one or more additional sexual PCR cycles.

- 15 Where insertion of mutations into the template polynucleotide is desired, single-stranded or double-stranded polynucleotides having a region of identity to the template polynucleotide and a region of heterology to the template polynucleotide may be added in a 20 fold excess by weight as compared to the total nucleic acid, more preferably the single-stranded polynucleotides may be added in a 10 fold excess by weight as compared to the total nucleic acid.

- 20 Where a mixture of different but related template polynucleotides is desired, populations of polynucleotides from each of the templates may be combined at a ratio of less than about 1:100, more preferably the ratio is less than about 1:40. For example, a backcross of the wild-type polynucleotide with a population of mutated polynucleotide may be desired to eliminate neutral mutations (e.g., mutations yielding an insubstantial alteration in the phenotypic property being selected for). In such an example, the ratio of randomly provided wild-type polynucleotides which may be added to the randomly provided sexual PCR cycle hybrid polynucleotides is approximately 1:1 to about 100:1, and more preferably from 1:1 to 40:1.

30

The mixed population of random polynucleotides are denatured to form single-stranded polynucleotides and then re-annealed. Only those single-stranded polynucleotides having regions of homology with other single-stranded polynucleotides will re-anneal.

5

The random polynucleotides may be denatured by heating. One skilled in the art could determine the conditions necessary to completely denature the double-stranded nucleic acid. Preferably the temperature is from 80 °C to 100 °C, more preferably the temperature is from 90 °C to 96 °C. other methods which may be used to denature the polynucleotides include pressure (36) and pH.

10

The polynucleotides may be re-annealed by cooling. Preferably the temperature is from 20 °C to 75 °C, more preferably the temperature is from 40 °C to 65 °C. If a high frequency of crossovers is needed based on an average of only 4 consecutive bases of homology, recombination can be forced by using a low annealing temperature, although the process becomes more difficult. The degree of renaturation which occurs will depend on the degree of homology between the population of single-stranded polynucleotides.

15

Renaturation can be accelerated by the addition of polyethylene glycol ("PEG") or salt. The salt concentration is preferably from 0 mM to 200 mM, more preferably the salt concentration is from 10 mM to 100 mM. The salt may be KCl or NaCl. The concentration of PEG is preferably from 0% to 20%, more preferably from 5% to 10%.

20

The annealed polynucleotides are next incubated in the presence of a nucleic acid polymerase and dNTP's (i.e. dATP, dCTP, dGTP and dTTP). The nucleic acid polymerase may be the Klenow fragment, the Taq polymerase or any other DNA polymerase known in the art.

25

The approach to be used for the assembly depends on the minimum degree of homology that should still yield crossovers. If the areas of identity are large, Taq

30

polymerase can be used with an annealing temperature of between 45-65 °C. If the areas of identity are small, Klenow polymerase can be used with an annealing temperature of between 20-30 °C. One skilled in the art could vary the temperature of annealing to increase the number of cross-overs achieved.

5

The polymerase may be added to the random polynucleotides prior to annealing, simultaneously with annealing or after annealing.

10 The cycle of denaturation, renaturation and incubation in the presence of polymerase is referred to herein as shuffling or reassembly of the nucleic acid. This cycle is repeated for a desired number of times. Preferably the cycle is repeated from 2 to 50 times, more preferably the sequence is repeated from 10 to 40 times.

15 The resulting nucleic acid is a larger double-stranded polynucleotide of from about 50 bp to about 100 kb, preferably the larger polynucleotide is from 500 bp to 50 kb.

20 This larger polynucleotides may contain a number of copies of a polynucleotide having the same size as the template polynucleotide in tandem. This concatemeric polynucleotide is then denatured into single copies of the template polynucleotide. The result will be a population of polynucleotides of approximately the same size as the template polynucleotide. The population will be a mixed population where single or double-stranded polynucleotides having an area of identity and an area of heterology have been added to the template polynucleotide prior to shuffling.

25

These polynucleotides are then cloned into the appropriate vector and the ligation mixture used to transform bacteria.

30 It is contemplated that the single polynucleotides may be obtained from the larger concatemeric polynucleotide by amplification of the single polynucleotide prior to cloning by a variety of methods including PCR (USPN 4,683,195 and USPN 4,683,202), rather than by digestion of the concatemer.

The vector used for cloning is not critical provided that it will accept a polynucleotide of the desired size. If expression of the particular polynucleotide is desired, the cloning vehicle should further comprise transcription and translation signals
5 next to the site of insertion of the polynucleotide to allow expression of the polynucleotide in the host cell. Preferred vectors include the pUC series and the pBR series of plasmids.

The resulting bacterial population will include a number of recombinant
10 polynucleotides having random mutations. This mixed population may be tested to identify the desired recombinant polynucleotides. The method of selection will depend on the polynucleotide desired.

For example, if a polynucleotide which encodes a protein with increased binding
15 efficiency to a ligand is desired, the proteins expressed by each of the portions of the polynucleotides in the population or library may be tested for their ability to bind to the ligand by methods known in the art (i.e. panning, affinity chromatography). If a polynucleotide which encodes for a protein with increased drug resistance is desired, the proteins expressed by each of the polynucleotides in the population or library may
20 be tested for their ability to confer drug resistance to the host organism. One skilled in the art, given knowledge of the desired protein, could readily test the population to identify polynucleotides which confer the desired properties onto the protein.

It is contemplated that one skilled in the art could use a phage display system in
25 which fragments of the protein are expressed as fusion proteins on the phage surface (Pharmacia, Milwaukee WI). The recombinant DNA molecules are cloned into the phage DNA at a site which results in the transcription of a fusion protein a portion of which is encoded by the recombinant DNA molecule. The phage containing the recombinant nucleic acid molecule undergoes replication and transcription in the cell.
30 The leader sequence of the fusion protein directs the transport of the fusion protein to the tip of the phage particle. Thus the fusion protein which is partially encoded by the

recombinant DNA molecule is displayed on the phage particle for detection and selection by the methods described above.

5 It is further contemplated that a number of cycles of nucleic acid shuffling may be conducted with polynucleotides from a sub-population of the first population, which sub-population contains DNA encoding the desired recombinant protein. In this manner, proteins with even higher binding affinities or enzymatic activity could be achieved.

10 It is also contemplated that a number of cycles of nucleic acid shuffling may be conducted with a mixture of wild-type polynucleotides and a sub-population of nucleic acid from the first or subsequent rounds of nucleic acid shuffling in order to remove any silent mutations from the sub-population.

15 Any source of nucleic acid, in purified form can be utilized as the starting nucleic acid. Thus the process may employ DNA or RNA including messenger RNA, which DNA or RNA may be single or double stranded. In addition, a DNA-RNA hybrid which contains one strand of each may be utilized. The nucleic acid sequence may be of various lengths depending on the size of the nucleic acid sequence to be
20 mutated. Preferably the specific nucleic acid sequence is from 50 to 50000 base pairs. It is contemplated that entire vectors containing the nucleic acid encoding the protein of interest may be used in the methods of this invention.

The nucleic acid may be obtained from any source, for example, from plasmids
25 such a pBR322, from cloned DNA or RNA or from natural DNA or RNA from any source including bacteria, yeast, viruses and higher organisms such as plants or animals. DNA or RNA may be extracted from blood or tissue material. The template polynucleotide may be obtained by amplification using the polynucleotide chain reaction (PCR, see USPN 4,683,202 and USPN 4,683,195). Alternatively, the
30 polynucleotide may be present in a vector present in a cell and sufficient nucleic acid

may be obtained by culturing the cell and extracting the nucleic acid from the cell by methods known in the art.

Any specific nucleic acid sequence can be used to produce the population of
5 hybrids by the present process. It is only necessary that a small population of hybrid sequences of the specific nucleic acid sequence exist or be created prior to the present process.

The initial small population of the specific nucleic acid sequences having
10 mutations may be created by a number of different methods. Mutations may be created by error-prone PCR. Error-prone PCR uses low-fidelity polymerization conditions to introduce a low level of point mutations randomly over a long sequence. Alternatively, mutations can be introduced into the template polynucleotide by
15 oligonucleotide-directed mutagenesis. In oligonucleotide-directed mutagenesis, a short sequence of the polynucleotide is removed from the polynucleotide using restriction enzyme digestion and is replaced with a synthetic polynucleotide in which various bases have been altered from the original sequence. The polynucleotide sequence can also be altered by chemical mutagenesis. Chemical mutagens include, for example, sodium
20 bisulfite, nitrous acid, hydroxylamine, hydrazine or formic acid. other agents which are analogues of nucleotide precursors include nitrosoguanidine, 5-bromouracil, 2-aminopurine, or acridine. Generally, these agents are added to the PCR reaction in place of the nucleotide precursor thereby mutating the sequence. Intercalating agents such as proflavine, acriflavine, quinacrine and the like can also be used. Random
25 mutagenesis of the polynucleotide sequence can also be achieved by irradiation with X-rays or ultraviolet light. Generally, plasmid polynucleotides so mutagenized are introduced into *E. coli* and propagated as a pool or library of hybrid plasmids.

Alternatively the small mixed population of specific nucleic acids may be found
in nature in that they may consist of different alleles of the same gene or the same gene
30 from different related species (i.e., cognate genes). Alternatively, they may be related DNA sequences found within one species, for example, the immunoglobulin genes.

Once the mixed population of the specific nucleic acid sequences is generated, the polynucleotides can be used directly or inserted into an appropriate cloning vector, using techniques well-known in the art.

5

The choice of vector depends on the size of the polynucleotide sequence and the host cell to be employed in the methods of this invention. The templates of this invention may be plasmids, phages, cosmids, phagemids, viruses (e.g., retroviruses, parainfluenzavirus, herpesviruses, reoviruses, paramyxoviruses, and the like), or selected portions thereof (e.g., coat protein, spike glycoprotein, capsid protein). For example, cosmids and phagemids are preferred where the specific nucleic acid sequence to be mutated is larger because these vectors are able to stably propagate large polynucleotides.

15 If the mixed population of the specific nucleic acid sequence is cloned into a vector it can be clonally amplified by inserting each vector into a host cell and allowing the host cell to amplify the vector. This is referred to as clonal amplification because while the absolute number of nucleic acid sequences increases, the number of hybrids does not increase. Utility can be readily determined by screening expressed polypeptides.

20

The DNA shuffling method of this invention can be performed blindly on a pool of unknown sequences. By adding to the reassembly mixture oligonucleotides (with ends that are homologous to the sequences being reassembled) any sequence mixture can be incorporated at any specific position into another sequence mixture. Thus, it is contemplated that mixtures of synthetic oligonucleotides, PCR polynucleotides or even whole genes can be mixed into another sequence library at defined positions. The insertion of one sequence (mixture) is independent from the insertion of a sequence in another part of the template. Thus, the degree of recombination, the homology required, and the diversity of the library can be independently and simultaneously varied along the length of the reassembled DNA.

25

30

This approach of mixing two genes may be useful for the humanization of antibodies from murine hybridomas. The approach of mixing two genes or inserting alternative sequences into genes may be useful for any therapeutically used protein, for example, interleukin I, antibodies, tPA and growth hormone. The approach may also be useful in any nucleic acid for example, promoters or introns or 3' untranslated region or 5' untranslated regions of genes to increase expression or alter specificity of expression of proteins. The approach may also be used to mutate ribozymes or aptamers.

Shuffling requires the presence of homologous regions separating regions of diversity. Scaffold-like protein structures may be particularly suitable for shuffling. The conserved scaffold determines the overall folding by self-association, while displaying relatively unrestricted loops that mediate the specific binding. Examples of such scaffolds are the immunoglobulin beta-barrel, and the four-helix bundle which are well-known in the art. This shuffling can be used to create scaffold-like proteins with various combinations of mutated sequences for binding.

Saturation mutagenesis

In one aspect, this invention provides for the use of proprietary codon primers (containing a degenerate N,N,G/T sequence) to introduce point mutations into a polynucleotide, so as to generate a set of progeny polypeptides in which a full range of single amino acid substitutions is represented at each amino acid position. The oligos used are comprised contiguously of a first homologous sequence, a degenerate N,N,G/T sequence, and preferably but not necessarily a second homologous sequence. The downstream progeny translational products from the use of such oligos include all possible amino acid changes at each amino acid site along the polypeptide, because the degeneracy of the N,N,G/T sequence includes codons for all 20 amino acids.

In one aspect, one such degenerate oligo (comprised of one degenerate N,N,G/T cassette) is used for subjecting each original codon in a parental polynucleotide template to a full range of codon substitutions. In another aspect, at least two degenerate N,N,G/T cassettes are used – either in the same oligo or not, for subjecting at least two original codons in a parental polynucleotide template to a full range of codon substitutions. Thus, more than one N,N,G/T sequence can be contained in one oligo to introduce amino acid mutations at more than one site. This plurality of N,N,G/T sequences can be directly contiguous, or separated by one or more additional nucleotide sequence(s). In another aspect, oligos serviceable for introducing additions and deletions can be used either alone or in combination with the codons containing an N,N,G/T sequence, to introduce any combination or permutation of amino acid additions, deletions, and/or substitutions.

In a particular exemplification, it is possible to simultaneously mutagenize two or more contiguous amino acid positions using an oligo that contains contiguous N,N,G/T triplets, i.e. a degenerate $(N,N,G/T)_n$ sequence.

In another aspect, the present invention provides for the use of degenerate cassettes having less degeneracy than the N,N,G/T sequence. For example, it may be

desirable in some instances to use (e.g. in an oligo) a degenerate triplet sequence comprised of only one N, where said N can be in the first second or third position of the triplet. Any other bases including any combinations and permutations thereof can be used in the remaining two positions of the triplet. Alternatively, it may be desirable in
5 some instances to use (e.g. in an oligo) a degenerate N,N,N triplet sequence, or an N,N, G/C triplet sequence.

It is appreciated, however, that the use of a degenerate triplet (such as N,N,G/T or an N,N, G/C triplet sequence) as disclosed in the instant invention is advantageous
10 for several reasons. In one aspect, this invention provides a means to systematically and fairly easily generate the substitution of the full range of possible amino acids (for a total of 20 amino acids) into each and every amino acid position in a polypeptide. Thus, for a 100 amino acid polypeptide, the instant invention provides a way to systematically and fairly easily generate 2000 distinct species (i.e. 20 possible amino acids per position
15 X 100 amino acid positions). It is appreciated that there is provided, through the use of an oligo containing a degenerate N,N,G/T or an N,N,G/C triplet sequence, 32 individual sequences that code for 20 possible amino acids. Thus, in a reaction vessel in which a parental polynucleotide sequence is subjected to saturation mutagenesis using one such oligo, there are generated 32 distinct progeny polynucleotides encoding 20 distinct
20 polypeptides. In contrast, the use of a non-degenerate oligo in site-directed mutagenesis leads to only one progeny polypeptide product per reaction vessel.

This invention also provides for the use of nondegenerate oligos, which can optionally be used in combination with degenerate primers disclosed. It is appreciated
25 that in some situations, it is advantageous to use nondegenerate oligos to generate specific point mutations in a working polynucleotide. This provides a means to generate specific silent point mutations, point mutations leading to corresponding amino acid changes, and point mutations that cause the generation of stop codons and the corresponding expression of polypeptide fragments.

30

Thus, in a preferred embodiment of this invention, each saturation mutagenesis reaction vessel contains polynucleotides encoding at least 20 progeny polypeptide molecules such that all 20 amino acids are represented at the one specific amino acid position corresponding to the codon position mutagenized in the parental polynucleotide. The 32-fold degenerate progeny polypeptides generated from each saturation mutagenesis reaction vessel can be subjected to clonal amplification (e.g. cloned into a suitable *E. coli* host using an expression vector) and subjected to expression screening. When an individual progeny polypeptide is identified by screening to display a favorable change in property (when compared to the parental polypeptide), it can be sequenced to identify the correspondingly favorable amino acid substitution contained therein.

It is appreciated that upon mutagenizing each and every amino acid position in a parental polypeptide using saturation mutagenesis as disclosed herein, favorable amino acid changes may be identified at more than one amino acid position. One or more new progeny molecules can be generated that contain a combination of all or part of these favorable amino acid substitutions. For example, if 2 specific favorable amino acid changes are identified in each of 3 amino acid positions in a polypeptide, the permutations include 3 possibilities at each position (no change from the original amino acid, and each of two favorable changes) and 3 positions. Thus, there are $3 \times 3 \times 3$ or 27 total possibilities, including 7 that were previously examined - 6 single point mutations (i.e. 2 at each of three positions) and no change at any position.

In yet another aspect, site-saturation mutagenesis can be used together with shuffling, chimerization, recombination and other mutagenizing processes, along with screening. This invention provides for the use of any mutagenizing process(es), including saturation mutagenesis, in an iterative manner. In one exemplification, the iterative use of any mutagenizing process(es) is used in combination with screening.

Thus, in a non-limiting exemplification, this invention provides for the use of saturation mutagenesis in combination with additional mutagenization processes, such

as process where two or more related polynucleotides are introduced into a suitable host cell such that a hybrid polynucleotide is generated by recombination and reductive reassortment.

5 In addition to performing mutagenesis along the entire sequence of a gene, the instant invention provides that mutagenesis can be used to replace each of any number of bases in a polynucleotide sequence, wherein the number of bases to be mutagenized is preferably every integer from 15 to 100,000. Thus, instead of mutagenizing every position along a molecule, one can subject every a discrete number of bases (preferably
10 a subset totaling from 15 to 100,000) to mutagenesis. Preferably, a separate nucleotide is used for mutagenizing each position or group of positions along a polynucleotide sequence. A group of 3 positions to be mutagenized may be a codon. The mutations are preferably introduced using a mutagenic primer, containing a heterologous cassette, also referred to as a mutagenic cassette. Preferred cassettes can have from 1 to 500
15 bases. Each nucleotide position in such heterologous cassettes be N, A, C, G, T, A/C, A/G, A/T, C/G, C/T, G/T, C/G/T, A/G/T, A/C/T, A/C/G, or E, where E is any base that is not A, C, G, or T (E can be referred to as a designer oligo). The tables below show exemplary tri-nucleotide cassettes (there are over 3000 possibilities in addition to N,N,G/T and N,N,N and N,N,A/C).

20

In a general sense, saturation mutagenesis is comprised of mutagenizing a complete set of mutagenic cassettes (wherein each cassette is preferably 1-500 bases in length) in defined polynucleotide sequence to be mutagenized (wherein the sequence to be mutagenized is preferably from 15 to 100,000 bases in length). Thusly, a group of
25 mutations (ranging from 1 to 100 mutations) is introduced into each cassette to be mutagenized. A grouping of mutations to be introduced into one cassette can be different or the same from a second grouping of mutations to be introduced into a second cassette during the application of one round of saturation mutagenesis. Such groupings are exemplified by deletions, additions, groupings of particular codons, and
30 groupings of particular nucleotide cassettes.

Defined sequences to be mutagenized (see Fig. 20) include preferably a whole gene, pathway, cDNA, an entire open reading frame (ORF), and entire promoter, enhancer, repressor/transactivator, origin of replication, intron, operator, or any polynucleotide functional group. Generally, a preferred “defined sequences” for this purpose may be any polynucleotide that a 15 base-polynucleotide sequence, and polynucleotide sequences of lengths between 15 bases and 15,000 bases (this invention specifically names every integer in between). Considerations in choosing groupings of codons include types of amino acids encoded by a degenerate mutagenic cassette.

10 In a particularly preferred exemplification a grouping of mutations that can be introduced into a mutagenic cassette (see Tables 1-85), this invention specifically provides for degenerate codon substitutions (using degenerate oligos) that code for 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, and 20 amino acids at each position, and a library of polypeptides encoded thereby.

TABLE	Site 1	Site 2	Site 3	# of a.a.'s	NPL: POL: NEG:				
					POS:	STP			
1. N,N,G/T	N	N	G/T	20	15:	9:	2:	5:	1
2. N,N,G/C	N	N	G/C	20	15:	9:	2:	5:	1
3. N,N,G/A	N	N	G/A	14	15:	6:	2:	6:	3
4. N,N,A/C	N	N	A/C	18	14:	9:	2:	5:	2
5. N,N,A/T	N	N	A/T	18	14:	9:	2:	5:	2
6. N,N,C/T	N	N	C/T	15	14:	12:	2:	4:	0
7. N,N,N	N	N	N	20	29:	18:	4:	10:	3
8. N,N,G	N	N	G	13	8:	3:	1:	3:	1
9. N,N,A	N	N	A	12	7:	3:	1:	3:	2
10. N,N,C	N	N	C	15	7:	6:	1:	2:	0
11. N,N,T	N	N	T	15	7:	6:	1:	2:	0
12. N,N,C/G/T	N	N	C/G/T	20	22:	15:	3:	7:	1
13. N,N,A/G/T	N	N	A/G/T	20	22:	12:	3:	8:	3
14. N,N,A/C/T	N	N	A/C/T	18	21:	15:	3:	7:	2
15. N,N,A/C/G	N	N	A/C/G	20	22:	12:	3:	8:	3
16. N,A,A	N	A	A	3	0:	1:	1:	1:	1
17. N,A,C	N	A	C	4	0:	2:	1:	1:	0
18. N,A,G	N	A	G	3	0:	1:	1:	1:	1
19. N,A,T	N	A	T	4	0:	2:	1:	1:	0
20. N,C,A	N	C	A	4	2:	2:	0:	0:	0
21. N,C,C	N	C	C	4	2:	2:	0:	0:	0
22. N,C,G	N	C	G	4	2:	2:	0:	0:	0
23. N,C,T	N	C	T	4	2:	2:	0:	0:	0
24. N,G,A	N	G	A	2	1:	0:	0:	2:	1
25. N,G,C	N	G	C	4	1:	2:	0:	1:	0
26. N,G,G	N	G	G	3	2:	0:	0:	2:	0
27. N,G,T	N	G	T	4	1:	2:	0:	1:	0
28. N,T,A	N	T	A	3	4:	0:	0:	0:	0
29. N,T,C	N	T	C	4	4:	0:	0:	0:	0
30. N,T,G	N	T	G	3	4:	0:	0:	0:	0
31. N,T,T	N	T	T	4	4:	0:	0:	0:	0
32. N,A/C,A	N	A/C	A	7	2:	3:	1:	1:	1
33. N,A/G,A	N	A/G	A	5	1:	1:	1:	3:	2
34. N,A/T,A	N	A/T	A	6	4:	1:	1:	1:	1
35. N,C/G,A	N	C/G	A	6	3:	2:	0:	2:	1
36. N,C/T,A	N	C/T	A	7	6:	2:	0:	0:	0
37. N,T/G,A	N	T/G	A	5	5:	0:	0:	2:	1
38. N,C/G/T,A	N	C/G/T	A	9	7:	2:	0:	2:	1
39. N,A/G/T,A	N	A/G/T	A	8	5:	1:	1:	3:	3
40. N,A/C/T,A	N	A/C/T	A	10	6:	3:	1:	1:	1
41. N,A/C/G,A	N	A/C/G	A	9	3:	3:	1:	3:	2
42. A,N,N	A	N	N	7	4:	8:	0:	4:	0
43. C,N,N	C	N	N	5	8:	2:	0:	6:	0
44. G,N,N	G	N	N	5	12:	0:	4:	0:	0
45. T,N,N	T	N	N	6	5:	8:	0:	0:	3
46. A/C,N,N	A/C	N	N	11	12:	10:	0:	10:	0
47. A/G,N,N	A/G	N	N	12	16:	8:	4:	4:	0
48. A/T,N,N	A/T	N	N	12	9:	16:	0:	4:	3
49. C/G,N,N	C/G	N	N	10	20:	2:	4:	6:	0
50. C/T,N,N	C/T	N	N	10	13:	10:	0:	6:	3
51. G/T,N,N	G/T	N	N	11	17:	8:	4:	0:	3
52. N,A,N	N	A	N	7	0:	6:	4:	4:	2
53. N,C,N	N	C	N	4	8:	8:	0:	0:	0
54. N,G,N	N	G	N	5	5:	4:	0:	6:	1
55. N,T,N	N	T	N	5	16:	0:	0:	0:	0
56. N,A/C,N	N	A/C	N	11	8:	14:	4:	4:	2
57. N,A/G,N	N	A/G	N	12	5:	10:	4:	10:	3
58. N,A/T,N	N	A/T	N	12	16:	6:	4:	4:	2
59. N,C/G,N	N	C/G	N	8	13:	12:	0:	6:	1
60. N,C/T,N	N	C/T	N	9	24:	8:	0:	0:	0
61. N,G/T,N	N	G/T	N	10	21:	4:	0:	6:	1
62. N,A/C/G,N	N	A/C/G	N	15	13:	18:	4:	10:	3
63. N,A/C/T,N	N	A/C/T	N	16	24:	14:	4:	4:	2
64. N,A/G/T,N	N	A/G/T	N	17	21:	10:	4:	10:	3
65. N,C/G/T,N	N	C/G/T	N	13	29:	12:	0:	6:	1
66. C,C,N	C	C	N	1	4:	0:	0:	0:	0
67. G,G,N	G	G	N	1	4:	0:	0:	0:	0
68. G,C,N	G	C	N	1	4:	0:	0:	0:	0
69. G,T,N	G	T	N	1	4:	0:	0:	0:	0
70. C,G,N	C	G	N	1	0:	0:	0:	4:	0
71. C,T,N	C	T	N	1	4:	0:	0:	0:	0

TABLE	Site 1	Site 2	Site 3	# of a.a.'s	NPL: POL: NEG:				
					POS:	STP			
72. T,C,N	T	C	N	1	0:	4:	0:	0:	0
73. A,C,N	A	C	N	1	0:	4:	0:	0:	0
74. G,A,N	G	A	N	2	0:	0:	4:	0:	0
75. A,T,N	A	T	N	2	4:	0:	0:	0:	0
76. C,A,N	C	A	N	2	0:	2:	0:	2:	0
77. T,T,N	T	T	N	2	4:	0:	0:	0:	0
78. A,A,N	A	A	N	2	0:	2:	0:	2:	0
79. T,A,N	T	A	N	1	0:	2:	0:	0:	2
80. T,G,N	T	G	N	2	1:	2:	0:	0:	1
81. A,G,N	A	G	N	2	0:	2:	0:	2:	0
82. G/C,G,N	G/C	G	N	2	4:	0:	0:	4:	0
83. G/C,C,N	G/C	C	N	2	8:	0:	0:	0:	0
84. G/C,A,N	G/C	A	N	4	0:	2:	4:	2:	0
85. G/C,T,N	G/C	T	N	2	8:	0:	0:	0:	0

TABLE 1. Mutagenic Cassette: N, N, G/T

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)					
GGT	YES	GLYCINE 2	NONPOLAR (NPL) 15					
GGC	NO							
GGA	NO							
GGG	YES							
GCT	YES	ALANINE 2						
GCC	NO							
GCA	NO							
GCG	YES							
GTT	YES	VALINE 2						
GTC	NO							
GTA	NO							
GTG	YES							
TTA	NO	LEUCINE 3						
TTG	YES							
CTT	YES							
CTC	NO							
CTA	NO							
CTG	YES	ISOLEUCINE 1						
ATT	YES							
ATC	NO							
ATA	NO							
ATG	YES	METHIONINE 1						
TTT	YES	PHENYLALANINE 1						
TTC	NO							
TGG	YES	TRYPTOPHAN 1						
CCT	YES	PROLINE 2						
CCC	NO							
CCA	NO							
CCG	YES							
TCT	YES	SERINE 3	POLAR NONIONIZABLE (POL) 9					
TCC	NO							
TCA	NO							
TCG	YES							
AGT	YES							
AGC	NO	CYSTEINE 1						
TGT	YES							
TGC	NO							
AAT	YES	ASPARAGINE 1						
AAC	NO	GLUTAMINE 1						
CAA	NO							
CAG	YES	TYROSINE 1						
TAT	YES							
TAC	NO	THREONINE 2						
ACT	YES							
ACC	NO							
ACA	NO							
ACG	YES	ASPARTIC ACID 1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 2					
GAT	YES							
GAC	NO	GLUTAMIC ACID 1						
GAA	NO							
GAG	YES	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 5					
AAA	NO							
AAG	YES	ARGININE 3						
CGT	YES							
CGC	NO							
CGA	NO							
CGG	YES							
AGA	NO							
AGG	YES	HISTIDINE 1						
CAT	YES							
CAC	NO							
TAA	NO	STOP CODON 1	STOP SIGNAL (STP) 1					
TAG	YES							
TGA	NO							

TOTAL	64	32	20 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 15: 9: 2: 5: 1
-------	----	----	--------------------------------	---

TABLE 2. Mutagenic Cassette: N, N, G/C

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)					
GGT	NO	GLYCINE 2	NONPOLAR (NPL) 15					
GGC	YES							
GGA	NO							
GGG	YES							
GCT	NO	ALANINE 2						
GCC	YES							
GCA	NO							
GCG	YES							
GTT	NO	VALINE 2						
GTC	YES							
GTA	NO							
GTG	YES							
TTA	NO	LEUCINE 3						
TTG	YES							
CTT	NO							
CTC	YES							
CTA	NO							
CTG	YES							
ATT	NO	ISOLEUCINE 1						
ATC	YES							
ATA	NO							
ATG	YES	METHIONINE 1						
TTT	NO	PHENYLALANINE 1						
TTC	YES							
TGG	YES	TRYPTOPHAN 1						
CCT	NO	PROLINE 2						
CCC	YES							
CCA	NO							
CCG	YES							
TCT	NO	SERINE 3	POLAR NONIONIZABLE (POL) 9					
TCC	YES							
TCA	NO							
TCG	YES							
AGT	NO							
AGC	YES							
TGT	NO	CYSTEINE 1						
TGC	YES							
AAT	NO	ASPARAGINE 1						
AAC	YES							
CAA	NO	GLUTAMINE 1						
CAG	YES							
TAT	NO	TYROSINE 1						
TAC	YES							
ACT	NO	THREONINE 2						
ACC	YES							
ACA	NO							
ACG	YES							
GAT	NO	ASPARTIC ACID 1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 2					
GAC	YES							
GAA	NO	GLUTAMIC ACID 1						
GAG	YES							
AAA	NO	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 5					
AAG	YES							
CGT	NO	ARGININE 3						
CGC	YES							
CGA	NO							
CGG	YES							
AGA	NO							
AGG	YES							
CAT	NO	HISTIDINE 1						
CAC	YES							
TAA	NO	STOP CODON 1	STOP SIGNAL (STP) 1					
TAG	YES							
TGA	NO							
64	32	20 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 15: 9: 2: 5: 1					

TABLE 3. Mutagenic Cassette: N, N, G/A

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
GGT	NO	GLYCINE 2	NONPOLAR (NPL) 15
GGC	NO		
GGA	YES		
GGG	YES		
GCT	NO	ALANINE 2	
GCC	NO		
GCA	YES		
GCG	YES		
GTT	NO	VALINE 2	
GTC	NO		
GTA	YES		
GTG	YES		
TTA	YES	LEUCINE 4	
TTG	YES		
CTT	NO		
CTC	NO		
CTA	YES		
CTG	YES		
ATT	NO	ISOLEUCINE 1	
ATC	NO		
ATA	YES		
ATG	YES	METHIONINE 1	
TTT	NO	PHENYLALANINE 0	
TTC	NO		
TGG	YES	TRYPTOPHAN 1	
CCT	NO	PROLINE 2	
CCC	NO		
CCA	YES		
CCG	YES		
TCT	NO	SERINE 2	POLAR NONIONIZABLE (POL) 6
TCC	NO		
TCA	YES		
TCG	YES		
AGT	NO		
AGC	NO		
TGT	NO	CYSTEINE 0	
TGC	NO		
AAT	NO	ASPARAGINE 0	
AAC	NO		
CAA	YES	GLUTAMINE 2	
CAG	YES		
TAT	NO	TYROSINE 0	
TAC	NO		
ACT	NO	THREONINE 2	
ACC	NO		
ACA	YES		
ACG	YES		
GAT	NO	ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 2
GAC	NO		
GAA	YES	GLUTAMIC ACID 2	
GAG	YES		
AAA	YES	LYSINE 2	IONIZABLE: BASIC POSITIVE CHARGE (POS) 6
AAG	YES		
CGT	NO	ARGININE 4	
CGC	NO		
CGA	YES		
CGG	YES		
AGA	YES		
AGG	YES		
CAT	NO	HISTIDINE 0	
CAC	NO		
TAA	YES	STOP CODON 3	STOP SIGNAL (STP) 3
TAG	YES		
TGA	YES		

WO 00/53744

PCT/US00/06497

TOTAL	64	32	14 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 15: 6: 2: 6: 3
-------	----	----	--------------------------------	---

TABLE 4. Mutagenic Cassette: N, N, A/C

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
GGT	NO	GLYCINE 2	NONPOLAR (NPL) 14
GGC	YES		
GGA	YES		
GGG	NO		
GCT	NO	ALANINE 2	
GCC	YES		
GCA	YES		
GCG	NO		
GTT	NO	VALINE 2	
GTC	YES		
GTA	YES		
GTG	NO		
TTA	YES	LEUCINE 3	
TTG	NO		
CTT	NO		
CTC	YES		
CTA	YES		
CTG	NO		
ATT	NO	ISOLEUCINE 2	
ATC	YES		
ATA	YES		
ATG	NO	METHIONINE 0	
TTT	NO	PHENYLALANINE 1	
TTC	YES		
TGG	NO	TRYPTOPHAN 0	
CCT	NO	PROLINE 2	
CCC	YES		
CCA	YES		
CCG	NO		
TCT	NO	SERINE 3	POLAR NONIONIZABLE (POL) 9
TCC	YES		
TCA	YES		
TCG	NO		
AGT	NO	CYSTEINE 1	
AGC	YES		
TGT	NO	ASPARAGINE 1	
TGC	YES		
AAT	NO	GLUTAMINE 1	
AAC	YES		
CAA	YES	TYROSINE 1	
CAG	NO		
TAT	NO	THREONINE 2	
TAC	YES		
ACT	NO		
ACC	YES		
ACA	YES	ASPARTIC ACID 1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 2
ACG	NO		
GAT	NO		
GAC	YES		
GAA	YES	GLUTAMIC ACID 1	
GAG	NO		
AAA	YES	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 5
AAG	NO		
CGT	NO	ARGININE 3	
CGC	YES		
CGA	YES		
CGG	NO		
AGA	YES		
AGG	NO		
CAT	NO	HISTIDINE 1	
CAC	YES		
TAA	YES	STOP CODON 2	STOP SIGNAL (STP) 2
TAG	NO		
TGA	YES		
64	32	18 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 14: 9: 2: 5: 2

TABLE 5. Mutagenic Cassette: N, N, A/T

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)	
GGT	YES	GLYCINE 2	NONPOLAR (NPL) 14	
GGC	NO			
GGA	YES			
GGG	NO			
GCT	YES	ALANINE 2		
GCC	NO			
GCA	YES			
GCG	NO			
GTT	YES	VALINE 2		
GTC	NO			
GTA	YES			
GTG	NO			
TTA	YES	LEUCINE 3		
TTG	NO			
CTT	YES			
CTC	NO			
CTA	YES			
CTG	NO			
ATT	YES	ISOLEUCINE 2		
ATC	NO			
ATA	YES			
ATG	NO	METHIONINE 0		
TTT	YES	PHENYLALANINE 1		
TTC	NO	TRYPTOPHAN 0		
TGG	NO			
CCT	YES			PROLINE 2
CCC	NO			
CCA	YES			
CCG	NO			
TCT	YES	SERINE 3	POLAR NONIONIZABLE (POL) 9	
TCC	NO			
TCA	YES			
TCG	NO			
AGT	YES			
AGC	NO			
TGT	YES	CYSTEINE 1		
TGC	NO			
AAT	YES	ASPARAGINE 1		
AAC	NO			
CAA	YES	GLUTAMINE 1		
CAG	NO			
TAT	YES	TYROSINE 1		
TAC	NO			
ACT	YES	THREONINE 2		
ACC	NO			
ACA	YES			
ACG	NO			
GAT	YES	ASPARTIC ACID 1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 2	
GAC	NO	GLUTAMIC ACID 1		
GAA	YES			
GAG	NO			
AAA	YES	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 5	
AAG	NO	ARGININE 3		
CGT	YES			
CGC	NO			
CGA	YES			
CGG	NO			
AGA	YES	HISTIDINE 1		
AGG	NO			
CAT	YES			
CAC	NO			
TAA	YES	STOP CODON 2	STOP SIGNAL (STP) 2	
TAG	NO			
TGA	YES			
64	32	18 Amino Acids Are Represented	NPL: 14: POL: 9: NEG: 2: POS: 5: STP: 2	

TOTAL

TABLE 6. Mutagenic Cassette: N, N, C/T

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)							
GGT	YES	GLYCINE 2	NONPOLAR (NPL) 14							
GGC	YES									
GGA	NO									
GGG	NO									
GCT	YES	ALANINE 2								
GCC	YES									
GCA	NO									
GCG	NO									
GTT	YES	VALINE 2								
GTC	YES									
GTA	NO									
GTG	NO									
TTA	NO	LEUCINE 2								
TTG	NO									
CTT	YES									
CTC	YES									
CTA	NO									
CTG	NO									
ATT	YES	ISOLEUCINE 2								
ATC	YES									
ATA	NO									
ATG	NO	METHIONINE 0								
TTT	YES	PHENYLALANINE 2								
TTC	YES									
TGG	NO	TRYPTOPHAN 0								
CCT	YES	PROLINE 2								
CCC	YES									
CCA	NO									
CCG	NO									
TCT	YES		SERINE 4							POLAR NONIONIZABLE (POL) 12
TCC	YES									
TCA	NO									
TCG	NO									
AGT	YES									
AGC	YES									
TGT	YES	CYSTEINE 2								
TGC	YES									
AAT	YES	ASPARAGINE 2								
AAC	YES									
CAA	NO	GLUTAMINE 0								
CAG	NO									
TAT	YES	TYROSINE 2								
TAC	YES									
ACT	YES	THREONINE 2								
ACC	YES									
ACA	NO									
ACG	NO									
GAT	YES			ASPARTIC ACID 2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 2					
GAC	YES	GLUTAMIC ACID 0								
GAA	NO									
GAG	NO									
AAA	NO	LYSINE 0		IONIZABLE: BASIC POSITIVE CHARGE (POS) 4						
AAG	NO									
CGT	YES	ARGININE 2								
CGC	YES									
CGA	NO									
CGG	NO									
AGA	NO									
AGG	NO									
CAT	YES	HISTIDINE 2								
CAC	YES									
TAA	NO	STOP CODON 0	STOP SIGNAL (STP) 0							
TAG	NO									
TGA	NO									

WO 00/53744

PCT/US00/06497

TOTAL

64	32	15 Amino Acids Are Represented	NPL: POL: NEG: POS: STP
			14: 12: 2: 4: 0

TABLE 7. Mutagenic Cassette: N, N, N

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)						
GGT	YES	GLYCINE 4	NONPOLAR (NPL) 29						
GGC	YES								
GGA	YES								
GGG	YES								
GCT	YES	ALANINE 4							
GCC	YES								
GCA	YES								
GCG	YES								
GTT	YES	VALINE 4							
GTC	YES								
GTA	YES								
GTG	YES								
TTA	YES	LEUCINE 6							
TTG	YES								
CTT	YES								
CTC	YES								
CTA	YES								
CTG	YES								
ATT	YES	ISOLEUCINE 3							
ATC	YES								
ATA	YES								
ATG	YES	METHIONINE 1							
TTT	YES	PHENYLALANINE 2							
TTC	YES								
TGG	YES	TRYPTOPHAN 1							
CCT	YES	PROLINE 4							
CCC	YES								
CCA	YES								
CCG	YES								
TCT	YES	SERINE 6	POLAR NONIONIZABLE (POL) 18						
TCC	YES								
TCA	YES								
TCG	YES								
AGT	YES								
AGC	YES								
TGT	YES	CYSTEINE 2							
TGC	YES								
AAT	YES	ASPARAGINE 2							
AAC	YES								
CAA	YES	GLUTAMINE 2							
CAG	YES								
TAT	YES	TYROSINE 2							
TAC	YES								
ACT	YES	THREONINE 4							
ACC	YES								
ACA	YES								
ACG	YES								
GAT	YES	ASPARTIC ACID 2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 4						
GAC	YES								
GAA	YES	GLUTAMIC ACID 2	IONIZABLE: BASIC POSITIVE CHARGE (POS) 10						
GAG	YES								
AAA	YES	LYSINE 2							
AAG	YES								
CGT	YES	ARGININE 6							
CGC	YES								
CGA	YES								
CGG	YES								
AGA	YES								
AGG	YES								
CAT	YES	HISTIDINE 2							
CAC	YES								
TAA	YES	STOP CODON 3	STOP SIGNAL (STP) 3						
TAG	YES								
TGA	YES								
64	64	20 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 29: 18: 4: 10: 3						

TABLE 8. Mutagenic Cassette: N, N, G

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)					
GGT	NO	GLYCINE 1	NONPOLAR (NPL) 8					
GGC	NO							
GGA	NO							
GGG	YES							
GCT	NO	ALANINE 1						
GCC	NO							
GCA	NO							
GCG	YES							
GTT	NO	VALINE 1						
GTC	NO							
GTA	NO							
GTG	YES							
TTA	NO	LEUCINE 2						
TTG	YES							
CTT	NO							
CTC	NO							
CTA	NO							
CTG	YES							
ATT	NO	ISOLEUCINE 0						
ATC	NO							
ATA	NO							
ATG	YES	METHIONINE 1						
TTT	NO	PHENYLALANINE 0						
TTC	NO	TRYPTOPHAN 1						
TGG	YES							
CCT	NO	PROLINE 1						
CCC	NO							
CCA	NO							
CCG	YES							
TCT	NO	SERINE 1	POLAR NONIONIZABLE (POL) 3					
TCC	NO							
TCA	NO							
TCG	YES							
AGT	NO	CYSTEINE 0						
AGC	NO							
TGT	NO							
TGC	NO	ASPARAGINE 0						
AAT	NO							
AAC	NO	GLUTAMINE 1						
CAA	NO							
CAG	YES	TYROSINE 0						
TAT	NO							
TAC	NO	THREONINE 1						
ACT	NO							
ACC	NO							
ACA	NO							
ACG	YES	ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 1					
GAT	NO							
GAC	NO	GLUTAMIC ACID 1						
GAA	NO							
GAG	YES	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 3					
AAA	NO							
AAG	YES	ARGININE 2						
CGT	NO							
CGC	NO							
CGA	NO							
CGG	YES							
AGA	NO							
AGG	YES	HISTIDINE 0						
CAT	NO							
CAC	NO							
TAA	NO	STOP CODON 1	STOP SIGNAL (STP) 1					
TAG	YES							
TGA	NO							
64	16	13 Amino Acids Are Represented	NPL: 8: POL: 3: NEG: 1: POS: 3: STP = 1					

TABLE 9. Mutagenic Cassette: N, N, A

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
GGT	NO	GLYCINE 1	NONPOLAR (NPL) 7
GGC	NO		
GGA	YES		
GGG	NO		
GCT	NO	ALANINE 1	
GCC	NO		
GCA	YES		
GCG	NO		
GTT	NO	VALINE 1	
GTC	NO		
GTA	YES		
GTG	NO		
TTA	YES	LEUCINE 2	
TTG	NO		
CTT	NO		
CTC	NO		
CTA	YES	ISOLEUCINE 1	
CTG	NO		
ATT	NO		
ATC	NO	METHIONINE 0	
ATA	YES		
ATG	NO	PHENYLALANINE 0	
TTT	NO		
TTC	NO	TRYPTOPHAN 0	
TGG	NO		
CCT	NO	PROLINE 1	
CCC	NO		
CCA	YES		
CCG	NO		
TCT	NO	SERINE 1	POLAR NONIONIZABLE (POL) 3
TCC	NO		
TCA	YES		
TCG	NO		
AGT	NO	CYSTEINE 0	
AGC	NO		
TGT	NO	TGC 0	
TGC	NO		
AAT	NO	ASPARAGINE 0	
AAC	NO		
CAA	YES	GLUTAMINE 1	
CAG	NO		
TAT	NO	TYROSINE 0	
TAC	NO		
ACT	NO	THREONINE 1	
ACC	NO		
ACA	YES		
ACG	NO		
GAT	NO	ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 1
GAC	NO	GLUTAMIC ACID 1	
GAA	YES		
GAG	NO	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 3
AAA	YES		
AAG	NO	ARGININE 2	
CGT	NO		
CGC	NO		
CGA	YES		
CGG	NO		
AGA	YES		
AGG	NO	HISTIDINE 0	
CAT	NO		
CAC	NO		
TAA	YES	STOP CODON 2	STOP SIGNAL (STP) 2
TAG	NO		
TGA	YES		
64	16	12 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 7: 3: 1: 3: 2

TABLE 10. Mutagenic Cassette: N, N, C

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)									
GGT	NO	GLYCINE 1	NONPOLAR (NPL) 7									
GGC	YES											
GGA	NO											
GGG	NO											
GCT	NO	ALANINE 1										
GCC	YES											
GCA	NO											
GCG	NO											
GTT	NO	VALINE 1										
GTC	YES											
GTA	NO											
GTG	NO											
TTA	NO	LEUCINE 1										
TTG	NO											
CTT	NO											
CTC	YES											
CTA	NO	ISOLEUCINE 1										
CTG	NO											
ATT	NO											
ATC	YES											
ATA	NO	METHIONINE 0										
ATG	NO											
TTT	NO	PHENYLALANINE 1										
TTC	YES	TRYPTOPHAN 0										
TGG	NO											
CCT	NO	PROLINE 1										
CCC	YES											
CCA	NO											
CCG	NO											
TCT	NO	SERINE 2	POLAR NONIONIZABLE (POL) 6									
TCC	YES											
TCA	NO											
TCG	NO											
AGT	NO	CYSTEINE 1										
AGC	YES											
TGT	NO											
TGC	YES											
AAT	NO	ASPARAGINE 1										
AAC	YES											
CAA	NO	GLUTAMINE 0										
CAG	NO											
TAT	NO	TYROSINE 1										
TAC	YES											
ACT	NO	THREONINE 1										
ACC	YES											
ACA	NO											
ACG	NO											
GAT	NO	ASPARTIC ACID 1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 1									
GAC	YES											
GAA	NO	GLUTAMIC ACID 0										
GAG	NO											
AAA	NO	LYSINE 0	IONIZABLE: BASIC POSITIVE CHARGE (POS) 2									
AAG	NO											
CGT	NO	ARGININE 1										
CGC	YES											
CGA	NO											
CGG	NO											
AGA	NO											
AGG	NO											
CAT	NO	HISTIDINE 1										
CAC	YES											
TAA	NO	STOP CODON 0	STOP SIGNAL (STP) 0									
TAG	NO											
TGA	NO											
64	16	15 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 7: 6: 1: 2: 0									

TABLE 11. Mutagenic Cassette: N, N, T

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)						
GGT	YES	GLYCINE 1	NONPOLAR (NPL) 7						
GGC	NO								
GGA	NO								
GGG	NO								
GCT	YES	ALANINE 1							
GCC	NO								
GCA	NO								
GCG	NO								
GTT	YES	VALINE 1							
GTC	NO								
GTA	NO								
GTG	NO								
TTA	NO	LEUCINE 1							
TTG	NO								
CTT	YES								
CTC	NO								
CTA	NO								
CTG	NO								
ATT	YES	ISOLEUCINE 1							
ATC	NO								
ATA	NO								
ATG	NO	METHIONINE 0							
TTT	YES	PHENYLALANINE 1							
TTC	NO								
TGG	NO	TRYPTOPHAN 0							
CCT	YES	PROLINE 1							
CCC	NO								
CCA	NO								
CCG	NO								
TCT	YES	SERINE 2	POLAR NONIONIZABLE (POL) 6						
TCC	NO								
TCA	NO								
TCG	NO								
AGT	YES								
AGC	NO								
TGT	YES	CYSTEINE 1							
TGC	NO								
AAT	YES	ASPARAGINE 1							
AAC	NO								
CAA	NO	GLUTAMINE 0							
CAG	NO								
TAT	YES	TYROSINE 1							
TAC	NO								
ACT	YES	THREONINE 1							
ACC	NO								
ACA	NO								
ACG	NO								
GAT	YES	ASPARTIC ACID 1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 1						
GAC	NO								
GAA	NO	GLUTAMIC ACID 0							
GAG	NO								
AAA	NO	LYSINE 0	IONIZABLE: BASIC POSITIVE CHARGE (POS) 2						
AAG	NO								
CGT	YES	ARGININE 1							
CGC	NO								
CGA	NO								
CGG	NO								
AGA	NO								
AGG	NO								
CAT	YES	HISTIDINE 1							
CAC	NO								
TAA	NO	STOP CODON 0	STOP SIGNAL (STP) 0						
TAG	NO								
TGA	NO								
64	16	15 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 7: 6: 1: 2: 0						

TABLE 12. Mutagenic Cassette: N, N, C/G/T

CODON		Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
GGT	YES	GLYCINE 3	NONPOLAR (NPL) 22	
GGC	YES			
GGA	NO			
GGG	YES			
GCT	YES	ALANINE 3		
GCC	YES			
GCA	NO			
GCG	YES			
GTT	YES	VALINE 3		
GTC	YES			
GTA	NO			
GTG	YES			
TTA	NO	LEUCINE 4		
TTG	YES			
CTT	YES			
CTC	YES			
CTA	NO			
CTG	YES			
ATT	YES	ISOLEUCINE 2		
ATC	YES			
ATA	NO			
ATG	YES	METHIONINE 1		
TTT	YES	PHENYLALANINE 2		
TTC	YES			
TGG	YES	TRYPTOPHAN 1		
CCT	YES	PROLINE 3		
CCC	YES			
CCA	NO			
CCG	YES			
TCT	YES	SERINE 5	POLAR NONIONIZABLE (POL) 15	
TCC	YES			
TCA	NO			
TCG	YES			
AGT	YES			
AGC	YES			
TGT	YES	CYSTEINE 2		
TGC	YES			
AAT	YES	ASPARAGINE 2		
AAC	YES			
CAA	NO	GLUTAMINE 1		
CAG	YES			
TAT	YES	TYROSINE 2		
TAC	YES	THREONINE 3		
ACT	YES			
ACC	YES			
ACA	NO			
ACG	YES			
GAT	YES	ASPARTIC ACID 2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 3	
GAC	YES			
GAA	NO	GLUTAMIC ACID 1		
GAG	YES			
AAA	NO	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 7	
AAG	YES			
CGT	YES	ARGININE 4		
CGC	YES			
CGA	NO			
CGG	YES			
AGA	NO			
AGG	YES			
CAT	YES	HISTIDINE 2		
CAC	YES			
TAA	NO	STOP CODON 1	STOP SIGNAL (STP) 1	
TAG	YES			
TGA	NO			
64	48	20 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 22: 15: 3: 7: 1	

TABLE 13. Mutagenic Cassette: N, N, A/G/T

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)	
GGT	YES	GLYCINE 3	NONPOLAR (NPL) 22	
GGC	NO			
GGA	YES			
GGG	YES			
GCT	YES	ALANINE 3		
GCC	NO			
GCA	YES			
GCG	YES			
GTT	YES	VALINE 3		
GTC	NO			
GTA	YES			
GTG	YES			
TTA	YES	LEUCINE 5		
TTG	YES			
CTT	YES			
CTC	NO			
CTA	YES	ISOLEUCINE 2		
CTG	YES			
ATT	YES			
ATC	NO			
ATA	YES	METHIONINE 1		
ATG	YES			
TTT	YES	PHENYLALANINE 1		
TTC	NO			
TGG	YES	TRYPTOPHAN 1		
CCT	YES	PROLINE 3		
CCC	NO			
CCA	YES			
CCG	YES			
TCT	YES	SERINE 4	POLAR NONIONIZABLE (POL) 12	
TCC	NO			
TCA	YES			
TCG	YES			
AGT	YES	CYSTEINE 1		
AGC	NO			
TGT	YES			
TGC	NO			
AAT	YES	ASPARAGINE 1		
AAC	NO			
CAA	YES	GLUTAMINE 2		
CAG	YES			
TAT	YES	TYROSINE 1		
TAC	NO			
ACT	YES	THREONINE 3		
ACC	NO			
ACA	YES			
ACG	YES			
GAT	YES	ASPARTIC ACID 1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 3	
GAC	NO	GLUTAMIC ACID 2		
GAA	YES			
GAG	YES			
AAA	YES	LYSINE 2	IONIZABLE: BASIC POSITIVE CHARGE (POS) 8	
AAG	YES			
CGT	YES	ARGININE 5		
CGC	NO			
CGA	YES			
CGG	YES			
AGA	YES			
AGG	YES			
CAT	YES	HISTIDINE 1		
CAC	NO			
TAA	YES	STOP CODON 3	STOP SIGNAL (STP) 3	
TAG	YES			
TGA	YES			
TOTAL	64	48	20 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 22: 12: 3: 8: 3

TABLE 14. Mutagenic Cassette: N, N, A/C/T

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)					
GGT	YES	GLYCINE 3	NONPOLAR (NPL) 21					
GGC	YES							
GGA	YES							
GGG	NO							
GCT	YES	ALANINE 3						
GCC	YES							
GCA	YES							
GCG	NO							
GTT	YES	VALINE 3						
GTC	YES							
GTA	YES							
GTG	NO							
TTA	YES	LEUCINE 4						
TTG	NO							
CTT	YES							
CTC	YES							
CTA	YES							
CTG	NO							
ATT	YES	ISOLEUCINE 3						
ATC	YES							
ATA	YES							
ATG	NO	METHIONINE 0						
TTT	YES	PHENYLALANINE 2						
TTC	YES							
TGG	NO	TRYPTOPHAN 0						
CCT	YES	PROLINE 3						
CCC	YES							
CCA	YES							
CCG	NO							
TCT	YES	SERINE 5	POLAR NONIONIZABLE (POL) 15					
TCC	YES							
TCA	YES							
TCG	NO							
AGT	YES							
AGC	YES							
TGT	YES	CYSTEINE 2						
TGC	YES							
AAT	YES	ASPARAGINE 2						
AAC	YES							
CAA	YES	GLUTAMINE 1						
CAG	NO							
TAT	YES	TYROSINE 2						
TAC	YES							
ACT	YES	THREONINE 3						
ACC	YES							
ACA	YES							
ACG	NO							
GAT	YES	ASPARTIC ACID 2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 3					
GAC	YES							
GAA	YES	GLUTAMIC ACID 1						
GAG	NO							
AAA	YES	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 7					
AAG	NO							
CGT	YES	ARGININE 4						
CGC	YES							
CGA	YES							
CGG	NO							
AGA	YES							
AGG	NO							
CAT	YES	HISTIDINE 2						
CAC	YES							
TAA	YES	STOP CODON 2	STOP SIGNAL (STP) 2					
TAG	NO							
TGA	YES							
64	48	18 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 21: 15: 3: 7: 2					

TABLE 15. Mutagenic Cassette: N, N, A/C/G

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
GGT	NO	GLYCINE 3	NONPOLAR (NPL) 22
GGC	YES		
GGA	YES		
GGG	YES		
GCT	NO	ALANINE 3	
GCC	YES		
GCA	YES		
GCG	YES		
GTT	NO	VALINE 3	
GTC	YES		
GTA	YES		
GTG	YES		
TTA	YES	LEUCINE 5	
TTG	YES		
CTT	NO		
CTC	YES		
CTA	YES		
CTG	YES	ISOLEUCINE 2	
ATT	NO		
ATC	YES		
ATA	YES	METHIONINE 1	
ATG	YES		
TTT	NO	PHENYLALANINE 1	
TTC	YES	TRYPTOPHAN 1	
TGG	YES		
CCT	NO	PROLINE 3	
CCC	YES		
CCA	YES		
CCG	YES		
TCT	NO	SERINE 4	POLAR NONIONIZABLE (POL) 12
TCC	YES		
TCA	YES		
TCG	YES		
AGT	NO		
AGC	YES	CYSTEINE 1	
TGT	NO		
TGC	YES	ASPARAGINE 1	
AAT	NO		
AAC	YES	GLUTAMINE 2	
CAA	YES		
CAG	YES	TYROSINE 1	
TAT	NO		
TAC	YES	THREONINE 3	
ACT	NO		
ACC	YES		
ACA	YES		
ACG	YES	ASPARTIC ACID 1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 3
GAT	NO		
GAC	YES	GLUTAMIC ACID 2	
GAA	YES		
GAG	YES	LYSINE 2	IONIZABLE: BASIC POSITIVE CHARGE (POS) 8
AAA	YES		
AAG	YES	ARGININE 5	
CGT	NO		
CGC	YES		
CGA	YES		
CGG	YES		
AGA	YES		
AGG	YES		
CAT	NO	HISTIDINE 1	
CAC	YES		
TAA	YES	STOP CODON 3	STOP SIGNAL (STP) 3
TAG	YES		
TGA	YES		
64	48	20 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 22: 12: 3: 8: 3

TABLE 16. Mutagenic Cassette: N, A, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	0
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0	POLAR NONIONIZABLE (POL)	1
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	1		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0		
GAA	YES	GLUTAMIC ACID	1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	1
AAA	YES	LYSINE	1		
		ARGININE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	1
		HISTIDINE	0		
TAA	YES	STOP CODON	1	STOP SIGNAL (STP)	1
TOTAL	4	3 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 0: 1: 1: 1: 1	

5 TABLE 17. Mutagenic Cassette: N, A, C

CODON		Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
			GLYCINE	0	NONPOLAR (NPL)	0
			ALANINE	0		
			VALINE	0		
			LEUCINE	0		
			ISOLEUCINE	0		
			METHIONINE	0		
			PHENYLALANINE	0		
			TRYPTOPHAN	0		
			PROLINE	0		
			SERINE	0		
			CYSTEINE	0	POLAR NONIONIZABLE (POL)	2
AAC	YES	ASPARAGINE	1			
		GLUTAMINE	0			
TAC	YES	TYROSINE	1			
		THREONINE	0			
GAC	YES	ASPARTIC ACID	1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	1	
		GLUTAMIC ACID	0			
			LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	1
			ARGININE	0		
CAC	YES	HISTIDINE	1			
			STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL		4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 0: 2: 1: 1: 0	

TABLE 18. Mutagenic Cassette: N, A, G

TABLE 16. Mutagenic Cassette: P ₁ , P ₂ , P ₃						
CODON		Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
			GLYCINE	0	NONPOLAR (NPL)	0
			ALANINE	0		
			VALINE	0		
			LEUCINE	0		
			ISOLEUCINE	0		
			METHIONINE	0		
			PHENYLALANINE	0		
			TRYPTOPHAN	0		
			PROLINE	0		
			SERINE	0	POLAR NONIONIZABLE (POL)	1
			CYSTEINE	0		
			ASPARAGINE	0		
CAG	YES		GLUTAMINE	1		
			TYROSINE	0		
			THREONINE	0		
			ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	1
GAG	YES		GLUTAMIC ACID	1		
AAG	YES		LYSINE	1	IONIZABLE: BASIC POSITIVE CHARGE (POS)	1
			ARGININE	0		
			HISTIDINE	0		
TAG	YES		STOP CODON	1	STOP SIGNAL (STP)	1
TOTAL		4	3 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 0: 1: 1: 1: 1	

5 TABLE 19. Mutagenic Cassette: N, A, T

TABLE 19: Mutagenic Cassette: A, T, C, G						
CODON		Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
			GLYCINE	0	NONPOLAR (NPL)	0
			ALANINE	0		
			VALINE	0		
			LEUCINE	0		
			ISOLEUCINE	0		
			METHIONINE	0		
			PHENYLALANINE	0		
			TRYPTOPHAN	0		
			PROLINE	0		
			SERINE	0	POLAR NONIONIZABLE (POL)	2
			CYSTEINE	0		
AAT	YES		ASPARAGINE	1		
			GLUTAMINE	0		
TAT	YES		TYROSINE	1		
			THREONINE	0		
GAT	YES		ASPARTIC ACID	1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	1
			GLUTAMIC ACID	0		
			LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	1
			ARGININE	0		
CAT	YES		HISTIDINE	1		
			STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL		4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 0: 2: 1: 1: 0	

TABLE 20. Mutagenic Cassette: N, C, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	2
GCA	YES	ALANINE	1		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCA	YES	PROLINE	1		
TCA	YES	SERINE	1	POLAR NONIONIZABLE (POL)	2
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACA	YES	THREONINE	1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 2: 2: 0: 0: 0	

5 TABLE 21. Mutagenic Cassette: N, C, C

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	2
GCC	YES	ALANINE	1		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCC	YES	PROLINE	1		
TCC	YES	SERINE	1	POLAR NONIONIZABLE (POL)	2
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACC	YES	THREONINE	1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 2: 2: 0: 0: 0	

TABLE 22. Mutagenic Cassette: N, C, G

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	2
GCG	YES	ALANINE	1		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCG	YES	PROLINE	1	POLAR NONIONIZABLE (POL)	2
TCG	YES	SERINE	1		
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACG	YES	THREONINE	1		
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 2: 2: 0: 0: 0	

5 TABLE 23. Mutagenic Cassette: N, C, T

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	2
GCT	YES	ALANINE	1		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCT	YES	PROLINE	1	POLAR NONIONIZABLE (POL)	2
TCT	YES	SERINE	1		
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACT	YES	THREONINE	1		
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 2: 2: 0: 0: 0	

TABLE 24. Mutagenic Cassette: N, G, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGA	YES	GLYCINE	1	NONPOLAR (NPL)	1
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	2
CGA	YES	ARGININE	2		
AGA	YES	HISTIDINE	0		
TGA	YES	STOP CODON	1	STOP SIGNAL (STP)	1
TOTAL	4	2 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 1: 0: 0: 2: 1	

5 TABLE 25. Mutagenic Cassette: N, G, C

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGC	YES	GLYCINE	1	NONPOLAR (NPL)	1
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
AGC	YES	SERINE	1	POLAR NONIONIZABLE (POL)	2
TGC	YES	CYSTEINE	1		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	1
CGC	YES	ARGININE	1		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 1: 2: 0: 1: 0	

TABLE 26. Mutagenic Cassette: N, G, G

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGG	YES	GLYCINE	1	NONPOLAR (NPL)	2
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
TGG	YES	TRYPTOPHAN	1		
		PROLINE	0	POLAR NONIONIZABLE (POL)	0
		SERINE	0		
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	2
CGG	YES	LYSINE	0		
AGG	YES	ARGININE	2		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	3 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 2: 0: 0: 2: 0	

5 TABLE 27. Mutagenic Cassette: N, G, T

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	1	NONPOLAR (NPL)	1
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
AGT	YES	SERINE	1	POLAR NONIONIZABLE (POL)	2
TGT	YES	CYSTEINE	1		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	1
CGT	YES	ARGININE	1		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 1: 2: 0: 1: 0	

TABLE 28. Mutagenic Cassette: N, T, A

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
		GLYCINE 0	NONPOLAR (NPL) 4
		ALANINE 0	
GTA	YES	VALINE 1	
TTA	YES	LEUCINE 2	
CTA	YES		
ATA	YES	ISOLEUCINE 1	
		METHIONINE 0	
		PHENYLALANINE 0	
		TRYPTOPHAN 0	
		PROLINE 0	
		SERINE 0	POLAR NONIONIZABLE (POL) 0
		CYSTEINE 0	
		ASPARAGINE 0	
		GLUTAMINE 0	
		TYROSINE 0	
		THREONINE 0	
		ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 0
		GLUTAMIC ACID 0	
		LYSINE 0	IONIZABLE: BASIC POSITIVE CHARGE (POS) 0
		ARGININE 0	
		HISTIDINE 0	
		STOP CODON 0	STOP SIGNAL (STP) 0
TOTAL	4	3 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 4: 0: 0: 0: 0

5 TABLE 29. Mutagenic Cassette: N, T, C

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
		GLYCINE 0	NONPOLAR (NPL) 4
		ALANINE 0	
GTC	YES	VALINE 1	
CTC	YES	LEUCINE 1	
ATC	YES	ISOLEUCINE 1	
		METHIONINE 0	
TTC	YES	PHENYLALANINE 1	
		TRYPTOPHAN 0	
		PROLINE 0	
		SERINE 0	POLAR NONIONIZABLE (POL) 0
		CYSTEINE 0	
		ASPARAGINE 0	
		GLUTAMINE 0	
		TYROSINE 0	
		THREONINE 0	
		ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 0
		GLUTAMIC ACID 0	
		LYSINE 0	IONIZABLE: BASIC POSITIVE CHARGE (POS) 0
		ARGININE 0	
		HISTIDINE 0	
		STOP CODON 0	STOP SIGNAL (STP) 0
TOTAL	4	4 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 4: 0: 0: 0: 0

TABLE 30. Mutagenic Cassette: N, T, G

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	4
		ALANINE	0		
GTG	YES	VALINE	1		
TTG	YES	LEUCINE	2		
CTG	YES				
		ISOLEUCINE	0		
ATG	YES	METHIONINE	1		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	3 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 4: 0: 0: 0: 0	

5 TABLE 31. Mutagenic Cassette: N, T, T

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	4
		ALANINE	0		
GTT	YES	VALINE	1		
CTT	YES	LEUCINE	1		
ATT	YES	ISOLEUCINE	1		
		METHIONINE	0		
TTT	YES	PHENYLALANINE	1		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 4: 0: 0: 0: 0	

TABLE 32. Mutagenic Cassette: N, A/C, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	2
GCA	YES	ALANINE	1		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCA	YES	PROLINE	1	POLAR NONIONIZABLE (POL)	3
TCA	YES	SERINE	1		
		CYSTEINE	0		
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	1		
		TYROSINE	0		
ACA	YES	THREONINE	1		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	1
GAA	YES	GLUTAMIC ACID	1		
AAA	YES	LYSINE	1	IONIZABLE: BASIC POSITIVE CHARGE (POS)	1
		ARGININE	0		
		HISTIDINE	0		
TAA	YES	STOP CODON	1	STOP SIGNAL (STP)	1
TOTAL	8	7 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 2: 3: 1: 1: 1	

5 TABLE 33. Mutagenic Cassette: N, A/G, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGA	YES	GLYCINE	1	NONPOLAR (NPL)	1
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	1
		CYSTEINE	0		
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	1		
		TYROSINE	0		
		THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	1
		ASPARTIC ACID	0		
GAA	YES	GLUTAMIC ACID	1	IONIZABLE: BASIC POSITIVE CHARGE (POS)	3
AAA	YES	LYSINE	1		
CGA	YES	ARGININE	2		
AGA	YES	HISTIDINE	0		
		STOP CODON	2		
TAA	YES	STOP CODON	2	STOP SIGNAL (STP)	2
TGA	YES				
TOTAL	8	5 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 1: 1: 1: 3: 2	

TABLE 34. Mutagenic Cassette: N, A/T, A

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
		GLYCINE 0	NONPOLAR (NPL) 4
		ALANINE 0	
GTA	YES	VALINE 1	
TTA	YES	LEUCINE 2	
CTA	YES		
ATA	YES	ISOLEUCINE 1	
		METHIONINE 0	
		PHENYLALANINE 0	
		TRYPTOPHAN 0	
		PROLINE 0	
		SERINE 0	POLAR NONIONIZABLE (POL) 1
		CYSTEINE 0	
		ASPARAGINE 0	
CAA	YES	GLUTAMINE 1	
		TYROSINE 0	
		THREONINE 0	
		ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 1
GAA	YES	GLUTAMIC ACID 1	
AAA	YES	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 1
		ARGININE 0	
		HISTIDINE 0	
TAA	YES	STOP CODON 1	STOP SIGNAL (STP) 1
TOTAL	8	6 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 4: 1: 1: 1: 1

5 TABLE 35. Mutagenic Cassette: N, C/G, A

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
GGA	YES	GLYCINE 1	NONPOLAR (NPL) 3
GCA	YES	ALANINE 1	
		VALINE 0	
		LEUCINE 0	
		ISOLEUCINE 0	
		METHIONINE 0	
		PHENYLALANINE 0	
		TRYPTOPHAN 0	
CCA	YES	PROLINE 1	
TCA	YES	SERINE 1	POLAR NONIONIZABLE (POL) 2
		CYSTEINE 0	
		ASPARAGINE 0	
		GLUTAMINE 0	
		TYROSINE 0	
ACA	YES	THREONINE 1	
		ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 0
		GLUTAMIC ACID 0	
		LYSINE 0	IONIZABLE: BASIC POSITIVE CHARGE (POS) 2
CGA	YES	ARGININE 2	
AGA	YES	HISTIDINE 0	
TGA	YES	STOP CODON 1	STOP SIGNAL (STP) 1
TOTAL	8	6 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 3: 2: 0: 2: 1

TABLE 36. Mutagenic Cassette: N, C/T, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	6
GCA	YES	ALANINE	1		
GTA	YES	VALINE	1		
TTA	YES	LEUCINE	2		
CTA	YES				
ATA	YES	ISOLEUCINE	1		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCA	YES	PROLINE	1		
TCA	YES	SERINE	1	POLAR NONIONIZABLE (POL)	2
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACA	YES	THREONINE	1	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	8	7 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 6: 2: 0: 0: 0	

5 TABLE 37. Mutagenic Cassette: N, T/G, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGA	YES	GLYCINE	1	NONPOLAR (NPL)	5
		ALANINE	0		
GTA	YES	VALINE	1		
TTA	YES	LEUCINE	2		
CTA	YES				
ATA	YES	ISOLEUCINE	1		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	2
CGA	YES	LYSINE	0		
AGA	YES	ARGININE	2		
		HISTIDINE	0	STOP SIGNAL (STP)	1
TGA	YES	STOP CODON	1		
TOTAL	8	5 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 5: 0: 0: 2: 1	

TABLE 38. Mutagenic Cassette: N, C/G/T, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGA	YES	GLYCINE	1	NONPOLAR (NPL)	7
GCA	YES	ALANINE	1		
GTA	YES	VALINE	1		
TTA	YES	LEUCINE	2		
CTA	YES				
ATA	YES	ISOLEUCINE	1		
		METHIONINE	0		
		PHENYLALANINE	0	POLAR NONIONIZABLE (POL)	2
		TRYPTOPHAN	0		
CCA	YES	PROLINE	1		
TCA	YES	SERINE	1		
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
ACA	YES	THREONINE	1		
		ASPARTIC ACID	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	2
		GLUTAMIC ACID	0		
		LYSINE	0		
CGA	YES	ARGININE	2		
AGA	YES			STOP SIGNAL (STP)	1
		HISTIDINE	0		
TGA	YES	STOP CODON	1	STOP SIGNAL (STP)	1
TOTAL	12	9 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 7: 2: 0: 2: 1	

5 TABLE 39. Mutagenic Cassette: N, A/G/T, A

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGA	YES	GLYCINE	1	NONPOLAR (NPL)	5
		ALANINE	0		
GTA	YES	VALINE	1		
TTA	YES	LEUCINE	2		
CTA	YES				
ATA	YES	ISOLEUCINE	1		
		METHIONINE	0		
		PHENYLALANINE	0	POLAR NONIONIZABLE (POL)	1
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0		
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	1		
		TYROSINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	1
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	3
GAA	YES	GLUTAMIC ACID	1		
AAA	YES	LYSINE	1		
CGA	YES	ARGININE	2		
AGA	YES			STOP SIGNAL (STP)	2
		HISTIDINE	0		
TAA	YES	STOP CODON	2	STOP SIGNAL (STP)	2
TGA	YES				
TOTAL	12	8 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 5: 1: 1: 3: 2	

TABLE 40. Mutagenic Cassette: N, A/C/T, A

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
		GLYCINE 0	NONPOLAR (NPL) 6
GCA	YES	ALANINE 1	
GTA	YES	VALINE 1	
TTA	YES	LEUCINE 2	
CTA	YES		
ATA	YES	ISOLEUCINE 1	
		METHIONINE 0	
		PHENYLALANINE 0	
		TRYPTOPHAN 0	
CCA	YES	PROLINE 1	
TCA	YES	SERINE 1	POLAR NONIONIZABLE (POL) 3
		CYSTEINE 0	
		ASPARAGINE 0	
CAA	YES	GLUTAMINE 1	
		TYROSINE 0	
ACA	YES	THREONINE 1	
		ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 1
GAA	YES	GLUTAMIC ACID 1	
AAA	YES	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 1
		ARGININE 0	
		HISTIDINE 0	
TAA	YES	STOP CODON 1	STOP SIGNAL (STP) 1
TOTAL	12	10 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 6: 3: 1: 1: 1

5 TABLE 41. Mutagenic Cassette: N, A/C/G, A

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
GGA	YES	GLYCINE 1	NONPOLAR (NPL) 3
GCA	YES	ALANINE 1	
		VALINE 0	
		LEUCINE 0	
		ISOLEUCINE 0	
		METHIONINE 0	
		PHENYLALANINE 0	
		TRYPTOPHAN 0	
CCA	YES	PROLINE 1	
TCA	YES	SERINE 1	POLAR NONIONIZABLE (POL) 3
		CYSTEINE 0	
		ASPARAGINE 0	
CAA	YES	GLUTAMINE 1	
		TYROSINE 0	
ACA	YES	THREONINE 1	
		ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 1
GAA	YES	GLUTAMIC ACID 1	
AAA	YES	LYSINE 1	IONIZABLE: BASIC POSITIVE CHARGE (POS) 3
CGA	YES	ARGININE 2	
AGA	YES		
		HISTIDINE 0	
TAA	YES	STOP CODON 2	STOP SIGNAL (STP) 2
TGA	YES		
TOTAL	12	9 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 3: 3: 1: 3: 2

TABLE 42. Mutagenic Cassette: A, N, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	4
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
ATT	YES	ISOLEUCINE	3		
ATC	YES				
ATA	YES				
ATG	YES				
		METHIONINE	1		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
AGT	YES	SERINE	2	POLAR NONIONIZABLE (POL)	8
AGC	YES				
		CYSTEINE	0		
AAT	YES	ASPARAGINE	2		
AAC	YES				
		GLUTAMINE	0		
		TYROSINE	0		
ACT	YES	THREONINE	4		
ACC	YES				
ACA	YES				
ACG	YES				
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
AAA	YES	LYSINE	2	IONIZABLE: BASIC POSITIVE CHARGE (POS)	4
AAG	YES				
AGA	YES	ARGININE	2		
AGG	YES				
		HISTIDINE	0	STOP SIGNAL (STP)	0
		STOP CODON	0		
TOTAL	16	7 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 4: 8: 0: 4: 0	

5 TABLE 43. Mutagenic Cassette: C, N, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	8
		ALANINE	0		
		VALINE	0		
CTT	YES	LEUCINE	4		
CTC	YES				
CTA	YES				
CTG	YES				
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
		SERINE	0	POLAR NONIONIZABLE (POL)	2
		CYSTEINE	0		
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	2		
CAG	YES				
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	6
CGT	YES	ARGININE	4		
CGC	YES				
CGA	YES				
CGG	YES				
CAT	YES	HISTIDINE	2		
CAC	YES				

WO 00/53744

PCT/US00/06497

TOTAL		STOP CODON 0	STOP SIGNAL (STP) 0
	16	5 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 8: 2: 0: 6: 0

TABLE 44. Mutagenic Cassette: G, N, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	12
GGC	YES				
GGA	YES				
GGG	YES				
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
		LEUCINE	0	POLAR NONIONIZABLE (POL)	0
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0		
		ASPARAGINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
GAT	YES	ASPARTIC ACID	2		
GAC	YES				
GAA	YES	GLUTAMIC ACID	2		
GAG	YES			IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		LYSINE	0		
		ARGININE	0		
		HISTIDINE	0	STOP SIGNAL (STP)	0
		STOP CODON	0		
TOTAL	16	5 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 12: 0: 4: 0: 0	

5 TABLE 45. Mutagenic Cassette: T, N, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	5
		ALANINE	0		
		VALINE	0		
TTA	YES	LEUCINE	2		
TTG	YES				
		ISOLEUCINE	0		
		METHIONINE	0		
TTT	YES	PHENYLALANINE	2	POLAR NONIONIZABLE (POL)	8
TTC	YES				
TGG	YES	TRYPTOPHAN	1		
		PROLINE	0		
TCT	YES	SERINE	4		
TCC	YES				
TCA	YES				
TCG	YES				
TGT	YES	CYSTEINE	2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
TGC	YES				
		ASPARAGINE	0		
		GLUTAMINE	0		
TAT	YES	TYROSINE	2	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
TAC	YES				
		THREONINE	0		
		ASPARTIC ACID	0	STOP SIGNAL (STP)	3
		GLUTAMIC ACID	0		
		LYSINE	0	STOP SIGNAL (STP)	3
		ARGININE	0		
		HISTIDINE	0		
TAA	YES	STOP CODON	3	STOP SIGNAL (STP)	3
TAG	YES				

WO 00/53744

PCT/US00/06497

TOTAL	TGA	YES						
		16	6 Amino Acids Are Represented	NPL:	POL:	NEG:	POS:	STP =
				5:	8:	0:	0:	3

TABLE 46. Mutagenic Cassette: A/C, N, N

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
		GLYCINE 0	NONPOLAR (NPL) 12
		ALANINE 0	
		VALINE 0	
CTT	YES	LEUCINE 4	
CTC	YES		
CTA	YES		
CTG	YES		
ATT	YES	ISOLEUCINE 3	
ATC	YES		
ATA	YES		
ATG	YES		
		METHIONINE 1	
		PHENYLALANINE 0	
		TRYPTOPHAN 0	
CCT	YES	PROLINE 4	
CCC	YES		
CCA	YES		
CCG	YES		
AGT	YES	SERINE 2	POLAR NONIONIZABLE (POL) 10
AGC	YES		
		CYSTEINE 0	
AAT	YES	ASPARAGINE 2	
AAC	YES		
CAA	YES	GLUTAMINE 2	
CAG	YES		
		TYROSINE 0	
ACT	YES	THREONINE 4	
ACC	YES		
ACA	YES		
ACG	YES		
		ASPARTIC ACID 0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 0
		GLUTAMIC ACID 0	
AAA	YES	LYSINE 2	IONIZABLE: BASIC POSITIVE CHARGE (POS) 10
AAG	YES		
CGT	YES	ARGININE 6	
CGC	YES		
CGA	YES		
CGG	YES		
AGA	YES		
AGG	YES		
CAT	YES	HISTIDINE 2	
CAC	YES		
		STOP CODON 0	STOP SIGNAL (STP) 0
	32	11 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 12: 10: 0: 10: 0

TABLE 47. Mutagenic Cassette: A/G, N, N

CODON	Represented	AMINO ACID (Frequency)	CATEGORY (Frequency)
GGT	YES	GLYCINE 4	NONPOLAR (NPL) 16
GGC	YES		
GGA	YES		
GGG	YES		
GCT	YES	ALANINE 4	
GCC	YES		
GCA	YES		
GCG	YES		
GTT	YES	VALINE 4	
GTC	YES		
GTA	YES		
GTG	YES		
		LEUCINE 0	
ATT	YES	ISOLEUCINE 3	
ATC	YES		
ATA	YES		
ATG	YES	METHIONINE 1	
		PHENYLALANINE 0	
		TRYPTOPHAN 0	
		PROLINE 0	
AGT	YES	SERINE 2	POLAR NONIONIZABLE (POL) 8
AGC	YES		
		CYSTEINE 0	
AAT	YES	ASPARAGINE 2	
AAC	YES		
		GLUTAMINE 0	
		TYROSINE 0	
ACT	YES	THREONINE 4	
ACC	YES		
ACA	YES		
ACG	YES		
GAT	YES	ASPARTIC ACID 2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 4
GAC	YES		
GAA	YES	GLUTAMIC ACID 2	
GAG	YES		
AAA	YES	LYSINE 2	IONIZABLE: BASIC POSITIVE CHARGE (POS) 4
AAG	YES		
AGA	YES	ARGININE 2	
AGG	YES		
		HISTIDINE 0	
		STOP CODON 0	STOP SIGNAL (STP) 0
	32	12 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 16: 8: 4: 4: 0

TABLE 48. Mutagenic Cassette: A/T, N, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)		
		GLYCINE	0	NONPOLAR (NPL)	9		
		ALANINE	0				
		VALINE	0				
TTA	YES	LEUCINE	2				
TTG	YES						
ATT	YES	ISOLEUCINE	3				
ATC	YES						
ATA	YES						
ATG	YES						
TTT	YES	METHIONINE	1				
TTC	YES	PHENYLALANINE	2				
TGG	YES	TRYPTOPHAN	1				
		PROLINE	0				
TCT	YES	SERINE	6	POLAR NONIONIZABLE (POL)	16		
TCC	YES						
TCA	YES						
TCG	YES						
AGT	YES						
AGC	YES						
TGT	YES	CYSTEINE	2				
TGC	YES						
AAT	YES	ASPARAGINE	2				
AAC	YES						
		GLUTAMINE	0				
TAT	YES	TYROSINE	2				
TAC	YES						
ACT	YES	THREONINE	4				
ACC	YES						
ACA	YES						
ACG	YES						
		ASPARTIC ACID	0			IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0				
AAA	YES	LYSINE	2	IONIZABLE: BASIC POSITIVE CHARGE (POS)	4		
AAG	YES						
AGA	YES					ARGININE	2
AGG	YES						
		HISTIDINE	0	STOP SIGNAL (STP)	3		
TAA	YES	STOP CODON	3				
TAG	YES						
TGA	YES						
	32	12 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 9: 16: 0: 4: 3			

TABLE 49. Mutagenic Cassette: C/G, N, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	20
GGC	YES				
GGA	YES				
GGG	YES				
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
CTT	YES	LEUCINE	4		
CTC	YES				
CTA	YES				
CTG	YES				
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
		SERINE	0	POLAR NONIONIZABLE (POL)	2
		CYSTEINE	0		
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	2		
CAG	YES				
		TYROSINE	0		
		THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
GAT	YES	ASPARTIC ACID	2		
GAC	YES				
GAA	YES	GLUTAMIC ACID	2		
GAG	YES			IONIZABLE: BASIC POSITIVE CHARGE (POS)	6
		LYSINE	0		
CGT	YES	ARGININE	4		
CGC	YES				
CGA	YES				
CGG	YES				
CAT	YES	HISTIDINE	2		
CAC	YES				
		STOP CODON	0	STOP SIGNAL (STP)	0
	32	10 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 20: 2: 4: 6: 0	

TABLE 50. Mutagenic Cassette: C/T, N, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	13
		ALANINE	0		
		VALINE	0		
TTA	YES	LEUCINE	6		
TTG	YES				
CTT	YES				
CTC	YES				
CTA	YES				
CTG	YES				
		ISOLEUCINE	0		
		METHIONINE	0		
TTT	YES	PHENYLALANINE	2		
TTC	YES				
TGG	YES	TRYPTOPHAN	1		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
TCT	YES	SERINE	4	POLAR NONIONIZABLE (POL)	10
TCC	YES				
TCA	YES				
TCG	YES				
TGT	YES	CYSTEINE	2		
TGC	YES				
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	2		
CAG	YES				
TAT	YES	TYROSINE	2		
TAC	YES				
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	6
CGT	YES	ARGININE	4		
CGC	YES				
CGA	YES				
CGG	YES				
CAT	YES	HISTIDINE	2		
CAC	YES				
TAA	YES	STOP CODON	3	STOP SIGNAL (STP)	3
TAG	YES				
TGA	YES				
	32	10 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 13: 10: 0: 6: 3	

5

10

TABLE 51. Mutagenic Cassette: G/T, N, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	17
GGC	YES				
GGA	YES				
GGG	YES				
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
TTA	YES	LEUCINE	2		
TTG	YES				
		ISOLEUCINE	0		
		METHIONINE	0		
TTT	YES	PHENYLALANINE	2		
TTC	YES				
TGG	YES	TRYPTOPHAN	1		
		PROLINE	0		
TCT	YES	SERINE	4	POLAR NONIONIZABLE (POL)	8
TCC	YES				
TCA	YES				
TCG	YES				
TGT	YES	CYSTEINE	2		
TGC	YES				
		ASPARAGINE	0		
		GLUTAMINE	0		
TAT	YES	TYROSINE	2		
TAC	YES				
		THREONINE	0		
GAT	YES	ASPARTIC ACID	2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
GAC	YES				
GAA	YES	GLUTAMIC ACID	2		
GAG	YES				
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
TAA	YES	STOP CODON	3	STOP SIGNAL (STP)	3
TAG	YES				
TGA	YES				
	32	11 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 17: 8: 4: 0: 3	

TABLE 52. Mutagenic Cassette: N, A, N

TABLE 52. Mutagenic Cassette A, 1974					
CODON		Represented	AMINO ACID	(Frequency)	CATEGORY (Frequency)
			GLYCINE	0	NONPOLAR (NPL) 0
			ALANINE	0	
			VALINE	0	
			LEUCINE	0	
			ISOLEUCINE	0	
			METHIONINE	0	
			PHENYLALANINE	0	
			TRYPTOPHAN	0	
			PROLINE	0	
			SERINE	0	POLAR NONIONIZABLE (POL) 6
			CYSTEINE	0	
AAT	YES		ASPARAGINE	2	
AAC	YES				
CAA	YES		GLUTAMINE	2	
CAG	YES				
TAT	YES		TYROSINE	2	
TAC	YES				
			THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 4
GAT	YES		ASPARTIC ACID	2	
GAC	YES				
GAA	YES		GLUTAMIC ACID	2	
GAG	YES				IONIZABLE: BASIC POSITIVE CHARGE (POS) 4
AAA	YES		LYSINE	2	
AAG	YES				
			ARGININE	0	
CAT	YES		HISTIDINE	2	STOP SIGNAL (STP) 2
CAC	YES				
TAA	YES		STOP CODON	2	
TAG	YES				
TOTAL		16	7 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 0: 6: 4: 4: 2

TABLE 53. Mutagenic Cassette: N, C, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	8
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
TCT	YES	SERINE	4	POLAR NONIONIZABLE (POL)	8
TCC	YES				
TCA	YES				
TCG	YES				
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACT	YES	THREONINE	4		
ACC	YES				
ACA	YES				
ACG	YES				
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0

WO 00/53744

PCT/US00/06497

TOTAL

	16	4 Amino Acids Are Represented	NPL: POL: NEG: POS: STP =
			8: 8: 0: 0: 0

TABLE 54. Mutagenic Cassette: N, G, N

CODON		Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES		GLYCINE	4	NONPOLAR (NPL)	5
GGC	YES					
GGA	YES					
GGG	YES					
			ALANINE	0		
			VALINE	0		
			LEUCINE	0		
			ISOLEUCINE	0		
			METHIONINE	0		
			PHENYLALANINE	0		
TGG	YES		TRYPTOPHAN	1		
			PROLINE	0		
AGT	YES		SERINE	2	POLAR NONIONIZABLE (POL)	4
AGC	YES					
TGT	YES		CYSTEINE	2		
TGC	YES					
			ASPARAGINE	0		
			GLUTAMINE	0		
			TYROSINE	0		
			THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
			ASPARTIC ACID	0		
			GLUTAMIC ACID	0		
			LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	6
CGT	YES		ARGININE	6		
CGC	YES					
CGA	YES					
CGG	YES					
AGA	YES					
AGG	YES					
			HISTIDINE	0	STOP SIGNAL (STP)	1
TGA	YES		STOP CODON	1		
TOTAL		16	5 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 5: 4: 0: 6: 1	

TABLE 55. Mutagenic Cassette: N, T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)		
		GLYCINE	0	NONPOLAR (NPL)	16		
		ALANINE	0				
GTT	YES	VALINE	4				
GTC	YES						
GTA	YES						
GTG	YES						
TTA	YES	LEUCINE	6				
TTG	YES						
CTT	YES						
CTC	YES						
CTA	YES						
CTG	YES						
ATT	YES	ISOLEUCINE	3				
ATC	YES						
ATA	YES						
ATG	YES	METHIONINE	1				
TTT	YES	PHENYLALANINE	2				
TTC	YES						
		TRYPTOPHAN	0	POLAR NONIONIZABLE (POL)	0		
		PROLINE	0				
		SERINE	0				
		CYSTEINE	0				
		ASPARAGINE	0				
		GLUTAMINE	0				
		TYROSINE	0				
		THREONINE	0				
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0		
		GLUTAMIC ACID	0				
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0		
		ARGININE	0				
		HISTIDINE	0				
		STOP CODON	0	STOP SIGNAL (STP)	0		

WO 00/53744

PCT/US00/06497

TOTAL

	16	5 Amino Acids Are Represented	NPL: POL: NEG: POS: STP =
			16: 0: 0: 0: 0

TABLE 56. Mutagenic Cassette: N, A/C, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	8
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCT	YES	PROLINE	4		
CCC	YES			POLAR NONIONIZABLE (POL)	14
CCA	YES				
CCG	YES				
TCT	YES	SERINE	4		
TCC	YES				
TCA	YES				
TCG	YES				
		CYSTEINE	0		
AAT	YES	ASPARAGINE	2		
AAC	YES				
CAA	YES	GLUTAMINE	2		
CAG	YES				
TAT	YES	TYROSINE	2		
TAC	YES				
ACT	YES	THREONINE	4		
ACC	YES			IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
ACA	YES				
ACG	YES				
GAT	YES	ASPARTIC ACID	2		
GAC	YES			IONIZABLE: BASIC POSITIVE CHARGE (POS)	4
GAA	YES	GLUTAMIC ACID	2		
GAG	YES				
AAA	YES	LYSINE	2		
AAG	YES			STOP SIGNAL (STP)	2
		ARGININE	0		
CAT	YES	HISTIDINE	2		
CAC	YES				
TAA	YES	STOP CODON	2	TOTAL	NPL: 8: POL: 14: NEG: 4: POS: 4: STP = 2
TAG	YES				
	32	11 Amino Acids Are Represented			

TABLE 57. Mutagenic Cassette: N, A/G, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)		
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	5		
GGC	YES						
GGA	YES						
GGG	YES						
		ALANINE	0				
		VALINE	0				
		LEUCINE	0				
		ISOLEUCINE	0				
		METHIONINE	0				
		PHENYLALANINE	0				
TGG	YES	TRYPTOPHAN	1				
		PROLINE	0				
AGT	YES	SERINE	2	POLAR NONIONIZABLE (POL)	10		
AGC	YES	CYSTEINE	2				
TGT	YES						
TGC	YES	ASPARAGINE	2				
AAT	YES						
AAC	YES	GLUTAMINE	2				
CAA	YES						
CAG	YES	TYROSINE	2				
TAT	YES						
TAC	YES	THREONINE	0				
GAT	YES	ASPARTIC ACID	2			IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
GAC	YES	GLUTAMIC ACID	2				
GAA	YES						
GAG	YES	LYSINE	2	IONIZABLE: BASIC POSITIVE CHARGE (POS)	10		
AAA	YES						
AAG	YES						
CGT	YES					ARGININE	6
CGC	YES						
CGA	YES						
CGG	YES						
AGA	YES						
AGG	YES						
CAT	YES						
CAC	YES						
TAA	YES	STOP CODON	3	STOP SIGNAL (STP)	3		
TAG	YES						
TGA	YES						
	32	12 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 5: 10: 4: 10: 3			

TABLE 58. Mutagenic Cassette: N, A/T, N

CODON	Represented	CATEGORY	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	16
		ALANINE	0		
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
TTA	YES	LEUCINE	6		
TTG	YES				
CTT	YES				
CTC	YES				
CTA	YES				
CTG	YES				
ATT	YES	ISOLEUCINE	3		
ATC	YES				
ATA	YES				
ATG	YES	METHIONINE	1		
TTT	YES	PHENYLALANINE	2		
TTC	YES				
		TRYPTOPHAN	0	POLAR NONIONIZABLE (POL)	6
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0		
AAT	YES	ASPARAGINE	2		
AAC	YES				
CAA	YES	GLUTAMINE	2		
CAG	YES				
TAT	YES	TYROSINE	2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
TAC	YES				
		THREONINE	0		
GAT	YES	ASPARTIC ACID	2		
GAC	YES			IONIZABLE: BASIC POSITIVE CHARGE (POS)	4
GAA	YES	GLUTAMIC ACID	2		
GAG	YES				
AAA	YES	LYSINE	2		
AAG	YES			STOP SIGNAL (STP)	2
		ARGININE	0		
CAT	YES	HISTIDINE	2		
CAC	YES				
TAA	YES	STOP CODON	2	NPL: POL: NEG: POS: STP =	16: 6: 4: 4: 2
TAG	YES				
TOTAL	32	12 Amino Acids Are Represented			

TABLE 59. Mutagenic Cassette: N, C/G, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	13
GGC	YES				
GGA	YES				
GGG	YES				
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
TGG	YES	TRYPTOPHAN	1		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
TCT	YES	SERINE	6	POLAR NONIONIZABLE (POL)	12
TCC	YES				
TCA	YES				
TCG	YES				
AGT	YES	CYSTEINE	2		
AGC	YES				
TGT	YES				
TGC	YES				
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACT	YES	THREONINE	4		
ACC	YES				
ACA	YES				
ACG	YES				
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	6
CGT	YES	ARGININE	6		
CGC	YES				
CGA	YES				
CGG	YES				
AGA	YES				
AGG	YES				
		HISTIDINE	0		
TGA	YES	STOP CODON	1	STOP SIGNAL (STP)	1
	32	8 Amino Acids Are Represented		NPL: 13: POL: 12: NEG: 0: POS: 6: STP = 1	

TABLE 60. Mutagenic Cassette: N, C/T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	24
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
TTA	YES	LEUCINE	6		
TTG	YES				
CTT	YES				
CTC	YES				
CTA	YES				
CTG	YES				
ATT	YES	ISOLEUCINE	3		
ATC	YES				
ATA	YES				
ATG	YES	METHIONINE	1		
TTT	YES	PHENYLALANINE	2		
TTC	YES				
		TRYPTOPHAN	0	POLAR NONIONIZABLE (POL)	8
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
TCT	YES	SERINE	4		
TCC	YES				
TCA	YES				
TCG	YES				
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACT	YES	THREONINE	4		
ACC	YES				
ACA	YES				
ACG	YES				
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	32	9 Amino Acids Are Represented	NPL: POL: NEG: POS: STP = 24: 8: 0: 0: 0		

TABLE 61. Mutagenic Cassette: N, G/T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	21
GGC	YES				
GGA	YES				
GGG	YES				
		ALANINE	0		
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
TTA	YES	LEUCINE	6		
TTG	YES				
CTT	YES				
CTC	YES				
CTA	YES				
CTG	YES				
ATT	YES	ISOLEUCINE	3		
ATC	YES				
ATA	YES				
ATG	YES	METHIONINE	1		
TTT	YES	PHENYLALANINE	2		
TTC	YES				
TGG	YES	TRYPTOPHAN	1		
		PROLINE	0		
AGT	YES	SERINE	2	POLAR NONIONIZABLE (POL)	4
AGC	YES				
TGT	YES	CYSTEINE	2		
TGC	YES				
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	6
CGT	YES	ARGININE	6		
CGC	YES				
CGA	YES				
CGG	YES				
AGA	YES				
AGG	YES				
		HISTIDINE	0		
TGA	YES	STOP CODON	1	STOP SIGNAL (STP)	1
TOTAL	32	10 Amino Acids Are Represented		NPL: 21: POL: 4: NEG: 0: POS: 6: STP = 1	

TABLE 62. Mutagenic Cassette: N, A/C/G, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	13
GGC	YES				
GGA	YES				
GGG	YES				
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
TGG	YES	TRYPTOPHAN	1		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
TCT	YES	SERINE	6	POLAR NONIONIZABLE (POL)	18
TCC	YES				
TCA	YES				
TCG	YES				
AGT	YES	CYSTEINE	2		
AGC	YES				
TGT	YES				
TGC	YES				
AAT	YES	ASPARAGINE	2		
AAC	YES	GLUTAMINE	2		
CAA	YES				
CAG	YES	TYROSINE	2		
TAT	YES				
TAC	YES				
ACT	YES	THREONINE	4		
ACC	YES				
ACA	YES				
ACG	YES				
GAT	YES	ASPARTIC ACID	2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
GAC	YES	GLUTAMIC ACID	2		
GAA	YES				
GAG	YES				
AAA	YES	LYSINE	2	IONIZABLE: BASIC POSITIVE CHARGE (POS)	10
AAG	YES	ARGININE	6		
CGT	YES				
CGC	YES				
CGA	YES				
CGG	YES				
AGA	YES	HISTIDINE	2		
AGG	YES				
CAT	YES				
CAC	YES				
TAA	YES	STOP CODON	3	STOP SIGNAL (STP)	3
TAG	YES				
TGA	YES				
	48	15 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 13: 18: 4: 10: 3	

TABLE 63. Mutagenic Cassette: N, A/C/T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	24
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
TTA	YES	LEUCINE	6		
TTG	YES				
CTT	YES				
CTC	YES				
CTA	YES				
CTG	YES				
ATT	YES	ISOLEUCINE	3		
ATC	YES				
ATA	YES				
ATG	YES	METHIONINE	1		
TTT	YES	PHENYLALANINE	2		
TTC	YES				
		TRYPTOPHAN	0		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
TCT	YES	SERINE	4	POLAR NONIONIZABLE (POL)	14
TCC	YES				
TCA	YES				
TCG	YES				
		CYSTEINE	0		
AAT	YES	ASPARAGINE	2		
AAC	YES				
CAA	YES	GLUTAMINE	2		
CAG	YES				
TAT	YES	TYROSINE	2		
TAC	YES				
ACT	YES	THREONINE	4		
ACC	YES				
ACA	YES				
ACG	YES				
GAT	YES	ASPARTIC ACID	2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
GAC	YES				
GAA	YES	GLUTAMIC ACID	2		
GAG	YES				
AAA	YES	LYSINE	2	IONIZABLE: BASIC POSITIVE CHARGE (POS)	4
AAG	YES				
		ARGININE	0		
CAT	YES	HISTIDINE	2		
CAC	YES			STOP SIGNAL (STP)	2
TAA	YES	STOP CODON	2		
TAG	YES				
TOTAL	48	16 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 24: 14: 4: 4: 2	

TABLE 64. Mutagenic Cassette: N, A/G/T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)		
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	21		
GGC	YES						
GGA	YES						
GGG	YES						
		ALANINE	0				
GTT	YES	VALINE	4				
GTC	YES						
GTA	YES						
GTG	YES						
TTA	YES	LEUCINE	6				
TIG	YES						
CTT	YES						
CTC	YES						
CTA	YES						
CTG	YES						
ATT	YES	ISOLEUCINE	3				
ATC	YES						
ATA	YES						
ATG	YES	METHIONINE	1				
TTT	YES	PHENYLALANINE	2				
TTC	YES						
TGG	YES	TRYPTOPHAN	1				
		PROLINE	0				
AGT	YES	SERINE	2	POLAR NONIONIZABLE (POL)	10		
AGC	YES						
TGT	YES	CYSTEINE	2				
TGC	YES						
AAT	YES	ASPARAGINE	2				
AAC	YES						
CAA	YES	GLUTAMINE	2				
CAG	YES						
TAT	YES	TYROSINE	2				
TAC	YES						
		THREONINE	0				
GAT	YES	ASPARTIC ACID	2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4		
GAC	YES						
GAA	YES	GLUTAMIC ACID	2				
GAG	YES						
AAA	YES	LYSINE	2	IONIZABLE: BASIC POSITIVE CHARGE (POS)	10		
AAG	YES						
CGT	YES	ARGININE	6				
CGC	YES						
CGA	YES						
CGG	YES						
AGA	YES						
AGG	YES						
CAT	YES	HISTIDINE	2				
CAC	YES						
TAA	YES	STOP CODON	3	STOP SIGNAL (STP)	3		
TAG	YES						
TGA	YES						
	48	17 Amino Acids Are Represented		NPL: 21: POL: 10: NEG: 4: POS: 10: STP = 3			

TABLE 65. Mutagenic Cassette: N, C/G/T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	29
GGC	YES				
GGA	YES				
GGG	YES				
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
TTA	YES	LEUCINE	6		
TTG	YES				
CTT	YES				
CTC	YES				
CTA	YES				
CTG	YES				
ATT	YES	ISOLEUCINE	3		
ATC	YES				
ATA	YES				
ATG	YES	METHIONINE	1		
TTT	YES	PHENYLALANINE	2		
TTC	YES				
TGG	YES	TRYPTOPHAN	1		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
TCT	YES	SERINE	6	POLAR NONIONIZABLE (POL)	12
TCC	YES				
TCA	YES				
TCG	YES				
AGT	YES				
AGC	YES				
TGT	YES	CYSTEINE	2		
TGC	YES				
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
ACT	YES	THREONINE	4		
ACC	YES				
ACA	YES				
ACG	YES				
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	6
CGT	YES	ARGININE	6		
CGC	YES				
CGA	YES				
CGG	YES				
AGA	YES				
AGG	YES				
		HISTIDINE	0		
TGA	YES	STOP CODON	1	STOP SIGNAL (STP)	1
	48	13 Amino Acids Are Represented		NPL: 29: POL: 12: NEG: 0: POS: 6: STP = 1	

TABLE 66. Mutagenic Cassette: C, C, N

TABLE 30. Mutagenesis Calculations					
CODON		Represented	AMINO ACID	(Frequency)	CATEGORY (Frequency)
			GLYCINE	0	NONPOLAR (NPL) 4
			ALANINE	0	
			VALINE	0	
			LEUCINE	0	
			ISOLEUCINE	0	
			METHIONINE	0	
			PHENYLALANINE	0	
			TRYPTOPHAN	0	
CCT	YES		PROLINE	4	
CCC	YES				
CCA	YES				
CCG	YES				
			SERINE	0	POLAR NONIONIZABLE (POL) 0
			CYSTEINE	0	
			ASPARAGINE	0	
			GLUTAMINE	0	
			TYROSINE	0	
			THREONINE	0	
			ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 0
			GLUTAMIC ACID	0	
			LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS) 0
			ARGININE	0	
			HISTIDINE	0	
			STOP CODON	0	STOP SIGNAL (STP) 0
TOTAL		4	1 Amino Acid Is Represented		NPL: 4: POL: 0: NEG: 0: POS: 0: STP = 0

5 TABLE 67. Mutagenic Cassette: G, G, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	4
GGC	YES				
GGA	YES				
GGG	YES				
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	0	4	1 Amino Acid Is Represented		NPL: 4: POL: 0: NEG: 0: POS: 0: STP = 0

TABLE 68. Mutagenic Cassette: G, C, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	4
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0	POLAR NONIONIZABLE (POL)	0
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		GLUTAMIC ACID	0		
		LYSINE	0		
		ARGININE	0	STOP SIGNAL (STP)	0
		HISTIDINE	0		
		STOP CODON	0	NPL: POL: NEG: POS: STP = 4: 0: 0: 0: 0	
TOTAL	4	1 Amino Acid Is Represented			

5 TABLE 69. Mutagenic Cassette: G, T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	4
		ALANINE	0		
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0	POLAR NONIONIZABLE (POL)	0
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		TYROSINE	0		
		THREONINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0		
		LYSINE	0	STOP SIGNAL (STP)	0
		ARGININE	0		
		HISTIDINE	0	NPL: POL: NEG: POS: STP = 4: 0: 0: 0: 0	
TOTAL	4	1 Amino Acid Is Represented			

TABLE 70. Mutagenic Cassette: C, G, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)		
		GLYCINE	0	NONPOLAR (NPL)	0		
		ALANINE	0				
		VALINE	0				
		LEUCINE	0				
		ISOLEUCINE	0				
		METHIONINE	0				
		PHENYLALANINE	0				
		TRYPTOPHAN	0				
		PROLINE	0				
		SERINE	0	POLAR NONIONIZABLE (POL)	0		
		CYSTEINE	0				
		ASPARAGINE	0				
		GLUTAMINE	0				
		TYROSINE	0				
		THREONINE	0				
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0		
		GLUTAMIC ACID	0				
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	4		
CGT	YES	ARGININE	4				
CGC	YES						
CGA	YES						
CGG	YES						
		HISTIDINE	0	STOP SIGNAL (STP)	0		
		STOP CODON	0				
TOTAL	4	1 Amino Acid Is Represented		NPL: POL: NEG: POS: STP = 0: 0: 0: 4: 0			

5 TABLE 71. Mutagenic Cassette: C, T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	4
		ALANINE	0		
		VALINE	0		
CTT	YES	LEUCINE	4		
CTC	YES				
CTA	YES				
CTG	YES				
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	1 Amino Acid Is Represented		NPL: POL: NEG: POS: STP = 4: 0: 0: 0: 0	

TABLE 72. Mutagenic Cassette: T, C, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	0
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
TCT	YES	SERINE	4	POLAR NONIONIZABLE (POL)	4
TCC	YES				
TCA	YES				
TCG	YES				
		CYSTEINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		LYSINE	0		
		ARGININE	0		
		HISTIDINE	0	STOP SIGNAL (STP)	0
		STOP CODON	0		
TOTAL	4	1 Amino Acid Is Represented		NPL: 0: POL: 4: NEG: 0: POS: 0: STP = 0	

5 TABLE 73. Mutagenic Cassette: A, C, N

CODON		Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
			GLYCINE	0	NONPOLAR (NPL)	0
			ALANINE	0		
			VALINE	0		
			LEUCINE	0		
			ISOLEUCINE	0		
			METHIONINE	0		
			PHENYLALANINE	0		
			TRYPTOPHAN	0		
			PROLINE	0		
			SERINE	0	POLAR NONIONIZABLE (POL)	4
			CYSTEINE	0		
			ASPARAGINE	0		
			GLUTAMINE	0		
			TYROSINE	0		
ACT	YES		THREONINE	4		
ACC	YES					
ACA	YES					
ACG	YES					
			ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
			GLUTAMIC ACID	0		
			LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
			ARGININE	0		
			HISTIDINE	0		
			STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL		4	1 Amino Acid Is Represented		NPL: 0: POL: 4: NEG: 0: POS: 0: STP = 0	

TABLE 74. Mutagenic Cassette: G, A, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	0
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
GAT	YES	ASPARTIC ACID	2	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
GAC	YES				
GAA	YES	GLUTAMIC ACID	2		
GAG	YES				
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	2 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 0: 0: 4: 0: 0	

5 TABLE 75. Mutagenic Cassette: A, T, N

TABLE 15. Mutations Observed					
CODON		Represented	AMINO ACID	(Frequency)	CATEGORY (Frequency)
			GLYCINE	0	NONPOLAR (NPL) 4
			ALANINE	0	
			VALINE	0	
			LEUCINE	0	
ATT	YES		ISOLEUCINE	3	
ATC	YES				
ATA	YES				
ATG	YES		METHIONINE	1	
			PHENYLALANINE	0	
			TRYPTOPHAN	0	
			PROLINE	0	
			SERINE	0	POLAR NONIONIZABLE (POL) 0
			CYSTEINE	0	
			ASPARAGINE	0	
			GLUTAMINE	0	
			TYROSINE	0	
			THREONINE	0	
			ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 0
			GLUTAMIC ACID	0	
			LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS) 0
			ARGININE	0	
			HISTIDINE	0	
			STOP CODON	0	STOP SIGNAL (STP) 0
TOTAL		4	2 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 4: 0: 0: 0: 0

TABLE 76. Mutagenic Cassette: C, A, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	0
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	2
		CYSTEINE	0		
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	2		
CAG	YES				
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ARGININE	0		
CAT	YES	HISTIDINE	2		
CAC	YES				
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	2 Amino Acids Are Represented		NPL: 0: POL: 2: NEG: 0: POS: 2: STP = 0	

5 TABLE 77. Mutagenic Cassette: T, T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	4
		ALANINE	0		
		VALINE	0		
TTA	YES	LEUCINE	2		
TTG	YES				
		ISOLEUCINE	0		
		METHIONINE	0		
TTT	YES	PHENYLALANINE	2		
TTC	YES				
		TRYPTOPHAN	0	POLAR NONIONIZABLE (POL)	0
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	4	2 Amino Acids Are Represented		NPL: 4: POL: 0: NEG: 0: POS: 0: STP = 0	

TABLE 78. Mutagenic Cassette: A, A, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	0
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0	POLAR NONIONIZABLE (POL)	2
AAT	YES	ASPARAGINE	2		
AAC	YES				
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
AAA	YES	LYSINE	2	IONIZABLE: BASIC POSITIVE CHARGE (POS)	2
AAG	YES				
		ARGININE	0		
		HISTIDINE	0	STOP SIGNAL (STP)	0
		STOP CODON	0		
TOTAL	4	2 Amino Acids Are Represented		NPL: POL: NEG: POS: STP =	0: 2: 0: 2: 0

5 TABLE 79. Mutagenic Cassette: T, A, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	0
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0	POLAR NONIONIZABLE (POL)	2
		ASPARAGINE	0		
		GLUTAMINE	0		
TAT	YES	TYROSINE	2		
TAC	YES				
		THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		LYSINE	0		
		ARGININE	0		
		HISTIDINE	0	STOP SIGNAL (STP)	2
TAA	YES	STOP CODON	2		
TAG	YES			STOP SIGNAL (STP)	2
TOTAL	4	1 Amino Acid Is Represented		NPL: POL: NEG: POS: STP =	0: 2: 0: 0: 2

TABLE 80. Mutagenic Cassette: T, G, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	1
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
TGG	YES	TRYPTOPHAN	1		
		PROLINE	0	POLAR NONIONIZABLE (POL)	2
		SERINE	0		
TGT	YES	CYSTEINE	2		
TGC	YES				
		TYROSINE	0		
		THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		LYSINE	0		
		ARGININE	0		
		HISTIDINE	0		
TGA	YES	STOP CODON	1	STOP SIGNAL (STP)	1
TOTAL	4	2 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 1: 2: 0: 0: 1	

5 TABLE 81. Mutagenic Cassette: A, G, N

TABLE 01: Mutagenesis Results, 3/11/11					
CODON		Represented	AMINO ACID	(Frequency)	CATEGORY (Frequency)
			GLYCINE	0	NONPOLAR (NPL) 0
			ALANINE	0	
			VALINE	0	
			LEUCINE	0	
			ISOLEUCINE	0	
			METHIONINE	0	
			PHENYLALANINE	0	
			TRYPTOPHAN	0	
			PROLINE	0	POLAR NONIONIZABLE (POL) 2
AGT	YES		SERINE	2	
AGC	YES				
			CYSTEINE	0	
			ASPARAGINE	0	
			GLUTAMINE	0	
			TYROSINE	0	
			THREONINE	0	
			ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG) 0
			GLUTAMIC ACID	0	
			LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS) 2
AGA	YES		ARGININE	2	
AGG	YES				
			HISTIDINE	0	STOP SIGNAL (STP) 0
			STOP CODON	0	
TOTAL		4	2 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 0: 2: 0: 2: 0

TABLE 82. Mutagenic Cassette: G/C, G, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
GGT	YES	GLYCINE	4	NONPOLAR (NPL)	4
GGC	YES				
GGA	YES				
GGG	YES				
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	4
CGT	YES	ARGININE	4		
CGC	YES				
CGA	YES				
CGG	YES				
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0
TOTAL	8	2 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 4: 0: 0: 4: 0	

TABLE 83. Mutagenic Cassette: G/C, C, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	8
GCT	YES	ALANINE	4		
GCC	YES				
GCA	YES				
GCG	YES				
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
CCT	YES	PROLINE	4		
CCC	YES				
CCA	YES				
CCG	YES				
		SERINE	0	POLAR NONIONIZABLE (POL)	0
		CYSTEINE	0		
		ASPARAGINE	0		
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0		
		ASPARTIC ACID	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMIC ACID	0		
		LYSINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	STOP SIGNAL (STP)	0

WO 00/53744

PCT/US00/06497

TOTAL

	8	2 Amino Acids Are Represented	NPL:	POL:	NEG:	POS:	STP =
			8:	0:	0:	0:	0

TABLE 84. Mutagenic Cassette: G/C, A, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	0
		ALANINE	0		
		VALINE	0		
		LEUCINE	0		
		ISOLEUCINE	0		
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0	POLAR NONIONIZABLE (POL)	2
		ASPARAGINE	0		
CAA	YES	GLUTAMINE	2		
CAG	YES				
		TYROSINE	0		
		THREONINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	4
GAT	YES	ASPARTIC ACID	2		
GAC	YES				
GAA	YES	GLUTAMIC ACID	2		
GAG	YES			IONIZABLE: BASIC POSITIVE CHARGE (POS)	2
		LYSINE	0		
		ARGININE	0		
CAT	YES	HISTIDINE	2		
CAC	YES			STOP SIGNAL (STP)	0
		STOP CODON	0		
TOTAL	8	4 Amino Acids Are Represented		NPL: POL: NEG: POS: STP = 0: 2: 4: 2: 0	

TABLE 85. Mutagenic Cassette: G/C, T, N

CODON	Represented	AMINO ACID	(Frequency)	CATEGORY	(Frequency)
		GLYCINE	0	NONPOLAR (NPL)	8
		ALANINE	0		
GTT	YES	VALINE	4		
GTC	YES				
GTA	YES				
GTG	YES				
CTT	YES	LEUCINE	4		
CTC	YES				
CTA	YES				
CTG	YES				
		ISOLEUCINE	0	POLAR NONIONIZABLE (POL)	0
		METHIONINE	0		
		PHENYLALANINE	0		
		TRYPTOPHAN	0		
		PROLINE	0		
		SERINE	0		
		CYSTEINE	0		
		ASPARAGINE	0	IONIZABLE: ACIDIC NEGATIVE CHARGE (NEG)	0
		GLUTAMINE	0		
		TYROSINE	0		
		THREONINE	0	IONIZABLE: BASIC POSITIVE CHARGE (POS)	0
		ASPARTIC ACID	0		
		GLUTAMIC ACID	0		
		LYSINE	0	STOP SIGNAL (STP)	0
		ARGININE	0		
		HISTIDINE	0		
		STOP CODON	0	NPL: POL: NEG: POS: STP =	8: 0: 0: 0: 0
TOTAL	8	2 Amino Acids Are Represented			

2.11.2.3. EXONUCLEASE-MEDIATED REASSEMBLY

In a particular embodiment, this invention provides for a method for shuffling, assembling, reassembling, recombining, &/or concatenating at least two
5 polynucleotides to form a progeny polynucleotide (e.g. a chimeric progeny polynucleotide that can be expressed to produce a polypeptide or a gene pathway). In a particular embodiment, a double stranded polynucleotide end (e.g. two single stranded sequences hybridized to each other as hybridization partners) is treated with an
10 exonuclease to liberate nucleotides from one of the two strands, leaving the remaining strand free of its original partner so that, if desired, the remaining strand may be used to achieve hybridization to another partner.

In a particular aspect, a double stranded polynucleotide end (that may be part of - or connected to - a polynucleotide or a nonpolynucleotide sequence) is subjected to a
15 source of exonuclease activity. Serviceable sources of exonuclease activity may be an enzyme with 3' exonuclease activity, an enzyme with 5' exonuclease activity, an enzyme with both 3' exonuclease activity and 5' exonuclease activity, and any combination thereof. An exonuclease can be used to liberate nucleotides from one or both ends of a linear double stranded polynucleotide, and from one to all ends of a
20 branched polynucleotide having more than two ends. The mechanism of action of this liberation is believed to be comprised of an enzymatically-catalyzed hydrolysis of terminal nucleotides, and can be allowed to proceed in a time-dependent fashion, allowing experimental control of the progression of the enzymatic process.

25 By contrast, a non-enzymatic step may be used to shuffle, assemble, reassemble, recombine, and/or concatenate polynucleotide building blocks that is comprised of subjecting a working sample to denaturing (or "melting") conditions (for example, by changing temperature, pH, and /or salinity conditions) so as to melt a working set of double stranded polynucleotides into single polynucleotide strands. For shuffling, it is
30 desirable that the single polynucleotide strands participate to some extent in annealment with different hybridization partners (i.e. and not merely revert to exclusive

reannealment between what were former partners before the denaturation step). The presence of the former hybridization partners in the reaction vessel, however, does not preclude, and may sometimes even favor, reannealment of a single stranded polynucleotide with its former partner, to recreate an original double stranded polynucleotide.

In contrast to this non-enzymatic shuffling step comprised of subjecting double stranded polynucleotide building blocks to denaturation, followed by annealment, the instant invention further provides an exonuclease-based approach requiring no denaturation – rather, the avoidance of denaturing conditions and the maintenance of double stranded polynucleotide substrates in annealed (i.e. non-denatured) state are necessary conditions for the action of exonucleases (e.g., exonuclease III and red alpha gene product). Additionally in contrast, the generation of single stranded polynucleotide sequences capable of hybridizing to other single stranded polynucleotide sequences is the result of covalent cleavage – and hence sequence destruction - in one of the hybridization partners. For example, an exonuclease III enzyme may be used to enzymatically liberate 3' terminal nucleotides in one hybridization strand (to achieve covalent hydrolysis in that polynucleotide strand); and this favors hybridization of the remaining single strand to a new partner (since its former partner was subjected to covalent cleavage).

By way of further illustration, a specific exonuclease, namely exonuclease III is provided herein as an example of a 3' exonuclease; however, other exonucleases may also be used, including enzymes with 5' exonuclease activity and enzymes with 3' exonuclease activity, and including enzymes not yet discovered and enzymes not yet developed. It is particularly appreciated that enzymes can be discovered, optimized (e.g. engineered by directed evolution), or both discovered and optimized specifically for the instantly disclosed approach that have more optimal rates &/or more highly specific activities &/or greater lack of unwanted activities. In fact it is expected that the instant invention may encourage the discovery &/or development of such designer enzymes. In sum, this invention may be practiced with a variety of currently available

exonuclease enzymes, as well as enzymes not yet discovered and enzymes not yet developed.

The exonuclease action of exonuclease III requires a working double stranded
5 polynucleotide end that is either blunt or has a 5' overhang, and the exonuclease action
is comprised of enzymatically liberating 3' terminal nucleotides, leaving a single
stranded 5' end that becomes longer and longer as the exonuclease action proceeds (see
Figure 1). Any 5' overhangs produced by this approach may be used to hybridize to
another single stranded polynucleotide sequence (which may also be a single stranded
10 polynucleotide or a terminal overhang of a partially double stranded polynucleotide)
that shares enough homology to allow hybridization. The ability of these exonuclease
III-generated single stranded sequences (e.g. in 5' overhangs) to hybridize to other
single stranded sequences allows two or more polynucleotides to be shuffled,
assembled, reassembled, &/or concatenated.

15

Furthermore, it is appreciated that one can protect the end of a double stranded
polynucleotide or render it susceptible to a desired enzymatic action of a serviceable
exonuclease as necessary. For example, a double stranded polynucleotide end having a
3' overhang is not susceptible to the exonuclease action of exonuclease III. However, it
20 may be rendered susceptible to the exonuclease action of exonuclease III by a variety of
means; for example, it may be blunted by treatment with a polymerase, cleaved to
provide a blunt end or a 5' overhang, joined (ligated or hybridized) to another double
stranded polynucleotide to provide a blunt end or a 5' overhang, hybridized to a single
stranded polynucleotide to provide a blunt end or a 5' overhang, or modified by any of a
25 variety of means).

According to one aspect, an exonuclease may be allowed to act on one or on
both ends of a linear double stranded polynucleotide and proceed to completion, to near
completion, or to partial completion. When the exonuclease action is allowed to go to
30 completion, the result will be that the length of each 5' overhang will extend far towards
the middle region of the polynucleotide in the direction of what might be considered a

“rendezvous point” (which may be somewhere near the polynucleotide midpoint).
Ultimately, this results in the production of single stranded polynucleotides (that can become dissociated) that are each about half the length of the original double stranded polynucleotide (see Figure 1). Alternatively, an exonuclease-mediated reaction can be terminated before proceeding to completion.

Thus this exonuclease-mediated approach is serviceable for shuffling, assembling &/or reassembling, recombining, and concatenating polynucleotide building blocks, which polynucleotide building blocks can be up to ten bases long or tens of bases long or hundreds of bases long or thousands of bases long or tens of thousands of bases long or hundreds of thousands of bases long or millions of bases long or even longer.

This exonuclease-mediated approach is based on the action of double stranded DNA specific exodeoxyribonuclease activity of *E. coli* exonuclease III. Substrates for exonuclease III may be generated by subjecting a double stranded polynucleotide to fragmentation. Fragmentation may be achieved by mechanical means (e.g., shearing, sonication, etc.), by enzymatic means (e.g. using restriction enzymes), and by any combination thereof. Fragments of a larger polynucleotide may also be generated by polymerase-mediated synthesis.

Exonuclease III is a 28K monomeric enzyme, product of the *xthA* gene of *E. coli* with four known activities: exodeoxyribonuclease (alternatively referred to as exonuclease herein), RNaseH, DNA-3'-phosphatase, and AP endonuclease. The exodeoxyribonuclease activity is specific for double stranded DNA. The mechanism of action is thought to involve enzymatic hydrolysis of DNA from a 3' end progressively towards a 5' direction, with formation of nucleoside 5'-phosphates and a residual single strand. The enzyme does not display efficient hydrolysis of single stranded DNA, single-stranded RNA, or double-stranded RNA; however it degrades RNA in an DNA-RNA hybrid releasing nucleoside 5'-phosphates. The enzyme also releases inorganic phosphate specifically from 3'phosphomonoester groups on DNA, but not from RNA or

short oligonucleotides. Removal of these groups converts the terminus into a primer for DNA polymerase action.

Additional examples of enzymes with exonuclease activity include red-alpha
5 and venom phosphodiesterases. Red alpha (*redα*) gene product (also referred to as
lambda exonuclease) is of bacteriophage λ origin. The *redα* gene is transcribed from
the leftward promoter and its product is involved (24 kD) in recombination. Red alpha
gene product acts processively from 5'-phosphorylated termini to liberate
mononucleotides from duplex DNA (Takahashi & Kobayashi, 1990). Venom
10 phosphodiesterases (Laskowski, 1980) is capable of rapidly opening supercoiled DNA.

2.11.2.3. NON-STOCHASTIC LIGATION REASSEMBLY

In one aspect, the present invention provides a non-stochastic method termed synthetic ligation reassembly (SLR), that is somewhat related to stochastic shuffling, save that the nucleic acid building blocks are not shuffled or concatenated or chimerized randomly, but rather are assembled non-stochastically.

A particularly glaring difference is that the instant SLR method does not depend on the presence of a high level of homology between polynucleotides to be shuffled. In contrast, prior methods, particularly prior stochastic shuffling methods require that presence of a high level of homology, particularly at coupling sites, between polynucleotides to be shuffled. Accordingly these prior methods favor the regeneration of the original progenitor molecules, and are suboptimal for generating large numbers of novel progeny chimeras, particularly full-length progenies. The instant invention, on the other hand, can be used to non-stochastically generate libraries (or sets) of progeny molecules comprised of over 10^{100} different chimeras. Conceivably, SLR can even be used to generate libraries comprised of over 10^{1000} different progeny chimeras with (no upper limit in sight).

Thus, in one aspect, the present invention provides a method, which method is non-stochastic, of producing a set of finalized chimeric nucleic acid molecules having an overall assembly order that is chosen by design, which method is comprised of the steps of generating by design a plurality of specific nucleic acid building blocks having serviceable mutually compatible ligatable ends, and assembling these nucleic acid building blocks, such that a designed overall assembly order is achieved.

The mutually compatible ligatable ends of the nucleic acid building blocks to be assembled are considered to be "serviceable" for this type of ordered assembly if they enable the building blocks to be coupled in predetermined orders. Thus, in one aspect, the overall assembly order in which the nucleic acid building blocks can be coupled is specified by the design of the ligatable ends and, if more than one assembly step is to be

used, then the overall assembly order in which the nucleic acid building blocks can be coupled is also specified by the sequential order of the assembly step(s). Figure 4, Panel C illustrates an exemplary assembly process comprised of 2 sequential steps to achieve a designed (non-stochastic) overall assembly order for five nucleic acid building blocks. In a preferred embodiment of this invention, the annealed building pieces are treated with an enzyme, such as a ligase (e.g. T4 DNA ligase), achieve covalent bonding of the building pieces.

In a preferred embodiment, the design of nucleic acid building blocks is obtained upon analysis of the sequences of a set of progenitor nucleic acid templates that serve as a basis for producing a progeny set of finalized chimeric nucleic acid molecules. These progenitor nucleic acid templates thus serve as a source of sequence information that aids in the design of the nucleic acid building blocks that are to be mutagenized, i.e. chimerized or shuffled.

In one exemplification, this invention provides for the chimerization of a family of related genes and their encoded family of related products. In a particular exemplification, the encoded products are enzymes. As a representative list of families of enzymes which may be mutagenized in accordance with the aspects of the present invention, there may be mentioned, the following enzymes and their functions:

1 Lipase/Esterase

- a. Enantioselective hydrolysis of esters (lipids)/ thioesters
 - 1) Resolution of racemic mixtures
 - 2) Synthesis of optically active acids or alcohols from *meso*-diesters
- b. Selective syntheses
 - 1) Regiospecific hydrolysis of carbohydrate esters
 - 2) Selective hydrolysis of cyclic secondary alcohols
- c. Synthesis of optically active esters, lactones, acids, alcohols
 - 1) Transesterification of activated/nonactivated esters
 - 2) Interesterification

- 3) Optically active lactones from hydroxyesters
 - 4) Regio- and enantioselective ring opening of anhydrides
 - d. Detergents
 - e. Fat/Oil conversion
 - 5 f. Cheese ripening
- 2 Protease**
- a. Ester/amide synthesis
 - b. Peptide synthesis
 - 10 c. Resolution of racemic mixtures of amino acid esters
 - d. Synthesis of non-natural amino acids
 - e. Detergents/protein hydrolysis
- 3 Glycosidase/Glycosyl transferase**
- 15 a. Sugar/polymer synthesis
 - b. Cleavage of glycosidic linkages to form mono, di-and oligosaccharides
 - c. Synthesis of complex oligosaccharides
 - d. Glycoside synthesis using UDP-galactosyl transferase
 - e. Transglycosylation of disaccharides, glycosyl fluorides, aryl galactosides
 - 20 f. Glycosyl transfer in oligosaccharide synthesis
 - g. Diastereoselective cleavage of β -glucosylsulfoxides
 - h. Asymmetric glycosylations
 - i. Food processing
 - j. Paper processing
- 4 Phosphatase/Kinase**
- a. Synthesis/hydrolysis of phosphate esters
 - 1) Regio-, enantioselective phosphorylation
 - 2) Introduction of phosphate esters
 - 30 3) Synthesize phospholipid precursors
 - 4) Controlled polynucleotide synthesis

- b. Activate biological molecule
- c. Selective phosphate bond formation without protecting groups

5 Mono/Dioxygenase

- 5 a. Direct oxyfunctionalization of unactivated organic substrates
- b. Hydroxylation of alkane, aromatics, steroids
- c. Epoxidation of alkenes
- d. Enantioselective sulphoxidation
- e. Regio- and stereoselective Bayer-Villiger oxidations

10

6 Haloperoxidase

- a. Oxidative addition of halide ion to nucleophilic sites
- b. Addition of hypohalous acids to olefinic bonds
- c. Ring cleavage of cyclopropanes
- 15 d. Activated aromatic substrates converted to *ortho* and *para* derivatives
- e. 1,3 diketones converted to 2-halo-derivatives
- f. Heteroatom oxidation of sulfur and nitrogen containing substrates
- g. Oxidation of enol acetates, alkynes and activated aromatic rings

20 7 Lignin peroxidase/Diarylpropane peroxidase

- a. Oxidative cleavage of C-C bonds
- b. Oxidation of benzylic alcohols to aldehydes
- c. Hydroxylation of benzylic carbons
- d. Phenol dimerization
- 25 e. Hydroxylation of double bonds to form diols
- f. Cleavage of lignin aldehydes

8 Epoxide hydrolase

- a. Synthesis of enantiomerically pure bioactive compounds
- 30 b. Regio- and enantioselective hydrolysis of epoxide
- c. Aromatic and olefinic epoxidation by monooxygenases to form epoxides

- d. Resolution of racemic epoxides
- e. Hydrolysis of steroid epoxides

9 Nitrile hydratase/nitrilase

- 5 a. Hydrolysis of aliphatic nitriles to carboxamides
- b. Hydrolysis of aromatic, heterocyclic, unsaturated aliphatic nitriles to corresponding acids
- c. Hydrolysis of acrylonitrile
- d. Production of aromatic and carboxamides, carboxylic acids
- 10 (nicotinamide, picolinamide, isonicotinamide)
- e. Regioselective hydrolysis of acrylic dinitrile
- f. α -amino acids from α -hydroxynitriles

10 Transaminase

- 15 a. Transfer of amino groups into oxo-acids

11 Amidase/Acylase

- a. Hydrolysis of amides, amidines, and other C-N bonds
- b. Non-natural amino acid resolution and synthesis

20

These exemplifications, while illustrating certain specific aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

- 25 Thus according to one aspect of this invention, the sequences of a plurality of progenitor nucleic acid templates are aligned in order to select one or more demarcation points, which demarcation points can be located at an area of homology, and are comprised of one or more nucleotides, and which demarcation points are shared by at least two of the progenitor templates. The demarcation points can be used to delineate
- 30 the boundaries of nucleic acid building blocks to be generated. Thus, the demarcation

points identified and selected in the progenitor molecules serve as potential chimerization points in the assembly of the progeny molecules.

Preferably a serviceable demarcation point is an area of homology (comprised of
5 at least one homologous nucleotide base) shared by at least two progenitor templates.
More preferably a serviceable demarcation point is an area of homology that is shared
by at least half of the progenitor templates. More preferably still a serviceable
demarcation point is an area of homology that is shared by at least two thirds of the
progenitor templates. Even more preferably a serviceable demarcation points is an area
10 of homology that is shared by at least three fourths of the progenitor templates. Even
more preferably still a serviceable demarcation points is an area of homology that is
shared by at almost all of the progenitor templates. Even more preferably still a
serviceable demarcation point is an area of homology that is shared by all of the
progenitor templates.

15

The process of designing nucleic acid building blocks and of designing the
mutually compatible ligatable ends of the nucleic acid building blocks to be assembled
is illustrated in Figures 6 and 7. As shown, the alignment of a set of progenitor
templates reveals several naturally occurring demarcation points, and the identification
20 of demarcation points shared by these templates helps to non-stochastically determine
the building blocks to be generated and used for the generation of the progeny chimeric
molecules.

In a preferred embodiment, this invention provides that the ligation reassembly
25 process is performed exhaustively in order to generate an exhaustive library. In other
words, all possible ordered combinations of the nucleic acid building blocks are
represented in the set of finalized chimeric nucleic acid molecules. At the same time, in
a particularly preferred embodiment, the assembly order (i.e. the order of assembly of
each building block in the 5' to 3' sequence of each finalized chimeric nucleic acid) in
30 each combination is by design (or non-stochastic). Because of the non-stochastic nature
of this invention, the possibility of unwanted side products is greatly reduced.

In another preferred embodiment, this invention provides that the ligation reassembly process is performed systematically, for example in order to generate a systematically compartmentalized library, with compartments that can be screened systematically, e.g. one by one. In other words this invention provides that, through the selective and judicious use of specific nucleic acid building blocks, coupled with the selective and judicious use of sequentially stepped assembly reactions, an experimental design can be achieved where specific sets of progeny products are made in each of several reaction vessels. This allows a systematic examination and screening procedure to be performed. Thus, it allows a potentially very large number of progeny molecules to be examined systematically in smaller groups.

Because of its ability to perform chimerizations in a manner that is highly flexible yet exhaustive and systematic as well, particularly when there is a low level of homology among the progenitor molecules, the instant invention provides for the generation of a library (or set) comprised of a large number of progeny molecules. Because of the non-stochastic nature of the instant ligation reassembly invention, the progeny molecules generated preferably comprise a library of finalized chimeric nucleic acid molecules having an overall assembly order that is chosen by design. In a particularly preferred embodiment of this invention, such a generated library is comprised of preferably greater than 10^3 different progeny molecular species, more preferably greater than 10^5 different progeny molecular species, more preferably still greater than 10^{10} different progeny molecular species, more preferably still greater than 10^{15} different progeny molecular species, more preferably still greater than 10^{20} different progeny molecular species, more preferably still greater than 10^{30} different progeny molecular species, more preferably still greater than 10^{40} different progeny molecular species, more preferably still greater than 10^{50} different progeny molecular species, more preferably still greater than 10^{60} different progeny molecular species, more preferably still greater than 10^{70} different progeny molecular species, more preferably still greater than 10^{80} different progeny molecular species, more preferably still greater than 10^{100} different progeny molecular species, more preferably still greater

than 10^{110} different progeny molecular species, more preferably still greater than 10^{120} different progeny molecular species, more preferably still greater than 10^{130} different progeny molecular species, more preferably still greater than 10^{140} different progeny molecular species, more preferably still greater than 10^{150} different progeny molecular species, more preferably still greater than 10^{175} different progeny molecular species, more preferably still greater than 10^{200} different progeny molecular species, more preferably still greater than 10^{300} different progeny molecular species, more preferably still greater than 10^{400} different progeny molecular species, more preferably still greater than 10^{500} different progeny molecular species, and even more preferably still greater than 10^{1000} different progeny molecular species.

In one aspect, a set of finalized chimeric nucleic acid molecules, produced as described is comprised of a polynucleotide encoding a polypeptide. According to one preferred embodiment, this polynucleotide is a gene, which may be a man-made gene. According to another preferred embodiment, this polynucleotide is a gene pathway, which may be a man-made gene pathway. This invention provides that one or more man-made genes generated by this invention may be incorporated into a man-made gene pathway, such as a pathway operable in a eukaryotic organism (including a plant).

It is appreciated that the power of this invention is exceptional, as there is much freedom of choice and control regarding the selection of demarcation points, the size and number of the nucleic acid building blocks, and the size and design of the couplings. It is appreciated, furthermore, that the requirement for intermolecular homology is highly relaxed for the operability of this invention. In fact, demarcation points can even be chosen in areas of little or no intermolecular homology. For example, because of codon wobble, i.e. the degeneracy of codons, nucleotide substitutions can be introduced into nucleic acid building blocks without altering the amino acid originally encoded in the corresponding progenitor template. Alternatively, a codon can be altered such that the coding for an original amino acid is altered. This invention provides that such substitutions can be introduced into the nucleic acid building block in order to increase the incidence of intermolecularly homologous

demarcation points and thus to allow an increased number of couplings to be achieved among the building blocks, which in turn allows a greater number of progeny chimeric molecules to be generated.

5 In another exemplification, the synthetic nature of the step in which the building blocks are generated allows the design and introduction of nucleotides (e.g. one or more nucleotides, which may be, for example, codons or introns or regulatory sequences) that can later be optionally removed in an in vitro process (e.g. by mutagenesis) or in an in vivo process (e.g. by utilizing the gene splicing ability of a host organism). It is
10 appreciated that in many instances the introduction of these nucleotides may also be desirable for many other reasons in addition to the potential benefit of creating a serviceable demarcation point.

 Thus, according to another embodiment, this invention provides that a nucleic
15 acid building block can be used to introduce an intron. Thus, this invention provides that functional introns may be introduced into a man-made gene of this invention. This invention also provides that functional introns may be introduced into a man-made gene pathway of this invention. Accordingly, this invention provides for the generation of a chimeric polynucleotide that is a man-made gene containing one (or more) artificially
20 introduced intron(s).

 Accordingly, this invention also provides for the generation of a chimeric polynucleotide that is a man-made gene pathway containing one (or more) artificially introduced intron(s). Preferably, the artificially introduced intron(s) are functional in
25 one or more host cells for gene splicing much in the way that naturally-occurring introns serve functionally in gene splicing. This invention provides a process of producing man-made intron-containing polynucleotides to be introduced into host organisms for recombination and/or splicing.

30 The ability to achieve chimerizations, using couplings as described herein, in areas of little or no homology among the progenitor molecules, is particularly useful,

and in fact critical, for the assembly of novel gene pathways. This invention thus provides for the generation of novel man-made gene pathways using synthetic ligation reassembly. In a particular aspect, this is achieved by the introduction of regulatory sequences, such as promoters, that are operable in an intended host, to confer

5 operability to a novel gene pathway when it is introduced into the intended host. In a particular exemplification, this invention provides for the generation of novel man-made gene pathways that is operable in a plurality of intended hosts (e.g. in a microbial organism as well as in a plant cell). This can be achieved, for example, by the introduction of a plurality of regulatory sequences, comprised of a regulatory sequence

10 that is operable in a first intended host and a regulatory sequence that is operable in a second intended host. A similar process can be performed to achieve operability of a gene pathway in a third intended host species, etc. The number of intended host species can be each integer from 1 to 10 or alternatively over 10. Alternatively, for example, operability of a gene pathway in a plurality of intended hosts can be achieved by the

15 introduction of a regulatory sequence having intrinsic operability in a plurality of intended hosts.

Thus, according to a particular embodiment, this invention provides that a nucleic acid building block can be used to introduce a regulatory sequence, particularly

20 a regulatory sequence for gene expression. Preferred regulatory sequences include, but are not limited to, those that are man-made, and those found in archeal, bacterial, eukaryotic (including mitochondrial), viral, and prionic or prion-like organisms. Preferred regulatory sequences include but are not limited to, promoters, operators, and activator binding sites. Thus, this invention provides that functional regulatory

25 sequences may be introduced into a man-made gene of this invention. This invention also provides that functional regulatory sequences may be introduced into a man-made gene pathway of this invention.

Accordingly, this invention provides for the generation of a chimeric

30 polynucleotide that is a man-made gene containing one (or more) artificially introduced regulatory sequence(s). Accordingly, this invention also provides for the generation of

a chimeric polynucleotide that is a man-made gene pathway containing one (or more) artificially introduced regulatory sequence(s). Preferably, an artificially introduced regulatory sequence(s) is operatively linked to one or more genes in the man-made polynucleotide, and are functional in one or more host cells.

5

Preferred bacterial promoters that are serviceable for this invention include lacI, lacZ, T3, T7, gpt, lambda P_R, P_L and trp. Serviceable eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Particular plant regulatory sequences include promoters active in directing transcription in plants, either constitutively or stage and/or tissue specific, depending on the use of the plant or parts thereof. These promoters include, but are not limited to promoters showing constitutive expression, such as the 35S promoter of Cauliflower Mosaic Virus (CaMV) (Guilley et al., 1982), those for leaf-specific expression, such as the promoter of the ribulose biphosphate carboxylase small subunit gene (Coruzzi et al., 1984), those for root-specific expression, such as the promoter from the glutamin synthase gene (Tingey et al., 1987), those for seed-specific expression, such as the cruciferin A promoter from *Brassica napus* (Ryan et al., 1989), those for tuber-specific expression, such as the class-I patatin promoter from potato (Rocha-Sasa et al., 1989; Wenzler et al., 1989) or those for fruit-specific expression, such as the polygalacturonase (PG) promoter from tomato (Bird et al., 1988).

Other regulatory sequences that are preferred for this invention include terminator sequences and polyadenylation signals and any such sequence functioning as such in plants, the choice of which is within the level of the skilled artisan. An example of such sequences is the 3' flanking region of the nopaline synthase (nos) gene of *Agrobacterium tumefaciens* (Bevan, 1984). The regulatory sequences may also include enhancer sequences, such as found in the 35S promoter of CaMV, and mRNA stabilizing sequences such as the leader sequence of Alfalfa Mosaic Virus (AIMV) RNA4 (Brederode et al., 1980) or any other sequences functioning in a like manner.

30

Man-made genes produced using this invention can also serve as a substrate for recombination with another nucleic acid. Likewise, a man-made gene pathway produced using this invention can also serve as a substrate for recombination with another nucleic acid. In a preferred instance, the recombination is facilitated by, or
5 occurs at, areas of homology between the man-made intron-containing gene and a nucleic acid which serves as a recombination partner. In a particularly preferred instance, the recombination partner may also be a nucleic acid generated by this invention, including a man-made gene or a man-made gene pathway. Recombination may be facilitated by or may occur at areas of homology that exist at the one (or more)
10 artificially introduced intron(s) in the man-made gene.

The synthetic ligation reassembly method of this invention utilizes a plurality of nucleic acid building blocks, each of which preferably has two ligatable ends. The two ligatable ends on each nucleic acid building block may be two blunt ends (i.e. each
15 having an overhang of zero nucleotides), or preferably one blunt end and one overhang, or more preferably still two overhangs.

A serviceable overhang for this purpose may be a 3' overhang or a 5' overhang. Thus, a nucleic acid building block may have a 3' overhang or alternatively a 5'
20 overhang or alternatively two 3' overhangs or alternatively two 5' overhangs. The overall order in which the nucleic acid building blocks are assembled to form a finalized chimeric nucleic acid molecule is determined by purposeful experimental design and is not random.

25 According to one preferred embodiment, a nucleic acid building block is generated by chemical synthesis of two single-stranded nucleic acids (also referred to as single-stranded oligos) and contacting them so as to allow them to anneal to form a double-stranded nucleic acid building block.

30 A double-stranded nucleic acid building block can be of variable size. The sizes of these building blocks can be small or large depending on the choice of the

experimenter. Preferred sizes for building block range from 1 base pair (not including any overhangs) to 100,000 base pairs (not including any overhangs). Other preferred size ranges are also provided, which have lower limits of from 1 bp to 10,000 bp (including every integer value in between), and upper limits of from 2 bp to 100, 000 bp (including every integer value in between).

It is appreciated that current methods of polymerase-based amplification can be used to generate double-stranded nucleic acids of up to thousands of base pairs, if not tens of thousands of base pairs, in length with high fidelity. Chemical synthesis (e.g. phosphoramidite-based) can be used to generate nucleic acids of up to hundreds of nucleotides in length with high fidelity; however, these can be assembled, e.g. using overhangs or sticky ends, to form double-stranded nucleic acids of up to thousands of base pairs, if not tens of thousands of base pairs, in length if so desired.

A combination of methods (e.g. phosphoramidite-based chemical synthesis and PCR) can also be used according to this invention. Thus, nucleic acid building block made by different methods can also be used in combination to generate a progeny molecule of this invention.

The use of chemical synthesis to generate nucleic acid building blocks is particularly preferred in this invention & is advantageous for other reasons as well, including procedural safety and ease. No cloning or harvesting or actual handling of any biological samples is required. The design of the nucleic acid building blocks can be accomplished on paper. Accordingly, this invention teaches an advance in procedural safety in recombinant technologies.

Nonetheless, according to one preferred embodiment, a double-stranded nucleic acid building block according to this invention may also be generated by polymerase-based amplification of a polynucleotide template. In a non-limiting exemplification, as illustrated in Figure 2, a first polymerase-based amplification reaction using a first set of primers, F_2 and R_1 , is used to generate a blunt-ended product (labeled Reaction 1,

Product 1), which is essentially identical to Product A. A second polymerase-based amplification reaction using a second set of primers, F₁ and R₂, is used to generate a blunt-ended product (labeled Reaction 2, Product 2), which is essentially identical to Product B. These two products are mixed and allowed to melt and anneal, generating
5 potentially useful double-stranded nucleic acid building blocks with two overhangs. In the example of Fig. 2, the product with the 3' overhangs (Product C) is selected by nuclease-based degradation of the other 3 products using a 3' acting exonuclease, such as exonuclease III. It is appreciated that a 5' acting exonuclease (e.g. red alpha) may be also be used, for example to select Product D instead. It is also appreciated that other
10 selection means can also be used, including hybridization-based means, and that these means can incorporate a further means, such as a magnetic bead-based means, to facilitate separation of the desired product.

Many other methods exist by which a double-stranded nucleic acid building
15 block can be generated that is serviceable for this invention; and these are known in the art and can be readily performed by the skilled artisan.

According to particularly preferred embodiment, a double-stranded nucleic acid building block that is serviceable for this invention is generated by first generating two
20 single stranded nucleic acids and allowing them to anneal to form a double-stranded nucleic acid building block. The two strands of a double-stranded nucleic acid building block may be complementary at every nucleotide apart from any that form an overhang; thus containing no mismatches, apart from any overhang(s). According to another embodiment, the two strands of a double-stranded nucleic acid building block are
25 complementary at fewer than every nucleotide apart from any that form an overhang. Thus, according to this embodiment, a double-stranded nucleic acid building block can be used to introduce codon degeneracy. Preferably the codon degeneracy is introduced using the site-saturation mutagenesis described herein, using one or more N,N,G/T cassettes or alternatively using one or more N,N,N cassettes.

30

Contained within an exemplary experimental design for achieving an ordered assembly according to this invention are:

- 1) The design of specific nucleic acid building blocks.
- 5 2) The design of specific ligatable ends on each nucleic acid building block.
- 3) The design of a particular order of assembly of the nucleic acid building blocks.

An overhang may be a 3' overhang or a 5' overhang. An overhang may also
10 have a terminal phosphate group or alternatively may be devoid of a terminal phosphate group (having, e.g., a hydroxyl group instead). An overhang may be comprised of any number of nucleotides. Preferably an overhang is comprised of 0 nucleotides (as in a blunt end) to 10,000 nucleotides. Thus, a wide range of overhang sizes may be serviceable. Accordingly, the lower limit may be each integer from 1-200 and the upper
15 limit may be each integer from 2-10,000. According to a particular exemplification, an overhang may consist of anywhere from 1 nucleotide to 200 nucleotides (including every integer value in between).

The final chimeric nucleic acid molecule may be generated by sequentially
20 assembling 2 or more building blocks at a time until all the designated building blocks have been assembled. A working sample may optionally be subjected to a process for size selection or purification or other selection or enrichment process between the performance of two assembly steps. Alternatively, the final chimeric nucleic acid molecule may be generated by assembling all the designated building blocks at once in
25 one step.

In vitro Shuffling

The equivalents of some standard genetic matings may also be performed by shuffling *in vitro*. For example, a "molecular backcross" can be performed by repeatedly mixing the hybrid's nucleic acid with the wild-type nucleic acid while
5 selecting for the mutations of interest. As in traditional breeding, this approach can be used to combine phenotypes from different sources into a background of choice. It is useful, for example, for the removal of neutral mutations that affect unselected characteristics (i.e. immunogenicity). Thus it can be useful to determine which mutations in a protein are involved in the enhanced biological activity and which are
10 not, an advantage which cannot be achieved by error-prone mutagenesis or cassette mutagenesis methods.

Large, functional genes can be assembled correctly from a mixture of small random polynucleotides. This reaction may be of use for the reassembly of genes from
15 the highly fragmented DNA of fossils. In addition random nucleic acid fragments from fossils may be combined with polynucleotides from similar genes from related species.

It is also contemplated that the method of this invention can be used for the *in vitro* amplification of a whole genome from a single cell as is needed for a variety of
20 research and diagnostic applications. DNA amplification by PCR is in practice limited to a length of about 40 kb. Amplification of a whole genome such as that of *E. coli* (5,000 kb) by PCR would require about 250 primers yielding 125 forty kb polynucleotides. This approach is not practical due to the unavailability of sufficient sequence data. On the other hand, random production of polynucleotides of the genome with sexual PCR
25 cycles, followed by gel purification of small polynucleotides will provide a multitude of possible primers. Use of this mix of random small polynucleotides as primers in a PCR reaction alone or with the whole genome as the template should result in an inverse chain reaction with the theoretical endpoint of a single concatemer containing many copies of the genome.

30

100 fold amplification in the copy number and an average polynucleotide size of greater than 50 kb may be obtained when only random polynucleotides are used. It is thought that the larger concatemer is generated by overlap of many smaller polynucleotides. The quality of specific PCR products obtained using synthetic primers will be indistinguishable from the product obtained from unamplified DNA. It is expected that this approach will be useful for the mapping of genomes.

The polynucleotide to be shuffled can be produced as random or non-random polynucleotides, at the discretion of the practitioner.

10

In vivo Shuffling

In an embodiment of *in vivo* shuffling, the mixed population of the specific nucleic acid sequence is introduced into bacterial or eukaryotic cells under conditions such that at least two different nucleic acid sequences are present in each host cell. The polynucleotides can be introduced into the host cells by a variety of different methods. The host cells can be transformed with the smaller polynucleotides using methods known in the art, for example treatment with calcium chloride. If the polynucleotides are inserted into a phage genome, the host cell can be transfected with the recombinant phage genome having the specific nucleic acid sequences. Alternatively, the nucleic acid sequences can be introduced into the host cell using electroporation, transfection, lipofection, biolistics, conjugation, and the like.

In general, in this embodiment, the specific nucleic acids sequences will be present in vectors which are capable of stably replicating the sequence in the host cell. In addition, it is contemplated that the vectors will encode a marker gene such that host cells having the vector can be selected. This ensures that the mutated specific nucleic acid sequence can be recovered after introduction into the host cell. However, it is contemplated that the entire mixed population of the specific nucleic acid sequences need not be present on a vector sequence. Rather only a sufficient number of sequences need be cloned into vectors to ensure that after introduction of the polynucleotides into the host cells each host cell contains one vector having at least one specific nucleic acid sequence present therein. It is also contemplated that rather than having a subset of the population of the specific nucleic acids sequences cloned into vectors, this subset may be already stably integrated into the host cell.

It has been found that when two polynucleotides which have regions of identity are inserted into the host cells homologous recombination occurs between the two polynucleotides. Such recombination between the two mutated specific nucleic acid sequences will result in the production of double or triple hybrids in some situations.

It has also been found that the frequency of recombination is increased if some of the mutated specific nucleic acid sequences are present on linear nucleic acid molecules. Therefore, in a preferred embodiment, some of the specific nucleic acid sequences are present on linear polynucleotides.

5

After transformation, the host cell transformants are placed under selection to identify those host cell transformants which contain mutated specific nucleic acid sequences having the qualities desired. For example, if increased resistance to a particular drug is desired then the transformed host cells may be subjected to increased concentrations of the particular drug and those transformants producing mutated proteins able to confer increased drug resistance will be selected. If the enhanced ability of a particular protein to bind to a receptor is desired, then expression of the protein can be induced from the transformants and the resulting protein assayed in a ligand binding assay by methods known in the art to identify that subset of the mutated population which shows enhanced binding to the ligand. Alternatively, the protein can be expressed in another system to ensure proper processing.

10
15

Once a subset of the first recombined specific nucleic acid sequences (daughter sequences) having the desired characteristics are identified, they are then subject to a second round of recombination.

20

In the second cycle of recombination, the recombined specific nucleic acid sequences may be mixed with the original mutated specific nucleic acid sequences (parent sequences) and the cycle repeated as described above. In this way a set of second recombined specific nucleic acids sequences can be identified which have enhanced characteristics or encode for proteins having enhanced properties. This cycle can be repeated a number of times as desired.

25

It is also contemplated that in the second or subsequent recombination cycle, a backcross can be performed. A molecular backcross can be performed by mixing the desired specific nucleic acid sequences with a large number of the wild-type sequence,

30

such that at least one wild-type nucleic acid sequence and a mutated nucleic acid sequence are present in the same host cell after transformation. Recombination with the wild-type specific nucleic acid sequence will eliminate those neutral mutations that may affect unselected characteristics such as immunogenicity but not the selected
5 characteristics.

In another embodiment of this invention, it is contemplated that during the first round a subset of the specific nucleic acid sequences can be generated as smaller polynucleotides by slowing or halting their PCR amplification prior to introduction into
10 the host cell. The size of the polynucleotides must be large enough to contain some regions of identity with the other sequences so as to homologously recombine with the other sequences. The size of the polynucleotides will range from 0.03 kb to 100 kb more preferably from 0.2 kb to 10 kb. It is also contemplated that in subsequent rounds, all of the specific nucleic acid sequences other than the sequences selected from
15 the previous round may be utilized to generate PCR polynucleotides prior to introduction into the host cells.

The shorter polynucleotide sequences can be single-stranded or double-stranded. If the sequences were originally single-stranded and have become double-stranded they
20 can be denatured with heat, chemicals or enzymes prior to insertion into the host cell. The reaction conditions suitable for separating the strands of nucleic acid are well known in the art.

The steps of this process can be repeated indefinitely, being limited only by the
25 number of possible hybrids which can be achieved. After a certain number of cycles, all possible hybrids will have been achieved and further cycles are redundant.

In an embodiment the same mutated template nucleic acid is repeatedly recombined and the resulting recombinants selected for the desired characteristic.

30 Therefore, the initial pool or population of mutated template nucleic acid is cloned into a vector capable of replicating in a bacteria such as *E. coli*. The particular

vector is not essential, so long as it is capable of autonomous replication in *E. coli*. In a preferred embodiment, the vector is designed to allow the expression and production of any protein encoded by the mutated specific nucleic acid linked to the vector. It is also preferred that the vector contain a gene encoding for a selectable marker.

5

The population of vectors containing the pool of mutated nucleic acid sequences is introduced into the *E. coli* host cells. The vector nucleic acid sequences may be introduced by transformation, transfection or infection in the case of phage. The concentration of vectors used to transform the bacteria is such that a number of vectors is introduced into each cell. Once present in the cell, the efficiency of homologous recombination is such that homologous recombination occurs between the various vectors. This results in the generation of hybrids (daughters) having a combination of mutations which differ from the original parent mutated sequences.

10

The host cells are then clonally replicated and selected for the marker gene present on the vector. Only those cells having a plasmid will grow under the selection.

15

The host cells which contain a vector are then tested for the presence of favorable mutations. Such testing may consist of placing the cells under selective pressure, for example, if the gene to be selected is an improved drug resistance gene. If the vector allows expression of the protein encoded by the mutated nucleic acid sequence, then such selection may include allowing expression of the protein so encoded, isolation of the protein and testing of the protein to determine whether, for example, it binds with increased efficiency to the ligand of interest.

20

25

Once a particular daughter mutated nucleic acid sequence has been identified which confers the desired characteristics, the nucleic acid is isolated either already linked to the vector or separated from the vector. This nucleic acid is then mixed with the first or parent population of nucleic acids and the cycle is repeated.

30

It has been shown that by this method nucleic acid sequences having enhanced desired properties can be selected.

In an alternate embodiment, the first generation of hybrids are retained in the cells and the parental mutated sequences are added again to the cells. Accordingly, the
5 first cycle of Embodiment I is conducted as described above. However, after the daughter nucleic acid sequences are identified, the host cells containing these sequences are retained.

The parent mutated specific nucleic acid population, either as polynucleotides or
10 cloned into the same vector is introduced into the host cells already containing the daughter nucleic acids. Recombination is allowed to occur in the cells and the next generation of recombinants, or granddaughters are selected by the methods described above.

15 This cycle can be repeated a number of times until the nucleic acid or peptide having the desired characteristics is obtained. It is contemplated that in subsequent cycles, the population of mutated sequences which are added to the preferred hybrids may come from the parental hybrids or any subsequent generation.

20 In an alternative embodiment, the invention provides a method of conducting a "molecular" backcross of the obtained recombinant specific nucleic acid in order to eliminate any neutral mutations. Neutral mutations are those mutations which do not confer onto the nucleic acid or peptide the desired properties. Such mutations may however confer on the nucleic acid or peptide undesirable characteristics. Accordingly,
25 it is desirable to eliminate such neutral mutations. The method of this invention provide a means of doing so.

In this embodiment, after the hybrid nucleic acid, having the desired characteristics, is obtained by the methods of the embodiments, the nucleic acid, the
30 vector having the nucleic acid or the host cell containing the vector and nucleic acid is isolated.

- The nucleic acid or vector is then introduced into the host cell with a large excess of the wild-type nucleic acid. The nucleic acid of the hybrid and the nucleic acid of the wild-type sequence are allowed to recombine. The resulting recombinants are
- 5 placed under the same selection as the hybrid nucleic acid. Only those recombinants which retained the desired characteristics will be selected. Any silent mutations which do not provide the desired characteristics will be lost through recombination with the wild-type DNA. This cycle can be repeated a number of times until all of the silent mutations are eliminated.
- 10 Thus the methods of this invention can be used in a molecular backcross to eliminate unnecessary or silent mutations.

Utility

The *in vivo* recombination method of this invention can be performed blindly on a pool of unknown hybrids or alleles of a specific polynucleotide or sequence.

However, it is not necessary to know the actual DNA or RNA sequence of the specific
5 polynucleotide.

The approach of using recombination within a mixed population of genes can be useful for the generation of any useful proteins, for example, interleukin I, antibodies, tPA and growth hormone. This approach may be used to generate proteins having
10 altered specificity or activity. The approach may also be useful for the generation of hybrid nucleic acid sequences, for example, promoter regions, introns, exons, enhancer sequences, 3' untranslated regions or 5' untranslated regions of genes. Thus this approach may be used to generate genes having increased rates of expression. This approach may also be useful in the study of repetitive DNA sequences. Finally, this
15 approach may be useful to mutate ribozymes or aptamers.

Scaffold-like regions separating regions of diversity in proteins may be particularly suitable for the methods of this invention. The conserved scaffold determines the overall folding by self-association, while displaying relatively
20 unrestricted loops that mediate the specific binding. Examples of such scaffolds are the immunoglobulin beta barrel, and the four-helix bundle. The methods of this invention can be used to create scaffold-like proteins with various combinations of mutated sequences for binding.

25 The equivalents of some standard genetic matings may also be performed by the methods of this invention. For example, a "molecular" backcross can be performed by repeated mixing of the hybrid's nucleic acid with the wild-type nucleic acid while selecting for the mutations of interest. As in traditional breeding, this approach can be used to combine phenotypes from different sources into a background of choice. It is
30 useful, for example, for the removal of neutral mutations that affect unselected characteristics (i.e. immunogenicity). Thus it can be useful to determine which

mutations in a protein are involved in the enhanced biological activity and which are not.

5

Peptide Display Methods

The present method can be used to shuffle, by *in vitro* and/or *in vivo* recombination by any of the disclosed methods, and in any combination, polynucleotide sequences selected by peptide display methods, wherein an associated polynucleotide
5 encodes a displayed peptide which is screened for a phenotype (e.g., for affinity for a predetermined receptor (ligand)).

An increasingly important aspect of bio-pharmaceutical drug development and molecular biology is the identification of peptide structures, including the primary
10 amino acid sequences, of peptides or peptidomimetics that interact with biological macromolecules. one method of identifying peptides that possess a desired structure or functional property, such as binding to a predetermined biological macromolecule (e.g., a receptor), involves the screening of a large library or peptides for individual library members which possess the desired structure or functional property conferred by the
15 amino acid sequence of the peptide.

In addition to direct chemical synthesis methods for generating peptide libraries, several recombinant DNA methods also have been reported. One type involves the display of a peptide sequence, antibody, or other protein on the surface of a
20 bacteriophage particle or cell. Generally, in these methods each bacteriophage particle or cell serves as an individual library member displaying a single species of displayed peptide in addition to the natural bacteriophage or cell protein sequences. Each bacteriophage or cell contains the nucleotide sequence information encoding the particular displayed peptide sequence; thus, the displayed peptide sequence can be
25 ascertained by nucleotide sequence determination of an isolated library member.

A well-known peptide display method involves the presentation of a peptide sequence on the surface of a filamentous bacteriophage, typically as a fusion with a bacteriophage coat protein. The bacteriophage library can be incubated with an
30 immobilized, predetermined macromolecule or small molecule (e.g., a receptor) so that bacteriophage particles which present a peptide sequence that binds to the immobilized

macromolecule can be differentially partitioned from those that do not present peptide sequences that bind to the predetermined macromolecule. The bacteriophage particles (i.e., library members) which are bound to the immobilized macromolecule are then recovered and replicated to amplify the selected bacteriophage sub-population for a subsequent round of affinity enrichment and phage replication. After several rounds of affinity enrichment and phage replication, the bacteriophage library members that are thus selected are isolated and the nucleotide sequence encoding the displayed peptide sequence is determined, thereby identifying the sequence(s) of peptides that bind to the predetermined macromolecule (e.g., receptor). Such methods are further described in PCT patent publications WO 91/17271, WO 91/18980, WO 91/19818 and WO 93/08278.

The latter PCT publication describes a recombinant DNA method for the display of peptide ligands that involves the production of a library of fusion proteins with each fusion protein composed of a first polypeptide portion, typically comprising a variable sequence, that is available for potential binding to a predetermined macromolecule, and a second polypeptide portion that binds to DNA, such as the DNA vector encoding the individual fusion protein. When transformed host cells are cultured under conditions that allow for expression of the fusion protein, the fusion protein binds to the DNA vector encoding it. Upon lysis of the host cell, the fusion protein/vector DNA complexes can be screened against a predetermined macromolecule in much the same way as bacteriophage particles are screened in the phage-based display system, with the replication and sequencing of the DNA vectors in the selected fusion protein/vector DNA complexes serving as the basis for identification of the selected library peptide sequence(s).

Other systems for generating libraries of peptides and like polymers have aspects of both the recombinant and *in vitro* chemical synthesis methods. In these hybrid methods, cell-free enzymatic machinery is employed to accomplish the *in vitro* synthesis of the library members (i.e., peptides or polynucleotides). In one type of method, RNA molecules with the ability to bind a predetermined protein or a

predetermined dye molecule were selected by alternate rounds of selection and PCR amplification (**Tuerk and Gold**, 1990; **Ellington and Szostak**, 1990). A similar technique was used to identify DNA sequences which bind a predetermined human transcription factor (**Thiesen and Bach**, 1990; **Beaudry and Joyce**, 1992; PCT patent publications **WO 92/05258** and **WO 92/14843**). In a similar fashion, the technique of *in vitro* translation has been used to synthesize proteins of interest and has been proposed as a method for generating large libraries of peptides. These methods which rely upon *in vitro* translation, generally comprising stabilized polysome complexes, are described further in PCT patent publications **WO 88/08453**, **WO 90/05785**, **WO 90/07003**, **WO 91/02076**, **WO 91/05058**, and **WO 92/02536**. Applicants have described methods in which library members comprise a fusion protein having a first polypeptide portion with DNA binding activity and a second polypeptide portion having the library member unique peptide sequence; such methods are suitable for use in cell-free *in vitro* selection formats, among others.

15

The displayed peptide sequences can be of varying lengths, typically from 3-5000 amino acids long or longer, frequently from 5-100 amino acids long, and often from about 8-15 amino acids long. A library can comprise library members having varying lengths of displayed peptide sequence, or may comprise library members having a fixed length of displayed peptide sequence. Portions or all of the displayed peptide sequence(s) can be random, pseudorandom, defined set kernel, fixed, or the like. The present display methods include methods for *in vitro* and *in vivo* display of single-chain antibodies, such as nascent scFv on polysomes or scfv displayed on phage, which enable large-scale screening of scfv libraries having broad diversity of variable region sequences and binding specificities.

25

The present invention also provides random, pseudorandom, and defined sequence framework peptide libraries and methods for generating and screening those libraries to identify useful compounds (e.g., peptides, including single-chain antibodies) that bind to receptor molecules or epitopes of interest or gene products that modify peptides or RNA in a desired fashion. The random, pseudorandom, and defined

30

sequence framework peptides are produced from libraries of peptide library members that comprise displayed peptides or displayed single-chain antibodies attached to a polynucleotide template from which the displayed peptide was synthesized. The mode of attachment may vary according to the specific embodiment of the invention selected, and can include encapsulation in a phage particle or incorporation in a cell.

A method of affinity enrichment allows a very large library of peptides and single-chain antibodies to be screened and the polynucleotide sequence encoding the desired peptide(s) or single-chain antibodies to be selected. The polynucleotide can then be isolated and shuffled to recombine combinatorially the amino acid sequence of the selected peptide(s) (or predetermined portions thereof) or single-chain antibodies (or just VHI, VLI or CDR portions thereof). Using these methods, one can identify a peptide or single-chain antibody as having a desired binding affinity for a molecule and can exploit the process of shuffling to converge rapidly to a desired high-affinity peptide or scfv. The peptide or antibody can then be synthesized in bulk by conventional means for any suitable use (e.g., as a therapeutic or diagnostic agent).

A significant advantage of the present invention is that no prior information regarding an expected ligand structure is required to isolate peptide ligands or antibodies of interest. The peptide identified can have biological activity, which is meant to include at least specific binding affinity for a selected receptor molecule and, in some instances, will further include the ability to block the binding of other compounds, to stimulate or inhibit metabolic pathways, to act as a signal or messenger, to stimulate or inhibit cellular activity, and the like.

The present invention also provides a method for shuffling a pool of polynucleotide sequences selected by affinity screening a library of polysomes displaying nascent peptides (including single-chain antibodies) for library members which bind to a predetermined receptor (e.g., a mammalian proteinaceous receptor such as, for example, a peptidergic hormone receptor, a cell surface receptor, an intracellular protein which binds to other protein(s) to form intracellular protein complexes such as

hetero-dimers and the like) or epitope (e.g., an immobilized protein, glycoprotein, oligosaccharide, and the like).

Polynucleotide sequences selected in a first selection round (typically by affinity
5 selection for binding to a receptor (e.g., a ligand)) by any of these methods are pooled
and the pool(s) is/are shuffled by *in vitro* and/or *in vivo* recombination to produce a
shuffled pool comprising a population of recombined selected polynucleotide
sequences. The recombined selected polynucleotide sequences are subjected to at least
one subsequent selection round. The polynucleotide sequences selected in the
10 subsequent selection round(s) can be used directly, sequenced, and/or subjected to one
or more additional rounds of shuffling and subsequent selection. Selected sequences
can also be back-crossed with polynucleotide sequences encoding neutral sequences
(i.e., having insubstantial functional effect on binding), such as for example by back-
crossing with a wild-type or naturally-occurring sequence substantially identical to a
15 selected sequence to produce native-like functional peptides, which may be less
immunogenic. Generally, during back-crossing subsequent selection is applied to retain
the property of binding to the predetermined receptor (ligand).

Prior to or concomitant with the shuffling of selected sequences, the sequences
20 can be mutagenized. In one embodiment, selected library members are cloned in a
prokaryotic vector (e.g., plasmid, phagemid, or bacteriophage) wherein a collection of
individual colonies (or plaques) representing discrete library members are produced.
Individual selected library members can then be manipulated (e.g., by site-directed
mutagenesis, cassette mutagenesis, chemical mutagenesis, PCR mutagenesis, and the
25 like) to generate a collection of library members representing a kernel of sequence
diversity based on the sequence of the selected library member. The sequence of an
individual selected library member or pool can be manipulated to incorporate random
mutation, pseudorandom mutation, defined kernel mutation (i.e., comprising variant and
invariant residue positions and/or comprising variant residue positions which can
30 comprise a residue selected from a defined subset of amino acid residues), codon-based
mutation, and the like, either segmentally or over the entire length of the individual

selected library member sequence. The mutagenized selected library members are then shuffled by *in vitro* and/or *in vivo* recombinatorial shuffling as disclosed herein.

The invention also provides peptide libraries comprising a plurality of individual
5 library members of the invention, wherein (1) each individual library member of said plurality comprises a sequence produced by shuffling of a pool of selected sequences, and (2) each individual library member comprises a variable peptide segment sequence or single-chain antibody segment sequence which is distinct from the variable peptide segment sequences or single-chain antibody sequences of other individual library
10 members in said plurality (although some library members may be present in more than one copy per library due to uneven amplification, stochastic probability, or the like).

The invention also provides a product-by-process, wherein selected polynucleotide sequences having (or encoding a peptide having) a predetermined
15 binding specificity are formed by the process of: (1) screening a displayed peptide or displayed single-chain antibody library against a predetermined receptor (e.g., ligand) or epitope (e.g., antigen macromolecule) and identifying and/or enriching library members which bind to the predetermined receptor or epitope to produce a pool of selected library members, (2) shuffling by recombination the selected library members
20 (or amplified or cloned copies thereof) which binds the predetermined epitope and has been thereby isolated and/or enriched from the library to generate a shuffled library, and (3) screening the shuffled library against the predetermined receptor (e.g., ligand) or epitope (e.g., antigen macromolecule) and identifying and/or enriching shuffled library members which bind to the predetermined receptor or epitope to produce a pool of
25 selected shuffled library members.

Antibody Display and Screening Methods

The present method can be used to shuffle, by *in vitro* and/or *in vivo* recombination by any of the disclosed methods, and in any combination, polynucleotide sequences selected by antibody display methods, wherein an associated polynucleotide
5 encodes a displayed antibody which is screened for a phenotype (e.g., for affinity for binding a predetermined antigen (ligand)).

Various molecular genetic approaches have been devised to capture the vast immunological repertoire represented by the extremely large number of distinct variable
10 regions which can be present in immunoglobulin chains. The naturally-occurring germ line immunoglobulin heavy chain locus is composed of separate tandem arrays of variable segment genes located upstream of a tandem array of diversity segment genes, which are themselves located upstream of a tandem array of joining (i) region genes, which are located upstream of the constant region genes. During B lymphocyte
15 development, V-D-J rearrangement occurs wherein a heavy chain variable region gene (VH) is formed by rearrangement to form a fused D segment followed by rearrangement with a V segment to form a V-D-J joined product gene which, if productively rearranged, encodes a functional variable region (VH) of a heavy chain. Similarly, light chain loci rearrange one of several V segments with one of several J
20 segments to form a gene encoding the variable region (VL) of a light chain.

The vast repertoire of variable regions possible in immunoglobulins derives in part from the numerous combinatorial possibilities of joining V and i segments (and, in the case of heavy chain loci, D segments) during rearrangement in B cell development.
25 Additional sequence diversity in the heavy chain variable regions arises from non-uniform rearrangements of the D segments during V-D-J joining and from N region addition. Further, antigen-selection of specific B cell clones selects for higher affinity variants having non-germline mutations in one or both of the heavy and light chain variable regions; a phenomenon referred to as "affinity maturation" or "affinity
30 sharpening". Typically, these "affinity sharpening" mutations cluster in specific areas of

the variable region, most commonly in the complementarity-determining regions (CDRs).

In order to overcome many of the limitations in producing and identifying
5 high-affinity immunoglobulins through antigen-stimulated β cell development (i.e., immunization), various prokaryotic expression systems have been developed that can be manipulated to produce combinatorial antibody libraries which may be screened for high-affinity antibodies to specific antigens. Recent advances in the expression of antibodies in *Escherichia coli* and bacteriophage systems (see "alternative peptide
10 display methods", *infra*) have raised the possibility that virtually any specificity can be obtained by either cloning antibody genes from characterized hybridomas or by de novo selection using antibody gene libraries (e.g., from Ig cDNA).

Combinatorial libraries of antibodies have been generated in bacteriophage
15 lambda expression systems which may be screened as bacteriophage plaques or as colonies of lysogens (Huse et al, 1989); Caton and Koprowski, 1990; Mullinax et al, 1990; Persson et al, 1991). Various embodiments of bacteriophage antibody display libraries and lambda phage expression libraries have been described (Kang et al, 1991; Clackson et al, 1991; McCafferty et al, 1990; Burton et al, 1991; Hoogenboom et al,
20 1991; Chang et al, 1991; Breitling et al, 1991; Marks et al, 1991, p. 581; Barbas et al, 1992; Hawkins and Winter, 1992; Marks et al, 1992, p. 779; Marks et al, 1992, p. 16007; and Lowman et al, 1991; Lerner et al, 1992; all incorporated herein by reference). Typically, a bacteriophage antibody display library is screened with a receptor (e.g., polypeptide, carbohydrate, glycoprotein, nucleic acid) that is
25 immobilized (e.g., by covalent linkage to a chromatography resin to enrich for reactive phage by affinity chromatography) and/or labeled (e.g., to screen plaque or colony lifts).

One particularly advantageous approach has been the use of so-called single-chain fragment variable (scfv) libraries (Marks et al, 1992, p. 779; Winter and
30 Milstein, 1991; Clackson et al, 1991; Marks et al, 1991, p. 581; Chaudhary et al, 1990; Chiswell et al, 1992; McCafferty et al, 1990; and Huston et al, 1988). Various

embodiments of scfv libraries displayed on bacteriophage coat proteins have been described.

Beginning in 1988, single-chain analogues of Fv fragments and their fusion proteins have been reliably generated by antibody engineering methods. The first step generally involves obtaining the genes encoding VH and VL domains with desired binding properties; these V genes may be isolated from a specific hybridoma cell line, selected from a combinatorial V-gene library, or made by V gene synthesis. The single-chain Fv is formed by connecting the component V genes with an oligonucleotide that encodes an appropriately designed linker peptide, such as (Gly-Gly-Gly-Gly-Ser)₃ or equivalent linker peptide(s). The linker bridges the C-terminus of the first V region and N-terminus of the second, ordered as either VH-linker-VL or VL-linker-VH. In principle, the scfv binding site can faithfully replicate both the affinity and specificity of its parent antibody combining site.

Thus, scfv fragments are comprised of VH and VL domains linked into a single polypeptide chain by a flexible linker peptide. After the scfv genes are assembled, they are cloned into a phagemid and expressed at the tip of the M13 phage (or similar filamentous bacteriophage) as fusion proteins with the bacteriophage PIII (gene 3) coat protein. Enriching for phage expressing an antibody of interest is accomplished by panning the recombinant phage displaying a population scfv for binding to a predetermined epitope (e.g., target antigen, receptor).

The linked polynucleotide of a library member provides the basis for replication of the library member after a screening or selection procedure, and also provides the basis for the determination, by nucleotide sequencing, of the identity of the displayed peptide sequence or VH and VL amino acid sequence. The displayed peptide (s) or single-chain antibody (e. g., scfv) and/or its VH and VL domains or their CDRs can be cloned and expressed in a suitable expression system. Often polynucleotides encoding the isolated VH and VL domains will be ligated to polynucleotides encoding constant regions (CH and CL) to form polynucleotides encoding complete antibodies (e.g.,

chimeric or fully-human), antibody fragments, and the like. Often polynucleotides encoding the isolated CDRs will be grafted into polynucleotides encoding a suitable variable region framework (and optionally constant regions) to form polynucleotides encoding complete antibodies (e.g., humanized or fully-human), antibody fragments, and the like. Antibodies can be used to isolate preparative quantities of the antigen by immunoaffinity chromatography. Various other uses of such antibodies are to diagnose and/or stage disease (e.g., neoplasia) and for therapeutic application to treat disease, such as for example: neoplasia, autoimmune disease, AIDS, cardiovascular disease, infections, and the like.

10

Various methods have been reported for increasing the combinatorial diversity of a scfv library to broaden the repertoire of binding species (idiotype spectrum) The use of PCR has permitted the variable regions to be rapidly cloned either from a specific hybridoma source or as a gene library from non-immunized cells, affording combinatorial diversity in the assortment of VH and VL cassettes which can be combined. Furthermore, the VH and VL cassettes can themselves be diversified, such as by random, pseudorandom, or directed mutagenesis. Typically, VH and VL cassettes are diversified in or near the complementarity-determining regions (CDRS), often the third CDR, CDR3. Enzymatic inverse PCR mutagenesis has been shown to be a simple and reliable method for constructing relatively large libraries of scfv site-directed hybrids (Stemmer et al, 1993), as has error-prone PCR and chemical mutagenesis (Deng et al, 1994). Riechmann (Riechmann et al, 1993) showed semi-rational design of an antibody scfv fragment using site-directed randomization by degenerate oligonucleotide PCR and subsequent phage display of the resultant scfv hybrids. Barbas (Barbas et al, 1992) attempted to circumvent the problem of limited repertoire sizes resulting from using biased variable region sequences by randomizing the sequence in a synthetic CDR region of a human tetanus toxoid-binding Fab.

15

20

25

30

CDR randomization has the potential to create approximately 1×10^{20} CDRs for the heavy chain CDR3 alone, and a roughly similar number of variants of the heavy chain CDR1 and CDR2, and light chain CDR1-3 variants. Taken individually or

together, the combination possibilities of CDR randomization of heavy and/or light chains requires generating a prohibitive number of bacteriophage clones to produce a clone library representing all possible combinations, the vast majority of which will be non-binding. Generation of such large numbers of primary transformants is not feasible
5 with current transformation technology and bacteriophage display systems. For example, Barbas (**Barbas** et al, 1992) only generated 5×10^7 transformants, which represents only a tiny fraction of the potential diversity of a library of thoroughly randomized CDRS.

10 Despite these substantial limitations, bacteriophage display of scfv have already yielded a variety of useful antibodies and antibody fusion proteins. A bispecific single chain antibody has been shown to mediate efficient tumor cell lysis (**Gruber** et al, 1994). Intracellular expression of an anti-Rev scfv has been shown to inhibit HIV-1 virus replication *in vitro* (**Duan** et al, 1994), and intracellular expression of an
15 anti-p21rar, scfv has been shown to inhibit meiotic maturation of *Xenopus* oocytes (**Biocca** et al, 1993). Recombinant scfv which can be used to diagnose HIV infection have also been reported, demonstrating the diagnostic utility of scfv (**Lilley** et al, 1994). Fusion proteins wherein an scFv is linked to a second polypeptide, such as a toxin or fibrinolytic activator protein, have also been reported (**Holvost** et al, 1992; **Nicholls** et
20 al, 1993).

If it were possible to generate scfv libraries having broader antibody diversity and overcoming many of the limitations of conventional CDR mutagenesis and randomization methods which can cover only a very tiny fraction of the potential
25 sequence combinations, the number and quality of scfv antibodies suitable for therapeutic and diagnostic use could be vastly improved. To address this, the *in vitro* and *in vivo* shuffling methods of the invention are used to recombine CDRs which have been obtained (typically via PCR amplification or cloning) from nucleic acids obtained from selected displayed antibodies. Such displayed antibodies can be displayed on
30 cells, on bacteriophage particles, on polysomes, or any suitable antibody display system wherein the antibody is associated with its encoding nucleic acid(s). In a variation, the

CDRs are initially obtained from mRNA (or cDNA) from antibody-producing cells (e.g., plasma cells/splenocytes from an immunized wild-type mouse, a human, or a transgenic mouse capable of making a human antibody as in WO 92/03918, WO 93/12227, and WO 94/25585), including hybridomas derived therefrom.

5

Polynucleotide sequences selected in a first selection round (typically by affinity selection for displayed antibody binding to an antigen (e.g., a ligand) by any of these methods are pooled and the pool(s) is/are shuffled by *in vitro* and/or *in vivo* recombination, especially shuffling of CDRs (typically shuffling heavy chain CDRs with other heavy chain CDRs and light chain CDRs with other light chain CDRs) to produce a shuffled pool comprising a population of recombined selected polynucleotide sequences. The recombined selected polynucleotide sequences are expressed in a selection format as a displayed antibody and subjected to at least one subsequent selection round. The polynucleotide sequences selected in the subsequent selection round(s) can be used directly, sequenced, and/or subjected to one or more additional rounds of shuffling and subsequent selection until an antibody of the desired binding affinity is obtained. Selected sequences can also be back-crossed with polynucleotide sequences encoding neutral antibody framework sequences (i.e., having insubstantial functional effect on antigen binding), such as for example by back-crossing with a human variable region framework to produce human-like sequence antibodies. Generally, during back-crossing subsequent selection is applied to retain the property of binding to the predetermined antigen.

Alternatively, or in combination with the noted variations, the valency of the target epitope may be varied to control the average binding affinity of selected scfv library members. The target epitope can be bound to a surface or substrate at varying densities, such as by including a competitor epitope, by dilution, or by other method known to those in the art. A high density (valency) of predetermined epitope can be used to enrich for scfv library members which have relatively low affinity, whereas a low density (valency) can preferentially enrich for higher affinity scfv library members.

For generating diverse variable segments, a collection of synthetic oligonucleotides encoding random, pseudorandom, or a defined sequence kernel set of peptide sequences can be inserted by ligation into a predetermined site (e.g., a CDR). Similarly, the sequence diversity of one or more CDRs of the single-chain antibody cassette(s) can be expanded by mutating the CDR(s) with site-directed mutagenesis, CDR-replacement, and the like. The resultant DNA molecules can be propagated in a host for cloning and amplification prior to shuffling, or can be used directly (i.e., may avoid loss of diversity which may occur upon propagation in a host cell) and the selected library members subsequently shuffled.

10

Displayed peptide/polynucleotide complexes (library members) which encode a variable segment peptide sequence of interest or a single-chain antibody of interest are selected from the library by an affinity enrichment technique. This is accomplished by means of a immobilized macromolecule or epitope specific for the peptide sequence of interest, such as a receptor, other macromolecule, or other epitope species. Repeating the affinity selection procedure provides an enrichment of library members encoding the desired sequences, which may then be isolated for pooling and shuffling, for sequencing, and/or for further propagation and affinity enrichment.

20

The library members without the desired specificity are removed by washing. The degree and stringency of washing required will be determined for each peptide sequence or single-chain antibody of interest and the immobilized predetermined macromolecule or epitope. A certain degree of control can be exerted over the binding characteristics of the nascent peptide/DNA complexes recovered by adjusting the conditions of the binding incubation and the subsequent washing. The temperature, pH, ionic strength, divalent cations concentration, and the volume and duration of the washing will select for nascent peptide/DNA complexes within particular ranges of affinity for the immobilized macromolecule. Selection based on slow dissociation rate, which is usually predictive of high affinity, is often the most practical route. This may be done either by continued incubation in the presence of a saturating amount of free predetermined macromolecule, or by increasing the volume, number, and length of the

25
30

washes. In each case, the rebinding of dissociated nascent peptide/DNA or peptide/RNA complex is prevented, and with increasing time, nascent peptide/DNA or peptide/RNA complexes of higher and higher affinity are recovered.

5 Additional modifications of the binding and washing procedures may be applied to find peptides with special characteristics. The affinities of some peptides are dependent on ionic strength or cation concentration. This is a useful characteristic for peptides that will be used in affinity purification of various proteins when gentle conditions for removing the protein from the peptides are required.

10

One variation involves the use of multiple binding targets (multiple epitope species, multiple receptor species), such that a scfv library can be simultaneously screened for a multiplicity of scfv which have different binding specificities. Given that the size of a scfv library often limits the diversity of potential scfv sequences, it is typically desirable to use scfv libraries of as large a size as possible. The time and economic considerations of generating a number of very large polysome scFv-display libraries can become prohibitive. To avoid this substantial problem, multiple predetermined epitope species (receptor species) can be concomitantly screened in a single library, or sequential screening against a number of epitope species can be used.

15 In one variation, multiple target epitope species, each encoded on a separate bead (or subset of beads), can be mixed and incubated with a polysome-display scfv library under suitable binding conditions. The collection of beads, comprising multiple epitope species, can then be used to isolate, by affinity selection, scfv library members. Generally, subsequent affinity screening rounds can include the same mixture of beads, subsets thereof, or beads containing only one or two individual epitope species. This approach affords efficient screening, and is compatible with laboratory automation, batch processing, and high throughput screening methods.

20

25

A variety of techniques can be used in the present invention to diversify a peptide library or single-chain antibody library, or to diversify, prior to or concomitant with shuffling, around variable segment peptides found in early rounds of panning to

30

have sufficient binding activity to the predetermined macromolecule or epitope. In one approach, the positive selected peptide/polynucleotide complexes (those identified in an early round of affinity enrichment) are sequenced to determine the identity of the active peptides. Oligonucleotides are then synthesized based on these active peptide
5 sequences, employing a low level of all bases incorporated at each step to produce slight variations of the primary oligonucleotide sequences. This mixture of (slightly) degenerate oligonucleotides is then cloned into the variable segment sequences at the appropriate locations. This method produces systematic, controlled variations of the starting peptide sequences, which can then be shuffled. It requires, however, that
10 individual positive nascent peptide/polynucleotide complexes be sequenced before mutagenesis, and thus is useful for expanding the diversity of small numbers of recovered complexes and selecting variants having higher binding affinity and/or higher binding specificity. In a variation, mutagenic PCR amplification of positive selected peptide/polynucleotide complexes (especially of the variable region sequences, the
15 amplification products of which are shuffled *in vitro* and/or *in vivo* and one or more additional rounds of screening is done prior to sequencing. The same general approach can be employed with single-chain antibodies in order to expand the diversity and enhance the binding affinity/specificity, typically by diversifying CDRs or adjacent framework regions prior to or concomitant with shuffling. If desired, shuffling
20 reactions can be spiked with mutagenic oligonucleotides capable of *in vitro* recombination with the selected library members can be included. Thus, mixtures of synthetic oligonucleotides and PCR produced polynucleotides (synthesized by error-prone or high-fidelity methods) can be added to the *in vitro* shuffling mix and be incorporated into resulting shuffled library members (shufflants).

25

The present invention of shuffling enables the generation of a vast library of CDR-variant single-chain antibodies. One way to generate such antibodies is to insert synthetic CDRs into the single-chain antibody and/or CDR randomization prior to or concomitant with shuffling. The sequences of the synthetic CDR cassettes are selected
30 by referring to known sequence data of human CDR and are selected in the discretion of the practitioner according to the following guidelines: synthetic CDRs will have at least

40 percent positional sequence identity to known CDR sequences, and preferably will have at least 50 to 70 percent positional sequence identity to known CDR sequences. For example, a collection of synthetic CDR sequences can be generated by synthesizing a collection of oligonucleotide sequences on the basis of naturally-occurring human CDR sequences listed in Kabat (Kabat et al, 1991); the pool (s) of synthetic CDR sequences are calculated to encode CDR peptide sequences having at least 40 percent sequence identity to at least one known naturally-occurring human CDR sequence. Alternatively, a collection of naturally-occurring CDR sequences may be compared to generate consensus sequences so that amino acids used at a residue position frequently (i.e., in at least 5 percent of known CDR sequences) are incorporated into the synthetic CDRs at the corresponding position(s). Typically, several (e.g., 3 to about 50) known CDR sequences are compared and observed natural sequence variations between the known CDRs are tabulated, and a collection of oligonucleotides encoding CDR peptide sequences encompassing all or most permutations of the observed natural sequence variations is synthesized. For example but not for limitation, if a collection of human VH CDR sequences have carboxy-terminal amino acids which are either Tyr, Val, Phe, or Asp, then the pool(s) of synthetic CDR oligonucleotide sequences are designed to allow the carboxy-terminal CDR residue to be any of these amino acids. In some embodiments, residues other than those which naturally-occur at a residue position in the collection of CDR sequences are incorporated: conservative amino acid substitutions are frequently incorporated and up to 5 residue positions may be varied to incorporate non-conservative amino acid substitutions as compared to known naturally-occurring CDR sequences. Such CDR sequences can be used in primary library members (prior to first round screening) and/or can be used to spike *in vitro* shuffling reactions of selected library member sequences. Construction of such pools of defined and/or degenerate sequences will be readily accomplished by those of ordinary skill in the art.

The collection of synthetic CDR sequences comprises at least one member that is not known to be a naturally-occurring CDR sequence. It is within the discretion of the practitioner to include or not include a portion of random or pseudorandom

sequence corresponding to N region addition in the heavy chain CDR; the N region sequence ranges from 1 nucleotide to about 4 nucleotides occurring at V-D and D-J junctions. A collection of synthetic heavy chain CDR sequences comprises at least about 100 unique CDR sequences, typically at least about 1,000 unique CDR sequences, preferably at least about 10,000 unique CDR sequences, frequently more than 50,000 unique CDR sequences; however, usually not more than about 1×10^6 unique CDR sequences are included in the collection, although occasionally 1×10^7 to 1×10^8 unique CDR sequences are present, especially if conservative amino acid substitutions are permitted at positions where the conservative amino acid substituent is not present or is rare (i.e., less than 0.1 percent) in that position in naturally-occurring human CDRS. In general, the number of unique CDR sequences included in a library should not exceed the expected number of primary transformants in the library by more than a factor of 10. Such single-chain antibodies generally bind of about at least 1×10^6 M⁻¹, preferably with an affinity of about at least 5×10^7 M⁻¹, more preferably with an affinity of at least 1×10^8 M⁻¹ to 1×10^9 M⁻¹ or more, sometimes up to 1×10^{10} M⁻¹ or more. Frequently, the predetermined antigen is a human protein, such as for example a human cell surface antigen (e. g., CD4, CD8, IL-2 receptor, EGF receptor, PDGF receptor), other human biological macromolecule (e.g., thrombomodulin, protein C, carbohydrate antigen, sialyl Lewis antigen, Lselectin), or nonhuman disease associated macromolecule (e.g., bacterial LPS, virion capsid protein or envelope glycoprotein) and the like.

High affinity single-chain antibodies of the desired specificity can be engineered and expressed in a variety of systems. For example, scfv have been produced in plants (Firek et al, 1993) and can be readily made in prokaryotic systems (Owens and Young, 1994; Johnson and Bird, 1991). Furthermore, the single-chain antibodies can be used as a basis for constructing whole antibodies or various fragments thereof (Kettleborough et al, 1994). The variable region encoding sequence may be isolated (e.g., by PCR amplification or subcloning) and spliced to a sequence encoding a desired human constant region to encode a human sequence antibody more suitable for human therapeutic uses where immunogenicity is preferably minimized. The polynucleotide(s)

having the resultant fully human encoding sequence(s) can be expressed in a host cell (e.g., from an expression vector in a mammalian cell) and purified for pharmaceutical formulation.

5 The DNA expression constructs will typically include an expression control DNA sequence operably linked to the coding sequences, including naturally-associated or heterologous promoter regions. Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells. Once the vector has been incorporated into the appropriate host,
10 the host is maintained under conditions suitable for high level expression of the nucleotide sequences, and the collection and purification of the mutant "engineered" antibodies.

 As stated previously, the DNA sequences will be expressed in hosts after the
15 sequences have been operably linked to an expression control sequence (i.e., positioned to ensure the transcription and translation of the structural gene). These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. Commonly, expression vectors will contain selection markers, e.g., tetracycline or neomycin, to permit detection of those cells
20 transformed with the desired DNA sequences (see, e.g., **USPN 4,704,362**, which is incorporated herein by reference).

 In addition to eukaryotic microorganisms such as yeast, mammalian tissue cell culture may also be used to produce the polypeptides of the present invention (see
25 **Winnacker, 1987**), which is incorporated herein by reference). Eukaryotic cells are actually preferred, because a number of suitable host cell lines capable of secreting intact immunoglobulins have been developed in the art, and include the CHO cell lines, various COS cell lines, HeLa cells, and myeloma cell lines, but preferably transformed Bcells or hybridomas. Expression vectors for these cells can include expression control
30 sequences, such as an origin of replication, a promoter, an enhancer (**Queen et al, 1986**), and necessary processing information sites, such as ribosome binding sites, RNA

splice sites, polyadenylation sites, and transcriptional terminator sequences. Preferred expression control sequences are promoters derived from immunoglobulin genes, cytomegalovirus, SV40, Adenovirus, Bovine Papilloma Virus, and the like.

5 Eukaryotic DNA transcription can be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting sequences of between 10 to 300 bp that increase transcription by a promoter. Enhancers can effectively increase transcription when either 5' or 3' to the transcription unit. They are also effective if located within an intron or within the coding sequence itself. Typically, viral enhancers
10 are used, including SV40 enhancers, cytomegalovirus enhancers, polyoma enhancers, and adenovirus enhancers. Enhancer sequences from mammalian systems are also commonly used, such as the mouse immunoglobulin heavy chain enhancer.

Mammalian expression vector systems will also typically include a selectable
15 marker gene. Examples of suitable markers include, the dihydrofolate reductase gene (DHFR), the thymidine kinase gene (TK), or prokaryotic genes conferring drug resistance. The first two marker genes prefer the use of mutant cell lines that lack the ability to grow without the addition of thymidine to the growth medium. Transformed cells can then be identified by their ability to grow on non-supplemented media.
20 Examples of prokaryotic drug resistance genes useful as markers include genes conferring resistance to G418, mycophenolic acid and hygromycin.

The vectors containing the DNA segments of interest can be transferred into the host cell by well-known methods, depending on the type of cellular host. For example,
25 calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment, lipofection, or electroporation may be used for other cellular hosts. Other methods used to transform mammalian cells include the use of Polybrene, protoplast fusion, liposomes, electroporation, and micro-injection (see, generally, Sambrook et al, 1982 and 1989).

30

Once expressed, the antibodies, individual mutated immunoglobulin chains, mutated antibody fragments, and other immunoglobulin polypeptides of the invention can be purified according to standard procedures of the art, including ammonium sulfate precipitation, fraction column chromatography, gel electrophoresis and the like (see, generally, Scopes, 1982). Once purified, partially or to homogeneity as desired, the polypeptides may then be used therapeutically or in developing and performing assay procedures, immunofluorescent stainings, and the like (see, generally, **Lefkovits and Pernis, 1979 and 1981; Lefkovits, 1997**).

10 The antibodies generated by the method of the present invention can be used for diagnosis and therapy. By way of illustration and not limitation, they can be used to treat cancer, autoimmune diseases, or viral infections. For treatment of cancer, the antibodies will typically bind to an antigen expressed preferentially on cancer cells, such as erbB-2, CEA, CD33, and many other antigens and binding members well
15 known to those skilled in the art.

End-Selection

This invention provides a method for selecting a subset of polynucleotides from a starting set of polynucleotides, which method is based on the ability to discriminate one or more selectable features (or selection markers) present anywhere in a working
5 polynucleotide, so as to allow one to perform selection for (positive selection) &/or against (negative selection) each selectable polynucleotide. In a preferred aspect, a method is provided termed end-selection, which method is based on the use of a selection marker located in part or entirely in a terminal region of a selectable polynucleotide, and such a selection marker may be termed an "end-selection marker".

10

End-selection may be based on detection of naturally occurring sequences or on detection of sequences introduced experimentally (including by any mutagenesis procedure mentioned herein and not mentioned herein) or on both, even within the same polynucleotide. An end-selection marker can be a structural selection marker or a
15 functional selection marker or both a structural and a functional selection marker. An end-selection marker may be comprised of a polynucleotide sequence or of a polypeptide sequence or of any chemical structure or of any biological or biochemical tag, including markers that can be selected using methods based on the detection of radioactivity, of enzymatic activity, of fluorescence, of any optical feature, of a
20 magnetic property (e.g. using magnetic beads), of immunoreactivity, and of hybridization.

End-selection may be applied in combination with any method serviceable for performing mutagenesis. Such mutagenesis methods include, but are not limited to,
25 methods described herein (supra and infra). Such methods include, by way of non-limiting exemplification, any method that may be referred herein or by others in the art by any of the following terms: "saturation mutagenesis", "shuffling", "recombination", "re-assembly", "error-prone PCR", "assembly PCR", "sexual PCR", "crossover PCR", "oligonucleotide primer-directed mutagenesis", "recursive (&/or exponential) ensemble
30 mutagenesis (see Arkin and Youvan, 1992)", "cassette mutagenesis", "in vivo mutagenesis", and "in vitro mutagenesis". Moreover, end-selection may be performed

on molecules produced by any mutagenesis &/or amplification method (see, e.g., Arn Id, 1993; Caldwell and Joyce, 1992; Stemmer, 1994; following which method it is desirable to select for (including to screen for the presence of) desirable progeny molecules.

5

In addition, end-selection may be applied to a polynucleotide apart from any mutagenesis method. In a preferred embodiment, end-selection, as provided herein, can be used in order to facilitate a cloning step, such as a step of ligation to another polynucleotide (including ligation to a vector). This invention thus provides for end-
10 selection as a serviceable means to facilitate library construction, selection &/or enrichment for desirable polynucleotides, and cloning in general.

In a particularly preferred embodiment, end-selection can be based on (positive) selection for a polynucleotide; alternatively end-selection can be based on (negative)
15 selection against a polynucleotide; and alternatively still, end-selection can be based on both (positive) selection for, and on (negative) selection against, a polynucleotide. End-selection, along with other methods of selection &/or screening, can be performed in an iterative fashion, with any combination of like or unlike selection &/or screening methods and serviceable mutagenesis methods, all of which can be performed in an
20 iterative fashion and in any order, combination, and permutation.

It is also appreciated that, according to one embodiment of this invention, end-selection may also be used to select a polynucleotide is at least in part: circular (e.g. a plasmid or any other circular vector or any other polynucleotide that is partly circular),
25 &/or branched, &/or modified or substituted with any chemical group or moiety. In accord with this embodiment, a polynucleotide may be a circular molecule comprised of an intermediate or central region, which region is flanked on a 5' side by a 5' flanking region (which, for the purpose of end-selection, serves in like manner to a 5' terminal region of a non-circular polynucleotide) and on a 3' side by a 3' terminal region (which,
30 for the purpose of end-selection, serves in like manner to a 3' terminal region of a non-

circular polynucleotide). As used in this non-limiting exemplification, there may be sequence overlap between any two regions or even among all three regions.

In one non-limiting aspect of this invention, end-selection of a linear
5 polynucleotide is performed using a general approach based on the presence of at least one end-selection marker located at or near a polynucleotide end or terminus (that can be either a 5' end or a 3' end). In one particular non-limiting exemplification, end-selection is based on selection for a specific sequence at or near a terminus such as, but not limited to, a sequence recognized by an enzyme that recognizes a polynucleotide
10 sequence. An enzyme that recognizes and catalyzes a chemical modification of a polynucleotide is referred to herein as a polynucleotide-acting enzyme. In a preferred embodiment, serviceable polynucleotide-acting enzymes are exemplified non-exclusively by enzymes with polynucleotide-cleaving activity, enzymes with polynucleotide-methylating activity, enzymes with polynucleotide-ligating activity, and
15 enzymes with a plurality of distinguishable enzymatic activities (including non-exclusively, e.g., both polynucleotide-cleaving activity and polynucleotide-ligating activity).

Relevant polynucleotide-acting enzymes thus also include any commercially
20 available or non-commercially available polynucleotide endonucleases and their companion methylases including those catalogued at the website <http://www.neb.com/rebase>, and those mentioned in the following cited reference (Roberts and Macelis, 1996). Preferred polynucleotide endonucleases include – but are not limited to – type II restriction enzymes (including type IIS), and include
25 enzymes that cleave both strands of a double stranded polynucleotide (e.g. *Not* I, which cleaves both strands at 5'...GC/GGCCGC...3') and enzymes that cleave only one strand of a double stranded polynucleotide, i.e. enzymes that have polynucleotide-nicking activity, (e.g. *N. Bst*NB I, which cleaves only one strand at 5'...GAGTCNNNN/N...3'). Relevant polynucleotide-acting enzymes also include type
30 III restriction enzymes.

It is appreciated that relevant polynucleotide-acting enzymes also include any enzymes that may be developed in the future, though currently unavailable, that are serviceable for generating a ligation compatible end, preferably a sticky end, in a polynucleotide.

5

In one preferred exemplification, a serviceable selection marker is a restriction site in a polynucleotide that allows a corresponding type II (or type IIS) restriction enzyme to cleave an end of the polynucleotide so as to provide a ligatable end (including a blunt end or alternatively a sticky end with at least a one base overhang) that is serviceable for a desirable ligation reaction without cleaving the polynucleotide internally in a manner that destroys a desired internal sequence in the polynucleotide. Thus it is provided that, among relevant restriction sites, those sites that do not occur internally (i.e. that do not occur apart from the termini) in a specific working polynucleotide are preferred when the use of a corresponding restriction enzyme(s) is not intended to cut the working polynucleotide internally. This allows one to perform restriction digestion reactions to completion or to near completion without incurring unwanted internal cleavage in a working polynucleotide.

According to a preferred aspect, it is thus preferable to use restriction sites that are not contained, or alternatively that are not expected to be contained, or alternatively that unlikely to be contained (e.g. when sequence information regarding a working polynucleotide is incomplete) internally in a polynucleotide to be subjected to end-selection. In accordance with this aspect, it is appreciated that restriction sites that occur relatively infrequently are usually preferred over those that occur more frequently. On the other hand it is also appreciated that there are occasions where internal cleavage of a polypeptide is desired, e.g. to achieve recombination or other mutagenic procedures along with end-selection.

In accord with this invention, it is also appreciated that methods (e.g. mutagenesis methods) can be used to remove unwanted internal restriction sites. It is also appreciated that a partial digestion reaction (i.e. a digestion reaction that proceeds

to partial completion) can be used to achieve digestion at a recognition site in a terminal region while sparing a susceptible restriction site that occurs internally in a polynucleotide and that is recognized by the same enzyme. In one aspect, partial digest are useful because it is appreciated that certain enzymes show preferential cleavage of the same recognition sequence depending on the location and environment in which the recognition sequence occurs. For example, it is appreciated that, while lambda DNA has 5 *EcoR* I sites, cleavage of the site nearest to the right terminus has been reported to occur 10 times faster than the sites in the middle of the molecule. Also, for example, it has been reported that, while *Sac* II has four sites on lambda DNA, the three clustered centrally in lambda are cleaved 50 times faster than the remaining site near the terminus (at nucleotide 40,386). Summarily, site preferences have been reported for various enzymes by many investigators (e.g., **Thomas and Davis**, 1975; **Forsblum et al**, 1976; **Nath and Azzolina**, 1981; **Brown and Smith**, 1977; **Gingeras and Brooks**, 1983; **Krüger et al**, 1988; **Conrad and Topal**, 1989; **Oller et al**, 1991; **Topal**, 1991; and **Pein**, 1991; to name but a few). It is appreciated that any empirical observations as well as any mechanistic understandings of site preferences by any serviceable polynucleotide-acting enzymes, whether currently available or to be procured in the future, may be serviceable in end-selection according to this invention.

It is also appreciated that protection methods can be used to selectively protect specified restriction sites (e.g. internal sites) against unwanted digestion by enzymes that would otherwise cut a working polypeptide in response to the presence of those sites; and that such protection methods include modifications such as methylations and base substitutions (e.g. U instead of T) that inhibit an unwanted enzyme activity. It is appreciated that there are limited numbers of available restriction enzymes that are rare enough (e.g. having very long recognition sequences) to create large (e.g. megabase-long) restriction fragments, and that protection approaches (e.g. by methylation) are serviceable for increasing the rarity of enzyme cleavage sites. The use of *M.Fnu* II (mCGCG) to increase the apparent rarity of *Not* I approximately twofold is but one example among many (**Qiang et al**, 1990; **Nelson et al**, 1984; **Maxam and Gilbert**, 1980; **Raleigh and Wilson**, 1986).

According to a preferred aspect of this invention, it is provided that, in general, the use of rare restriction sites is preferred. It is appreciated that, in general, the frequency of occurrence of a restriction site is determined by the number of nucleotides contained therein, as well as by the ambiguity of the base requirements contained therein. Thus, in a non-limiting exemplification, it is appreciated that, in general, a restriction site composed of, for example, 8 specific nucleotides (e.g. the *Not* I site or GC/GGCCGC, with an estimated relative occurrence of 1 in 4^8 , i.e. 1 in 65,536, random 8-mers) is relatively more infrequent than one composed of, for example, 6 nucleotides (e.g. the *Sma* I site or CCC/GGG, having an estimated relative occurrence of 1 in 4^6 , i.e. 1 in 4,096, random 6-mers), which in turn is relatively more infrequent than one composed of, for example, 4 nucleotides (e.g. the *Msp* I site or C/CGG, having an estimated relative occurrence of 1 in 4^4 , i.e. 1 in 256, random 4-mers). Moreover, in another non-limiting exemplification, it is appreciated that, in general, a restriction site having no ambiguous (but only specific) base requirements (e.g. the *Fin* I site or GTCCC, having an estimated relative occurrence of 1 in 4^5 , i.e. 1 in 1024, random 5-mers) is relatively more infrequent than one having an ambiguous W (where W = A or T) base requirement (e.g. the *Ava* II site or G/GWCC, having an estimated relative occurrence of 1 in $4 \times 4 \times 2 \times 4 \times 4$ - i.e. 1 in 512 - random 5-mers), which in turn is relatively more infrequent than one having an ambiguous N (where N = A or C or G or T) base requirement (e.g. the *Asu* I site or G/GNCC, having an estimated relative occurrence of 1 in $4 \times 4 \times 1 \times 4 \times 4$, i.e. 1 in 256 - random 5-mers). These relative occurrences are considered general estimates for actual polynucleotides, because it is appreciated that specific nucleotide bases (not to mention specific nucleotide sequences) occur with dissimilar frequencies in specific polynucleotides, in specific species of organisms, and in specific groupings of organisms. For example, it is appreciated that the % G+C contents of different species of organisms are often very different and wide ranging.

The use of relatively more infrequent restriction sites as a selection marker include - in a non-limiting fashion - preferably those sites composed at least a 4

nucleotide sequence, more preferably those composed at least a 5 nucleotide sequence, more preferably still those composed at least a 6 nucleotide sequence (e.g. the *Bam*H I site or G/GATCC, the *Bgl* II site or A/GATCT, the *Pst* I site or CTGCA/G, and the *Xba* I site or T/CTAGA), more preferably still those composed at least a 7 nucleotide
5 sequence, more preferably still those composed of an 8 nucleotide sequence nucleotide sequence (e.g. the *Asc* I site or GG/CGCGCC, the *Not* I site or GC/GGCCGC, the *Pac* I site or TTAAT/TAA, the *Pme* I site or GTTT/AAAC, the *Srf* I site or GCCC/GGGC, the *Sse*838 I site or CCTGCA/GG, and the *Swa* I site or ATTT/AAAT), more preferably still those composed of a 9 nucleotide sequence, and even more preferably still those
10 composed of at least a 10 nucleotide sequence (e.g. the *Bsp*G I site or CG/CGCTGGAC). It is further appreciated that some restriction sites (e.g. for class IIS enzymes) are comprised of a portion of relatively high specificity (i.e. a portion containing a principal determinant of the frequency of occurrence of the restriction site) and a portion of relatively low specificity; and that a site of cleavage may or may not be
15 contained within a portion of relatively low specificity. For example, in the *Eco*57 I site or CTGAAG(16/14), there is a portion of relatively high specificity (i.e. the CTGAAG portion) and a portion of relatively low specificity (i.e. the N16 sequence) that contains a site of cleavage.

20 In another preferred embodiment of this invention, a serviceable end-selection marker is a terminal sequence that is recognized by a polynucleotide-acting enzyme that recognizes a specific polynucleotide sequence. In a preferred aspect of this invention, serviceable polynucleotide-acting enzymes also include other enzymes in addition to classic type II restriction enzymes. According to this preferred aspect of this invention,
25 serviceable polynucleotide-acting enzymes also include gyrases, helicases, recombinases, relaxases, and any enzymes related thereto.

Among preferred examples are topoisomerases (which have been categorized by some as a subset of the gyrases) and any other enzymes that have polynucleotide-
30 cleaving activity (including preferably polynucleotide-nicking activity) &/or polynucleotide-ligating activity. Among preferred topoisomerase enzymes are

topoisomerase I enzymes, which is available from many commercial sources (Epicentre Technologies, Madison, WI; Invitrogen, Carlsbad, CA; Life Technologies, Gathesburg, MD) and conceivably even more private sources. It is appreciated that similar enzymes may be developed in the future that are serviceable for end-selection as provided herein.

- 5 A particularly preferred topoisomerase I enzyme is a topoisomerase I enzyme of vaccinia virus origin, that has a specific recognition sequence (e.g. 5'...AAGGG...3') and has both polynucleotide-nicking activity and polynucleotide-ligating activity. Due to the specific nicking-activity of this enzyme (cleavage of one strand), internal recognition sites are not prone to polynucleotide destruction resulting from the nicking
- 10 activity (but rather remain annealed) at a temperature that causes denaturation of a terminal site that has been nicked. Thus for use in end-selection, it is preferable that a nicking site for topoisomerase-based end-selection be no more than 100 nucleotides from a terminus, more preferably no more than 50 nucleotides from a terminus, more preferably still no more than 25 nucleotides from a terminus, even more preferably still
- 15 no more than 20 nucleotides from a terminus, even more preferably still no more than 15 nucleotides from a terminus, even more preferably still no more than 10 nucleotides from a terminus, even more preferably still no more than 8 nucleotides from a terminus, even more preferably still no more than 6 nucleotides from a terminus, and even more preferably still no more than 4 nucleotides from a terminus.

20

- In a particularly preferred exemplification that is non-limiting yet clearly illustrative, it is appreciated that when a nicking site for topoisomerase-based end-selection is 4 nucleotides from a terminus, nicking produces a single stranded oligo of 4 bases (in a terminal region) that can be denatured from its complementary strand in an
- 25 end-selectable polynucleotide; this provides a sticky end (comprised of 4 bases) in a polynucleotide that is serviceable for an ensuing ligation reaction. To accomplish ligation to a cloning vector (preferably an expression vector), compatible sticky ends can be generated in a cloning vector by any means including by restriction enzyme-based means. The terminal nucleotides (comprised of 4 terminal bases in this specific
- 30 example) in an end-selectable polynucleotide terminus are thus wisely chosen to

provide compatibility with a sticky end generated in a cloning vector to which the polynucleotide is to be ligated.

On the other hand, internal nicking of an end-selectable polynucleotide, e.g. 500
5 bases from a terminus, produces a single stranded oligo of 500 bases that is not easily denatured from its complementary strand, but rather is serviceable for repair (e.g. by the same topoisomerase enzyme that produced the nick).

This invention thus provides a method - e.g. that is vaccinia topoisomerase-
10 based &/or type II (or IIS) restriction endonuclease-based &/or type III restriction endonuclease-based &/or nicking enzyme-based (e.g. using N. *Bst*NB I) - for producing a sticky end in a working polynucleotide, which end is ligation compatible, and which end can be comprised of at least a 1 base overhang. Preferably such a sticky end is comprised of at least a 2-base overhang, more preferably such a sticky end is comprised
15 of at least a 3-base overhang, more preferably still such a sticky end is comprised of at least a 4-base overhang, even more preferably still such a sticky end is comprised of at least a 5-base overhang, even more preferably still such a sticky end is comprised of at least a 6-base overhang. Such a sticky end may also be comprised of at least a 7-base overhang, or at least an 8-base overhang, or at least a 9-base overhang, or at least a 10-
20 base overhang, or at least 15-base overhang, or at least a 20-base overhang, or at least a 25-base overhang, or at least a 30-base overhang. These overhangs can be comprised of any bases, including A, C, G, or T.

It is appreciated that sticky end overhangs introduced using topoisomerase or a
25 nicking enzyme (e.g. using N. *Bst*NB I) can be designed to be unique in a ligation environment, so as to prevent unwanted fragment reassemblies, such as self-dimerizations and other unwanted concatamerizations.

According to one aspect of this invention, a plurality of sequences (which may
30 but do not necessarily overlap) can be introduced into a terminal region of an end-selectable polynucleotide by the use of an oligo in a polymerase-based reaction. In a

relevant, but by no means limiting example, such an oligo can be used to provide a preferred 5' terminal region that is serviceable for topoisomerase I-based end-selection, which oligo is comprised of: a 1-10 base sequence that is convertible into a sticky end (preferably by a vaccinia topoisomerase I), a ribosome binding site (i.e. and "RBS", that is preferably serviceable for expression cloning), and optional linker sequence followed by an ATG start site and a template-specific sequence of 0-100 bases (to facilitate annealment to the template in the a polymerase-based reaction). Thus, according to this example, a serviceable oligo (which may be termed a forward primer) can have the sequence: 5'[terminal sequence = (N)₁₋₁₀][topoisomerase I site & RBS =

5 AAGGGAGGAG][linker = (N)₁₋₁₀₀][start codon and template-specific sequence =

10 ATG(N)₀₋₁₀₀]3'.

Analogously, in a relevant, but by no means limiting example, an oligo can be used to provide a preferred 3' terminal region that is serviceable for topoisomerase I-based end-selection, which oligo is comprised of: a 1-10 base sequence that is convertible into a sticky end (preferably by a vaccinia topoisomerase I), and optional linker sequence followed by a template-specific sequence of 0-100 bases (to facilitate annealment to the template in the a polymerase-based reaction). Thus, according to this example, a serviceable oligo (which may be termed a reverse primer) can have the

15 sequence: 5'[terminal sequence = (N)₁₋₁₀][topoisomerase I site = AAGGG][linker =

20 (N)₁₋₁₀₀][template-specific sequence = (N)₀₋₁₀₀]3'.

It is appreciated that, end-selection can be used to distinguish and separate parental template molecules (e.g. to be subjected to mutagenesis) from progeny molecules (e.g. generated by mutagenesis). For example, a first set of primers, lacking in a topoisomerase I recognition site, can be used to modify the terminal regions of the parental molecules (e.g. in polymerase-based amplification). A different second set of primers (e.g. having a topoisomerase I recognition site) can then be used to generate mutated progeny molecules (e.g. using any polynucleotide chimerization method, such as interrupted synthesis,

25 template-switching polymerase-based amplification, or interrupted synthesis; or using saturation mutagenesis; or using any other method for introducing a topoisomerase I

30

recognition site into a mutagenized progeny molecule as disclosed herein) from the amplified template molecules. The use of topoisomerase I-based end-selection can then facilitate, not only discernment, but selective topoisomerase I-based ligation of the desired progeny molecules.

5

Annealment of a second set of primers to thusly amplified parental molecules can be facilitated by including sequences in a first set of primers (i.e. primers used for amplifying a set parental molecules) that are similar to a topoisomerase I recognition site, yet different enough to prevent functional topoisomerase I enzyme recognition. For example, sequences
10 that diverge from the AAGGG site by anywhere from 1 base to all 5 bases can be incorporated into a first set of primers (to be used for amplifying the parental templates prior to subjection to mutagenesis). In a specific, but non-limiting aspect, it is thus provided that a parental molecule can be amplified using the following exemplary – but by no means limiting – set of forward and reverse primers:

15

Forward Primer: 5' CTAGAAGAGAGGAGAAAACCATG(N)₁₀₋₁₀₀ 3', and
Reverse Primer: 5' GATCAAAGGCGCGCCTGCAGG(N)₁₀₋₁₀₀ 3'

According to this specific example of a first set of primers, (N)₁₀₋₁₀₀ represents
20 preferably a 10 to 100 nucleotide-long template-specific sequence, more preferably a 10 to 50 nucleotide-long template-specific sequence, more preferably still a 10 to 30 nucleotide-long template-specific sequence, and even more preferably still a 15 to 25 nucleotide-long template-specific sequence.

25 According to a specific, but non-limiting aspect, it is thus provided that, after this amplification (using a disclosed first set of primers lacking in a true topoisomerase I recognition site), amplified parental molecules can then be subjected to mutagenesis using one or more sets of forward and reverse primers that do have a true topoisomerase I recognition site. In a specific, but non-limiting aspect, it is thus provided that a parental
30 molecule can be used as templates for the generation of a mutagenized progeny molecule

using the following exemplary – but by no means limiting – second set of forward and reverse primers:

Forward Primer: 5' CTAGAAGGGAGGAGAAAACCATG 3'

5 Reverse Primer: 5' GATCAAAGGCGCGCCTGCAGG 3' (contains *Asc* I recognition sequence)

It is appreciated that any number of different primers sets not specifically mentioned can be used as first, second, or subsequent sets of primers for end-selection consistent with
10 this invention. Notice that type II restriction enzyme sites can be incorporated (e.g. an *Asc* I site in the above example). It is provided that, in addition to the other sequences mentioned, the experimentalist can incorporate one or more N,N,G/T triplets into a serviceable primer in order to subject a working polynucleotide to saturation mutagenesis. Summarily, use of a second and/or subsequent set of primers can achieve dual goals of introducing a
15 topoisomerase I site and of generating mutations in a progeny polynucleotide.

Thus, according to one use provided, a serviceable end-selection marker is an enzyme recognition site that allows an enzyme to cleave (including nick) a polynucleotide at a specified site, to produce a ligation-compatible end upon
20 denaturation of a generated single stranded oligo. Ligation of the produced polynucleotide end can then be accomplished by the same enzyme (e.g. in the case of vaccinia virus topoisomerase I), or alternatively with the use of a different enzyme. According to one aspect of this invention, any serviceable end-selection markers, whether like (e.g. two vaccinia virus topoisomerase I recognition sites) or unlike (e.g. a
25 class II restriction enzyme recognition site and a vaccinia virus topoisomerase I recognition site) can be used in combination to select a polynucleotide. Each selectable polynucleotide can thus have one or more end-selection markers, and they can be like or unlike end-selection markers. In a particular aspect, a plurality of end-selection markers can be located on one end of a polynucleotide and can have overlapping sequences with
30 each other.

It is important to emphasize that any number of enzymes, whether currently in existence or to be developed, can be serviceable in end-selection according to this invention. For example, in a particular aspect of this invention, a nicking enzyme (e.g. *N. Bst*NB I, which cleaves only one strand at 5'...GAGTCNNNN/N...3') can be used in conjunction with a source of polynucleotide-ligating activity in order to achieve end-selection. According to this embodiment, a recognition site for *N. Bst*N BI – instead of a recognition site for topoisomerase I – should be incorporated into an end-selectable polynucleotide (whether end-selection is used for selection of a mutagenized progeny molecule or whether end-selection is used apart from any mutagenesis procedure).

10

It is appreciated that the instantly disclosed end-selection approach using topoisomerase-based nicking and ligation has several advantages over previously available selection methods. In sum, this approach allows one to achieve direction cloning (including expression cloning). Specifically, this approach can be used for the achievement of: direct ligation (i.e. without subjection to a classic restriction-purification-ligation reaction, that is susceptible to a multitude of potential problems from an initial restriction reaction to a ligation reaction dependent on the use of T4 DNA ligase); separation of progeny molecules from original template molecules (e.g. original template molecules lack topoisomerase I sites that not introduced until after mutagenesis), obviation of the need for size separation steps (e.g. by gel chromatography or by other electrophoretic means or by the use of size-exclusion membranes), preservation of internal sequences (even when topoisomerase I sites are present), obviation of concerns about unsuccessful ligation reactions (e.g. dependent on the use of T4 DNA ligase, particularly in the presence of unwanted residual restriction enzyme activity), and facilitated expression cloning (including obviation of frame shift concerns). Concerns about unwanted restriction enzyme-based cleavages – especially at internal restriction sites (or even at often unpredictable sites of unwanted star activity) in a working polynucleotide – that are potential sites of destruction of a working polynucleotide can also be obviated by the instantly disclosed end-selection approach using topoisomerase-based nicking and ligation.

30

Two-Hybrid Based Screening Assays

Shuffling can also be used to recombinatorially diversify a pool of selected library members obtained by screening a two-hybrid screening system to identify library members which bind a predetermined polypeptide sequence. The selected library members are pooled and shuffled by *in vitro* and/or *in vivo* recombination. The shuffled pool can then be screened in a yeast two hybrid system to select library members which bind said predetermined polypeptide sequence (e. g., and SH2 domain) or which bind an alternate predetermined polypeptide sequence (e.g., an SH2 domain from another protein species).

An approach to identifying polypeptide sequences which bind to a predetermined polypeptide sequence has been to use a so-called "two-hybrid" system wherein the predetermined polypeptide sequence is present in a fusion protein (Chien et al, 1991). This approach identifies protein-protein interactions *in vivo* through reconstitution of a transcriptional activator (Fields and Song, 1989), the yeast Gal4 transcription protein. Typically, the method is based on the properties of the yeast Gal4 protein, which consists of separable domains responsible for DNA-binding and transcriptional activation. Polynucleotides encoding two hybrid proteins, one consisting of the yeast Gal4 DNA-binding domain fused to a polypeptide sequence of a known protein and the other consisting of the Gal4 activation domain fused to a polypeptide sequence of a second protein, are constructed and introduced into a yeast host cell. Intermolecular binding between the two fusion proteins reconstitutes the Gal4 DNA-binding domain with the Gal4 activation domain, which leads to the transcriptional activation of a reporter gene (e.g., *lacZ*, *HIS3*) which is operably linked to a Gal4 binding site. Typically, the two-hybrid method is used to identify novel polypeptide sequences which interact with a known protein (Silver and Hunt, 1993; Durfee et al, 1993; Yang et al, 1992; Luban et al, 1993; Hardy et al, 1992; Bartel et al, 1993; and Vojtek et al, 1993). However, variations of the two-hybrid method have been used to identify mutations of a known protein that affect its binding to a second known protein (Li and Fields, 1993; Lalo et al, 1993; Jackson et al, 1993; and

Madura et al, 1993). Two-hybrid systems have also been used to identify interacting structural domains of two known proteins (**Bardwell et al, 1993; Chakrabarty et al, 1992; Staudinger et al, 1993; and Milne and Weaver 1993**) or domains responsible for oligomerization of a single protein (**Iwabuchi et al, 1993; Bogerd et al, 1993**).

5 Variations of two-hybrid systems have been used to study the *in vivo* activity of a proteolytic enzyme (**Dasmahapatra et al, 1992**). Alternatively, an *E. coli*/BCCP interactive screening system (**Germino et al, 1993; Guarente, 1993**) can be used to identify interacting protein sequences (i.e., protein sequences which heterodimerize or form higher order heteromultimers). Sequences selected by a two-hybrid system can be
10 pooled and shuffled and introduced into a two-hybrid system for one or more subsequent rounds of screening to identify polypeptide sequences which bind to the hybrid containing the predetermined binding sequence. The sequences thus identified can be compared to identify consensus sequence(s) and consensus sequence kernels.

15 In general, standard techniques of recombination DNA technology are described in various publications (e.g. **Sambrook et al, 1989; Ausubel et al, 1987; and Berger and Kimmel, 1987**), each of which is incorporated herein in its entirety by reference. Polynucleotide modifying enzymes were used according to the manufacturer's recommendations. Oligonucleotides were synthesized on an Applied Biosystems Inc.
20 Model 394 DNA synthesizer using ABI chemicals. If desired, PCR amplimers for amplifying a predetermined DNA sequence may be selected at the discretion of the practitioner.

One microgram samples of template DNA are obtained and treated with U.V.
25 light to cause the formation of dimers, including TT dimers, particularly purine dimers. U.V. exposure is limited so that only a few photoproducts are generated per gene on the template DNA sample. Multiple samples are treated with U.V. light for varying periods of time to obtain template DNA samples with varying numbers of dimers from U.V. exposure.

30

A random priming kit which utilizes a non-proofreading polymease (for example, Prime-It II Random Primer Labeling kit by Stratagene Cloning Systems) is utilized to generate different size polynucleotides by priming at random sites on templates which are prepared by U.V. light (as described above) and extending along the templates. The priming protocols such as described in the Prime-It II Random Primer Labeling kit may be utilized to extend the primers. The dimers formed by U.V. exposure serve as a roadblock for the extension by the non-proofreading polymerase. Thus, a pool of random size polynucleotides is present after extension with the random primers is finished.

10

The present invention is further directed to a method for generating a selected mutant polynucleotide sequence (or a population of selected polynucleotide sequences) typically in the form of amplified and/or cloned polynucleotides, whereby the selected polynucleotide sequences(s) possess at least one desired phenotypic characteristic (e.g., encodes a polypeptide, promotes transcription of linked polynucleotides, binds a protein, and the like) which can be selected for. One method for identifying hybrid polypeptides that possess a desired structure or functional property, such as binding to a predetermined biological macromolecule (e.g., a receptor), involves the screening of a large library of polypeptides for individual library members which possess the desired structure or functional property conferred by the amino acid sequence of the polypeptide.

15
20

In one embodiment, the present invention provides a method for generating libraries of displayed polypeptides or displayed antibodies suitable for affinity interaction screening or phenotypic screening. The method comprises (1) obtaining a first plurality of selected library members comprising a displayed polypeptide or displayed antibody and an associated polynucleotide encoding said displayed polypeptide or displayed antibody, and obtaining said associated polynucleotides or copies thereof wherein said associated polynucleotides comprise a region of substantially identical sequences, optimally introducing mutations into said polynucleotides or copies, (2) pooling the polynucleotides or copies, (3) producing

25
30

smaller or shorter polynucleotides by interrupting a random or particularized priming and synthesis process or an amplification process, and (4) performing amplification, preferably PCR amplification, and optionally mutagenesis to homologously recombine the newly synthesized polynucleotides.

5

It is a particularly preferred object of the invention to provide a process for producing hybrid polynucleotides which express a useful hybrid polypeptide by a series of steps comprising:

- (a) producing polynucleotides by interrupting a polynucleotide amplification or synthesis process with a means for blocking or interrupting the amplification or synthesis process and thus providing a plurality of smaller or shorter polynucleotides due to the replication of the polynucleotide being in various stages of completion;
- (b) adding to the resultant population of single- or double-stranded polynucleotides one or more single- or double-stranded oligonucleotides, wherein said added oligonucleotides comprise an area of identity in an area of heterology to one or more of the single- or double-stranded polynucleotides of the population;
- (c) denaturing the resulting single- or double-stranded oligonucleotides to produce a mixture of single-stranded polynucleotides, optionally separating the shorter or smaller polynucleotides into pools of polynucleotides having various lengths and further optionally subjecting said polynucleotides to a PCR procedure to amplify one or more oligonucleotides comprised by at least one of said polynucleotide pools;
- (d) incubating a plurality of said polynucleotides or at least one pool of said polynucleotides with a polymerase under conditions which result in annealing of said single-stranded polynucleotides at regions of identity between the single-stranded polynucleotides and thus forming of a mutagenized double-stranded polynucleotide chain;
- (e) optionally repeating steps (c) and (d);
- (f) expressing at least one hybrid polypeptide from said polynucleotide chain, or chains; and
- (g) screening said at least one hybrid polypeptide for a useful activity.

30

In a preferred aspect of the invention, the means for blocking or interrupting the amplification or synthesis process is by utilization of U.V. light, DNA adducts, DNA binding proteins.

5 In one embodiment of the invention, the DNA adducts, or polynucleotides comprising the DNA adducts, are removed from the polynucleotides or polynucleotide pool, such as by a process including heating the solution comprising the DNA fragments prior to further processing.

10 Having thus disclosed exemplary embodiments of the present invention, it should be noted by those skilled in the art that the disclosures are exemplary only and that various other alternatives, adaptations and modifications may be made within the scope of the present invention. Accordingly, the present invention is not limited to the specific embodiments as illustrated herein.

15 Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following examples are to be considered illustrative and thus are not limiting of the remainder of the disclosure in any way whatsoever.

20

Example 1**Generation of Random Size Polynucleotides Using U.V. Induced Photoproducts**

- One microgram samples of template DNA are obtained and treated with U.V. light to cause the formation of dimers, including TT dimers, particularly purine dimers.
- 5 U.V. exposure is limited so that only a few photoproducts are generated per gene on the template DNA sample. Multiple samples are treated with U.V. light for varying periods of time to obtain template DNA samples with varying numbers of dimers from U.V. exposure.

- A random priming kit which utilizes a non-proofreading polymease (for
- 10 example, Prime-It II Random Primer Labeling kit by Stratagene Cloning Systems) is utilized to generate different size polynucleotides by priming at random sites on templates which are prepared by U.V. light (as described above) and extending along the templates. The priming protocols such as described in the Prime-It II Random Primer Labeling kit may be utilized to extend the primers. The dimers formed by U.V.
- 15 exposure serve as a roadblock for the extension by the non-proofreading polymerase. Thus, a pool of random size polynucleotides is present after extension with the random primers is finished.

Example 2**Isolation of Random Size Polynucleotides**

Polynucleotides of interest which are generated according to Example 1 are are
gel isolated on a 1.5% agarose gel. Polynucleotides in the 100-300 bp range are cut out
5 of the gel and 3 volumes of 6 M NaI is added to the gel slice. The mixture is incubated
at 50 °C for 10 minutes and 10 µl of glass milk (Bio 101) is added. The mixture is spun
for 1 minute and the supernatant is decanted. The pellet is washed with 500 µl of
Column Wash (Column Wash is 50% ethanol, 10mM Tris-HCl pH 7.5, 100 mM NaCl
and 2.5 mM EDTA) and spin for 1 minute, after which the supernatant is decanted. The
10 washing, spinning and decanting steps are then repeated. The glass milk pellet is
resuspended in 20µl of H₂O and spun for 1 minute. DNA remains in the aqueous phase.

Example 3**Shuffling of Isolated Random Size 100-300bp Polynucleotides**

The 100-300 bp polynucleotides obtained in Example 2 are recombined in an annealing mixture (0.2 mM each dNTP, 2.2 mM MgCl₂, 50 mM KCl, 10 mM Tris-HCl
5 ph 8.8, 0.1% Triton X-100, 0.3 μ; Taq DNA polymerase, 50 μl total volume) without adding primers. A Robocycler by Stratagene was used for the annealing step with the following program: 95 °C for 30 seconds, 25-50 cycles of [95 °C for 30 seconds, 50 - 60 °C (preferably 58 °C) for 30 seconds, and 72 °C for 30 seconds] and 5 minutes at 72 °C. Thus, the 100-300 bp polynucleotides combine to yield double-stranded
10 polynucleotides having a longer sequence. After separating out the reassembled double-stranded polynucleotides and denaturing them to form single stranded polynucleotides, the cycling is optionally again repeated with some samples utilizing the single strands as template and primer DNA and other samples utilizing random
15 primers in addition to the single strands.

Example 4**Screening of Polypeptides from Shuffled Polynucleotides**

The polynucleotides of Example 3 are separated and polypeptides are expressed therefrom. The original template DNA is utilized as a comparative control by obtaining
5 comparative polypeptides therefrom. The polypeptides obtained from the shuffled polynucleotides of Example 3 are screened for the activity of the polypeptides obtained from the original template and compared with the activity levels of the control. The shuffled polynucleotides coding for interesting polypeptides discovered during screening are compared further for secondary desirable traits. Some shuffled
10 polynucleotides corresponding to less interesting screened polypeptides are subjected to reshuffling.

Example 5

Directed Evolution an Enzyme by Saturation Mutagenesis

Site-Saturation Mutagenesis: To accomplish site-saturation mutagenesis every
5 residue (316) of a dehalogenase enzyme was converted into all 20 amino acids by site
directed mutagenesis using 32-fold degenerate oligonucleotide primers, as follows:

1. A culture of the dehalogenase expression construct was grown and a preparation of
the plasmid was made
- 10 2. Primers were made to randomize each codon - they have the common structure
 $X_{20}NN(G/T)X_{20}$
3. A reaction mix of 25 ul was prepared containing ~50 ng of plasmid template, 125 ng
of each primer, 1X native Pfu buffer, 200 uM each dNTP and 2.5 U native Pfu DNA
polymerase
- 15 4. The reaction was cycled in a Robo96 Gradient Cycler as follows:
Initial denaturation at 95°C for 1 min
20 cycles of 95°C for 45 sec, 53°C for 1 min and 72°C for 11 min
Final elongation step of 72°C for 10 min
5. The reaction mix was digested with 10 U of DpnI at 37°C for 1 hour to digest the
20 methylated template DNA
6. Two ul of the reaction mix were used to transform 50 ul of XL1-Blue MRF' cells
and the entire transformation mix was plated on a large LB-Amp-Met plate yielding
200-1000 colonies
7. Individual colonies were toothpicked into the wells of 96-well microtiter plates
25 containing LB-Amp-IPTG and grown overnight
8. The clones on these plates were assayed the following day

Screening: Approximately 200 clones of mutants for each position were grown in liquid media (384 well microtiter plates) and screened as follows:

- 5 1. Overnight cultures in 384-well plates were centrifuged and the media removed. To each well was added 0.06 mL 1 mM Tris/SO₄²⁻ pH 7.8.
2. Made 2 assay plates from each parent growth plate consisting of 0.02 mL cell suspension.
3. One assay plate was placed at room temperature and the other at elevated
10 temperature (initial screen used 55°C) for a period of time (initially 30 minutes).
4. After the prescribed time 0.08 mL room temperature substrate (TCP saturated 1 mM Tris/SO₄²⁻ pH 7.8 with 1.5 mM NaN₃ and 0.1 mM bromothymol blue) was added to each well.
5. Measurements at 620 nm were taken at various time points to generate a
15 progress curve for each well.
6. Data were analyzed and the kinetics of the cells heated to those not heated were compared. Each plate contained 1-2 columns (24 wells) of unmutated 20F12 controls.
7. Wells that appeared to have improved stability were re-grown and tested under
20 the same conditions.

Following this procedure nine single site mutations appeared to confer increased thermal stability on the enzyme. Sequence analysis was performed to determine of the exact amino acid changes at each position that were specifically responsible for the
25 improvement. In sum, the improvement was conferred at 7 sites by one amino acid change alone, at an eighth site by each of two amino acid changes, and at a ninth site by each of three amino acid changes. Several mutants were then made each having a plurality of these nine beneficial site mutations in combination; of these two mutants proved superior to all the other mutants, including those with single point mutations.

Example 6**Direct expression cloning using end-selection**

An esterase gene was amplified using 5'phosphorylated primers in a standard
5 PCR reaction (10 ng template; PCR conditions: 3' 94 C; [1' 94 C; 1' 50 C; 1'30'' 68 C]
x 30; 10' 68 C.

Forward Primer = 9511TopF

(CTAGAAGGGAGGAGAATTACATGAAGCGGCTTTTAGCCC)

10 Reverse Primer = 9511TopR (AGCTAAGGGTCAAGGCCGCACCCGAGG)

The resulting PCR product (ca.1000 bp) was gel purified and quantified.

A vector for expression cloning, pASK3 (Institut fuer Bioanalytik, Goettingen,
Germany), was cut with *Xba* I and *Bgl* II and dephosphorylated with CIP.

15

0.5 pmoles Vaccina Topoisomerase I (Invitrogen, Carlsbad, CA) was added to
60 ng (ca. 0.1 pmole) purified PCR product for 5' 37 C in buffer NEB I (New England
Biolabs, Beverly, MA) in 5 µl total volume.

The topogated PCR product was cloned into the vector pASK3 (5 µl, ca. 200 ng in NEB

20 I) for 5' at room temperature.

This mixture was dialyzed against H₂O for 30'.

2 µl were used for electroporation of DH10B cells (Gibco BRL, Gaithersburg, MD).

Efficiency: Based on the actual clone numbers this method can produce 2×10^6
25 clones per µg vector. All tested recombinants showed esterase activity after induction
with anhydrotetracycline.

Example 7

Dehalogenase Thermal Stability

This invention provides that a desirable property to be generated by directed evolution is exemplified in a limiting fashion by an improved residual activity (e.g. an enzymatic activity, an immunoreactivity, an antibiotic activity, etc.) of a molecule upon subjection to altered environment, including what may be considered a harsh environment, for a specified time. Such a harsh environment may comprise any combination of the following (iteratively or not, and in any order or permutation): an elevated temperature (including a temperature that may cause denaturation of a working enzyme), a decreased temperature, an elevated salinity, a decreased salinity, an elevated pH, a decreased pH, an elevated pressure, a decreased pressure, and an change in exposure to a radiation source (including uv radiation, visible light, as well as the entire electromagnetic spectrum).

15

The following example shows an application of directed evolution to evolve the ability of an enzyme to regain &/or retain activity upon exposure to an elevated temperature.

Every residue (316) of a dehalogenase enzyme was converted into all 20 amino acids by site directed mutagenesis using 32-fold degenerate oligonucleotide primers. These mutations were introduced into the already rate-improved variant Dhla 20F12. Approximately 200 clones of each position were grown in liquid media (384 well microtiter plates) to be screened. The screening procedure was as follows:

- 25 1. Overnight cultures in 384-well plates were centrifuged and the media removed. To each well was added 0.06 mL 1 mM Tris/SO₄²⁻ pH 7.8.
2. The robot made 2 assay plates from each parent growth plate consisting of 0.02 mL cell suspension.
3. One assay plate was placed at room temperature and the other at elevated
30 temperature (initial screen used 55°C) for a period of time (initially 30 minutes).

4. After the prescribed time 0.08 mL room temperature substrate (TCP saturated 1 mM Tris/SO₄²⁻ pH 7.8 with 1.5 mM NaN₃ and 0.1 mM bromothymol blue) was added to each well. TCP = trichloropropane.
5. Measurements at 620 nm were taken at various time points to generate a progress curve for each well.
6. Data were analyzed and the kinetics of the cells heated to those not heated were compared. Each plate contained 1-2 columns (24 wells) of un-mutated 20F12 controls.
7. Wells that appeared to have improved stability were regrown and tested under the same conditions.

Following this procedure nine single site mutations appeared to confer increased thermal stability on Dhla-20F12. Sequence analysis showed that the following changes were beneficial:

- 15 D89G
- F91S
- T159L
- G189Q, G189V
- 20 I220L
- N238T
- W251Y
- P302A, P302L, P302S, P302K
- P302R/S306R

- 25 Only two sites (189 and 302) had more than one substitution. The first 5 on the list were combined (using G189Q) into a single gene (this mutant is referred to as "Dhla5"). All changes but S306R were incorporated into another variant referred to as Dhla8.

- 30 Thermal stability was assessed by incubating the enzyme at the elevated temperature (55°C and 80°C) for some period of time and activity assay at 30°C. Initial rates were

plotted vs. time at the higher temperature. The enzyme was in 50 mM Tris/SO₄ pH 7.8 for both the incubation and the assay. Product (Cl⁻) was detected by a standard method using Fe(NO₃)₃ and HgSCN. Dhla 20F12 was used as the *de facto* wild type. The apparent half-life (T_{1/2}) was calculated by fitting the data to an exponential decay

5 function.

These results are shown in Figure 1.

LITERATURE CITED

Unless otherwise indicated, all references cited herein (supra and infra) are incorporated by reference in their entirety.

- 5 Barret AJ, et al., eds.: Enzyme Nomenclature: Recommendations of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology. San Diego: Academic Press, Inc., 1992.
- Boyce COL, ed.: Novo's Handbook of Practical Biotechnology. 2nd ed. Bagsvaerd,
10 Denmark, 1986.
- Drauz K, Waldman H, eds.: Enzyme Catalysis in Organic Synthesis: A Comprehensive Handbook. Vol. 1. New York: VCH Publishers, 1995.
- 15 Drauz K, Waldman H, eds.: Enzyme Catalysis in Organic Synthesis: A Comprehensive Handbook. Vol. 2. New York: VCH Publishers, 1995.
- Foster GD, Taylor SC, eds.: Plant Virology Protocols: From Virus Isolation to Transgenic Resistance. Methods in Molecular Biology, Vol. 81. New Jersey: Humana
20 Press Inc., 1998.
- Franks F, ed.: Protein Biotechnology: Isolation, Characterization, and Stabilization. New Jersey: Humana Press Inc., 1993.
- 25 Godfrey T, West S, eds.: Industrial Enzymology. 2nd ed. London: Macmillan Press Ltd, 1996.
- Gottschalk G: Bacterial Metabolism. 2nd ed. New York: Springer-Verlag Inc., 1986.
- 30 Gresshoff PM, ed.: Technology Transfer of Plant Biotechnology. Current Topics in Plant Molecular Biology. Boca Raton: CRC Press, 1997.

Griffin HG, Griffin AM, eds.: PCR Technology: Current Innovations. Boca Raton: CRC Press, Inc., 1994.

- 5 Hansen G, Chilton MD: Lessons in gene transfer to plants by a gifted microbe. *Curr Top Microbiol Immunol* 240:21-57, 1999.

Hartmann HT, et al.: Plant Propagation: Principles and Practices. 6th ed. New Jersey: Prentice Hall, Inc., 1997.

10

Perun TJ, Propst CL, eds.: Computer-Aided Drug Design: Methods and Applications. New York: Marcel Dekker, Inc., 1989.

- Owen MRL, Pen J: Transgenic Plants: A Production System for Industrial and
15 Pharmaceutical Proteins. Chichester: John Wiley & Sons, 1996.

Segel IH: Enzyme Kinetics: Behavior and Analysis of Rapid Equilibrium and Steady-State Enzyme Systems. New York: John Wiley & Sons, Inc., 1993.

- 20 White JS, White DC: Source Book of Enzymes. Boca Raton: CRC Press, 1997.

Wong CH, Whitesides GM: Enzymes in Synthetic Organic Chemistry. Vol. 12. New York: Elsevier Science Publications, 1995.

- 25 WO 97/35966; Filed Mar. 20, 1997, Published Oct. 2, 1997. Minshull J, Stemmer WP: Methods and Compositions for Cellular and Metabolic Engineering.

- WO 98/31837; Filed Jan. 16, 1998, Published July 23, 1998. Delcardayre SB, Tobin MB, Stemmer WP, Minshull, J: Evolution of Whole Cells and Organisms by Recursive
30 Sequence Recombination.

WO 98/37223; Filed Feb. 18 1998, Published Aug. 27, 1998. Pang SZ, Gonsalves D, Jan FJ: DNA Construct to Confer Multiple Traits on Plants.

Alting-Mecs MA and Short JM: Polycos vectors: a system for packaging filamentous
5 phage and phagemid vectors using lambda phage packaging extracts. *Gene* 137:1, 93-100, 1993.

Arkin AP and Youvan DC: An algorithm for protein engineering: simulations of recursive ensemble mutagenesis. *Proc Natl Acad Sci USA* 89(16):7811-7815, (Aug 15)
10 1992.

Arnold FH: Protein engineering for unusual environments. *Current Opinion in Biotechnology* 4(4):450-455, 1993.

15 Ausubel FM, et al Editors. Current Protocols in Molecular Biology, Vols. 1 and 2 and supplements. (a.k.a. "The Red Book") Greene Publishing Assoc., Brooklyn, NY, ©1987.

Ausubel FM, et al Editors. Current Protocols in Molecular Biology, Vols. 1 and 2 and
20 supplements. (a.k.a. "The Red Book") Greene Publishing Assoc., Brooklyn, NY, ©1989.

Ausubel FM, et al Editors. Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology. Greene Publishing Assoc.,
25 Brooklyn, NY, ©1989.

Ausubel FM, et al Editors. Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology, 2nd Edition. Greene Publishing Assoc., Brooklyn, NY, ©1992.

30

- Barbas CF 3d, Bain JD, Hoekstra DM, Lerner RA: Semisynthetic combinatorial antibody libraries: a chemical solution to the diversity problem. *Proc Natl Acad Sci USA* 89(10):4457-4461, 1992.
- 5 Bardwell AJ, Bardwell L, Johnson DK, Friedberg EC: Yeast DNA recombination and repair proteins Rad1 and Rad10 constitute a complex in vivo mediated by localized hydrophobic domains. *Mol Microbiol* 8(6):1177-1188, 1993.
- Bartel P, Chien CT, Sternglanz R, Fields S: Elimination of false positives that arise in
10 using the two-hybrid system. *Biotechniques* 14(6):920-924, 1993.
- Beaudry AA and Joyce GF: Directed evolution of an RNA enzyme. *Science* 257(5070):635-641, 1992.
- 15 Berger and Kimmel, Methods in Enzymology, Volume 152, Guide to Molecular Cloning Techniques. Academic Press, Inc., San Diego, CA, ©1987. (Cumulative Subject Index: Volumes 135-139, 141-167, 1990, 272 pp.)
- Bevan M: Binary Agrobacterium vectors for plant transformation. *Nucleic Acids
20 Research* 12(22):8711-21, 1984.
- Biocca S, Pierandrei-Amaldi P, Cattaneo A: Intracellular expression of anti-p21ras single chain Fv fragments inhibits meiotic maturation of xenopus oocytes. *Biochem Biophys Res Commun* 197(2):422-427, 1993.
- 25 Bird et al. *Plant Mol Biol* 11:651, 1988..
- Bogerd HP, Fridell RA, Blair WS, Cullen BR: Genetic evidence that the Tat proteins of human immunodeficiency virus types 1 and 2 can multimerize in the eukaryotic cell
30 nucleus. *J Virol* 67(8):5030-5034, 1993.

- Brederode FT, Koper-Zawrthoff EC, Bol JF: Complete nucleotide sequence of alfalfa mosaic virus RNA 4, *Nucleic Acids Research* 8(10):2213-23, 1980.
- Breitling F, Dubel S, Seehaus T, Klewinghaus I, Little M: A surface expression vector
5 for antibody screening. *Gene* 104(2):147-153, 1991.
- Brown NL, Smith M: Cleavage specificity of the restriction endonuclease isolated from *Haemophilus gallinarum* (Hga I). *Proc Natl Acad Sci U S A* 74(8):3213-6, (Aug) 1977.
- 10 Burton DR, Barbas CF 3d, Persson MA, Koenig S, Chanock RM, Lerner RA: A large array of human monoclonal antibodies to type 1 human immunodeficiency virus from combinatorial libraries of asymptomatic seropositive individuals. *Proc Natl Acad Sci U S A* 88(22):10134-7, (Nov 15) 1991.
- 15 Caldwell RC and Joyce GF: Randomization of genes by PCR mutagenesis. *PCR Methods Appl* 2(10):28-33, 1992.
- Caton AJ and Koprowski H: Influenze virus hemagglutinin-specific antibodies
isolatedf froma combinatorial expression library are closely related to the immune
20 response of the donor. *Proc Natl Acad Sci USA* 87(16):6450-6454, 1990.
- Chakraborty T, Martin JF, Olson EN: Analysis of the oligomerization of myogenin and E2A products in vivo using a two-hybrid assay system. *J Biol Chem* 267(25):17498-501, 1992.
- 25 Chang CN, Landolfi NF, Queen C: Expression of antibody Fab domains on bacteriophage surfaces. Potential use for antibody selection. *J Immunol* 147(10):3610-4, (Nov 15) 1991.

- Chaudhary VK, Batra JK, Gallo MG, Willingham MC, FitzGerald DJ, Pastan I: A rapid method of cloning functional variable-region antibody genes in *Escherichia coli* as single-chain immunotoxins. *Proc Natl Acad Sci USA* 87(3):1066-1070, 1990.
- 5 Chien CT, Bartel PL, Sternglanz R, Fields S: The two-hybrid system: a method to identify and clone genes for proteins that interact with a protein of interest. *Proc Natl Acad Sci USA* 88(21):9578-9582, 1991.
- Chiswell DJ, McCafferty J: Phage antibodies: will new 'coliclonal' antibodies replace monoclonal antibodies? *Trends Biotechnol* 10(3):80-84, 1992.
- 10 Chothia C and Lesk AM: Canonical structures for the hypervariable regions of immunoglobulins. *J Mol Biol* 196(4):901-917, 1987.
- 15 Chothia C, Lesk AM, Tramontano A, Levitt M, Smith-Gill SJ, Air G, Sheriff S, Padlan EA, Davies D, Tulip WR, et al: Conformations of immunoglobulin hypervariable regions. *Nature* 342(6252):877-883, 1989.
- Clackson T, Hoogenboom HR, Griffiths AD, Winter G: Making antibody fragments using phage display libraries. *Nature* 352(6336):624-628, 1991.
- 20 Conrad M, Topal MD: DNA and spermidine provide a switch mechanism to regulate the activity of restriction enzyme *Nae I*. *Proc Natl Acad Sci U S A* 86(24):9707-11, (Dec) 1989.
- 25 Coruzzi G, Broglie R, Edwards C, Chua NH: Tissue-specific and light-regulated expression of a pea nuclear gene encoding the small subunit of ribulose-1,5-bisphosphate carboxylase. *EMBO J* 3(8):1671-9, 1984.

- Dasmahapatra B, DiDomenico B, Dwyer S, Ma J, Sadowski I, Schwartz J: A genetic system for studying the activity of a proteolytic enzyme. *Proc Natl Acad Sci USA* 89(9):4159-4162, 1992.
- 5 Davis LG, Dibner MD, Battey JF. Basic Methods in Molecular Biology. Elsevier, New York, NY, ©1986.
- Delegrave S and Youvan DC. *Biotechnology Research* 11:1548-1552, 1993.
- 10 DeLong EF, Wu KY, Prezelin BB, Jovine RV: High abundance of Archaea in Antarctic marine picoplankton. *Nature* 371(6499):695-697, 1994.
- Deng SJ, MacKenzie CR, Sadowska J, Michniewicz J, Young NM, Bundle Dr, Narang SA: Selection of antibody single-chain variable fragments with improved carbohydrate binding by phage display. *J Biol Chem* 269(13):9533-9538, 1994.
- 15
- Duan L, Bagasra O, Laughlin MA, Oakes JW, Pomerantz RJ: Potent inhibition of human immunodeficiency virus type 1 replication by an intracellular anti-Rev single-chain antibody. *Proc Natl Acad Sci USA* 91(11):5075-5079, 1994.
- 20
- Durfee T, Becherer K, Chen PL, Yeh SH, Yang Y, Kilburn AE, Lee WH, Elledge SJ: The retinoblastoma protein associates with the protein phosphatase type 1 catalytic subunit. *Genes Dev* 7(4):555-569, 1993.
- 25
- Ellington AD and Szostak JW: In vitro selection of RNA molecules that bind specific ligands. *Nature* 346(6287):818-822, 1990.
- Fields S and Song O: A novel genetic system to detect protein-protein interactions. *Nature* 340(6230):245-246, 1989.
- 30

- Firek S, Draper J, Owen MR, Gandecha A, Cockburn B, Whitlam GC: Secretion of a functional single-chain Fv protein in transgenic tobacco plants and cell suspension cultures. *Plant Mol Biol* 23(4):861-870, 1993.
- 5 Forsblom S, Rigler R, Ehrenberg M, Philipson L: Kinetic studies on the cleavage of adenovirus DNA by restriction endonuclease Eco RI. *Nucleic Acids Res* 3(12):3255-69, (Dec) 1976.
- Germino FJ, Wang ZX, Weissman SM: Screening for in vivo protein-protein
10 interactions. *Proc Natl Acad Sci USA* 90(3):933-937, 1993.
- Gingeras TR, Brooks JE: Cloned restriction/modification system from *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 80(2):402-6, 1983 (Jan).
- 15 Gluzman Y: SV40-transformed simian cells support the replication of early SV40 mutants. *Cell* 23(1):175-182, 1981.
- Gruber M, Schodin BA, Wilson ER, Kranz DM: Efficient tumor cell lysis mediated by a bispecific single chain antibody expressed in *Escherichia coli*. *J Immunol*
20 152(11):5368-5374, 1994.
- Guarente L: Strategies for the identification of interacting proteins. *Proc Natl Acad Sci USA* 90(5):1639-1641, 1993.
- 25 Guilley H, Dudley RK, Jonard G, Balazs E, Richards KE: Transcription of Cauliflower mosaic virus DNA: detection of promoter sequences, and characterization of transcripts. *Cell* 30(3):763-73, 1982.
- Hardy CF, Sussel L, Shore D: A RAP1-interacting protein involved in transcriptional
30 silencing and telomere length regulation. *Genes Dev* 6(5):801-814, 1992.

- Hawkins RE and Winter G: Cell selection strategies for making antibodies from variable gene libraries: trapping the memory pool. *Eur J Immunol* 22(3):867-870, 1992.
- 5 Holvoet P, Laroche Y, Lijnen HR, Van Hoef B, Brouwers E, De Cock F, Lauwereys M, Gansemans Y, Collen D: Biochemical characterization of single-chain chimeric plasminogen activators consisting of a single-chain Fv fragment of a fibrin-specific antibody and single-chain urokinase. *Eur J Biochem* 210(3):945-952, 1992.
- 10 Honjo T, Alt FW, Rabbitts TH (eds): *Immunoglobulin genes*. Academic Press: San Diego, CA, pp. 361-368, ©1989.
- Hoogenboom HR, Griffiths AD, Johnson KS, Chiswell DJ, Judson P, Winter G: Multi-subunit proteins on the surface of filamentous phage: methodologies for displaying
- 15 antibody (Fab) heavy and light chains. *Nucleic Acids Res* 19(15):4133-4137, 1991.
- Huse WD, Sastry L, Iverson SA, Kang AS, Alting-Mees M, Burton DR, Benkovic SJ, Lerner RA: Generation of a large combinatorial library of the immunoglobulin repertoire in phage lambda. *Science* 246(4935):1275-1281, 1989.
- 20 Huston JS, Levinson D, Mudgett-Hunter M, Tai MS, Novotney J, Margolies MN, Ridge RJ, Brucoleri RE, Haber E, Crea R, et al: Protein engineering of antibody binding sites: recovery of specific activity in an anti-digoxin single-chain Fv analogue produced in *Escherichia coli*. *Proc Natl Acad Sci USA* 85(16):5879-5883, 1988.
- 25 Iwabuchi K, Li B, Bartel P, Fields S: Use of the two-hybrid system to identify the domain of p53 involved in oligomerization. *Oncogene* 8(6):1693-1696, 1993.
- Jackson AL, Pahl PM, Harrison K, Rosamond J, Sclafani RA: Cell cycle regulation of
- 30 the yeast Cdc7 protein kinase by association with the Dbf4 protein. *Mol Cell Biol* 13(5):2899-2908, 1993.

Johnson S and Bird RE: *Methods Enzymol* 203:88, 1991.

5 Kabat et al: Sequences of Proteins of Immunological Interest, 4th Ed. U.S. Department of Health and Human Services, Bethesda, MD (1987)

Kang AS, Barbas CF, Janda KD, Benkovic SJ, Lerner RA: Linkage of recognition and replication functions by assembling combinatorial antibody Fab libraries along phage surfaces. *Proc Natl Acad Sci USA* 88(10):4363-4366, 1991.

10 Kettleborough CA, Ansell KH, Allen RW, Rosell-Vives E, Gussow DH, Bendig MM: Isolation of tumor cell-specific single-chain Fv from immunized mice using phage-antibody libraries and the re-construction of whole antibodies from these antibody fragments. *Eur J Immunol* 24(4):952-958, 1994.

15 Kruger DH, Barcak GJ, Reuter M, Smith HO: EcoRII can be activated to cleave refractory DNA recognition sites. *Nucleic Acids Res* 16(9):3997-4008, (May 11) 1988.

20 Lalo D, Carles C, Sentenac A, Thuriaux P: Interactions between three common subunits of yeast RNA polymerases I and III. *Proc Natl Acad Sci USA* 90(12):5524-5528, 1993.

25 Laskowski M Sr: Purification and properties of venom phosphodiesterase. *Methods Enzymol* 65(1):276-84, 1980.

Lefkovits I and Pernis B, Editors. Immunological Methods, Vols. I and II. Academic Press, New York, NY. Also Vol. III published in Orlando and Vol. IV published in San Diego. ©1979-.

30 Ivan Lefkovits, Editor. Immunology methods manual : the comprehensive sourcebook of techniques. Academic Press, San Diego, ©1997.

- Lerner RA, Kang AS, Bain JD, Burton DR, Barbas CF 3d: Antibodies without immunization. *Science* 258(5086):1313-1314, 1992.
- Leung, D.W., et al, *Technique*, 1:11-15, 1989.
- 5 Li B and Fields S: Identification of mutations in p53 that affect its binding to SV40 large T antigen by using the yeast two-hybrid system. *FASEB J* 7(10):957-963, 1993.
- Lilley GG, Doelzal O, Hillyard CJ, Bernard C, Hudson PJ: Recombinant single-chain
10 antibody peptide conjugates expressed in Escherichia coli for the rapid diagnosis of HIV. *J Immunol Methods* 171(2):211-226, 1994.
- Lowman HB, Bass SH, Simpson N, Wells JA: Selecting high-affinity binding proteins by monovalent phage display. *Biochemistry* 30(45):10832-10838, 1991.
- 15 Luban J, Bossolt KL, Franke EK, Kalpana GV, Goff SP: Human immunodeficiency virus type 1 Gag protein binds to cyclophilins A and B. *Cell* 73(6):1067-1078, 1993.
- Madura K, Dohmen RJ, Varshavsky A: N-recognin/Ubc2 interactions in the N-end rule
20 pathway. *J Biol Chem* 268(16):12046-54, (Jun 5) 1993.
- Marks JD, Hoogenboom HR, Bonnert TP, McCafferty J, Griffiths AD, Winter G: By-passing immunization. Human antibodies from V-gene libraries displayed on phage. *J Mol Biol* 222(3):581-597, 1991.
- 25 Marks JD, Griffiths Ad, Malmqvist M, Clackson TP, Bye JM, Winter G: By-passing immunization: building high affinity human antibodies by chain shuffling. *Biotechnology (N Y)* 10(7):779-783, 1992.
- 30 Marks JD, Hoogenboom HR, Griffiths AD, Winter G: Molecular evolution of proteins on filamentous phage. Mimicking the strategy of the immune system. *J Biol Chem* 267(23):16007-16010, 1992.

- Maxam AM, Gilbert W: Sequencing end-labeled DNA with base-specific chemical cleavages. *Methods Enzymol* 65(1):499-560, 1980.
- 5 McCafferty J, Griffiths AD, Winter G, Chiswell DJ: Phage antibodies: filamentous phage displaying antibody variable domains. *Nature* 348(6301):552-554, 1990.
- Miller JH. A Short Course in Bacterial Genetics: A Laboratory Manual and Handbook for Escherichia coli and Related Bacteria (see inclusively p. 445). Cold Spring Harbor
10 Laboratory Press, Plainview, NY, ©1992.
- Milne GT and Weaver DT: Dominant negative alleles of RAD52 reveal a DNA repair/recombination complex including Rad51 and Rad52. *Genes Dev* 7(9):1755-1765, 1993.
- 15 Mullinax RL, Gross EA, Amberg JR, Hay BN, Hogrefe HH, Kubtiz MM, Greener A, Alting-Mees M, Ardourel D, Short JM, et al: Identification of human antibody fragment clones specific for tetanus toxoid in a bacteriophage lambda immunoexpression library. *Proc natl Acad Sci USA* 87(20):8095-9099, 1990.
- 20 Nath K, Azzolina BA: in *Gene Amplification and Analysis* (ed. Chirikjian JG), vol. 1, p. 113, Elsevier North Holland, Inc., New York, New York, ©1981.
- Needleman SB and Wunsch CD: A general method applicable to the search for
25 similarities in the amino acid sequence of two proteins. *J Mol Biol* 48(3):443-453, 1970.
- Nelson M, Christ C, Schildkraut I: Alteration of apparent restriction endonuclease recognition specificities by DNA methylases. *Nucleic Acids Res* 12(13):5165-73, 1984
30 (Jul 11).

Nicholls PJ, Johnson VG, Andrew SM, Hoogenboom HR, Raus JC, Youle RJ:
Characterization of single-chain antibody (sFv)-toxin fusion proteins produced in vitro
in rabbit reticulocyte lysate. *J Biol Chem* 268(7):5302-5308, 1993.

- 5 Oller AR, Vanden Broek W, Conrad M, Topal MD: Ability of DNA and spermidine to
affect the activity of restriction endonucleases from several bacterial species.
Biochemistry 30(9):2543-9, (Mar 5) 1991.

- Owens RJ and Young RJ: The genetic engineering of monoclonal antibodies. *J*
10 *Immunol Methods* 168(2):149-165, 1994.

Pearson WR and Lipman DJ: Improved tools for biological sequence comparison.
Proc Natl Acad Sci USA 85(8):2444-2448, 1988.

- 15 Pein CD, Reuter M, Meisel A, Cech D, Kruger DH: Activation of restriction
endonuclease EcoRII does not depend on the cleavage of stimulator DNA. *Nucleic*
Acids Res 19(19):5139-42, (Oct 11) 1991.

- Persson MA, Caothien RH, Burton DR: Generation of diverse high-affinity human
20 monoclonal antibodies by repertoire cloning. *Proc Natl Acad Sci USA* 88(6):2432-
2436, 1991.

- Queen C, Foster J, Stauber C, Stafford J: Cell-type specific regulation of a kappa
immunoglobulin gene by promoter and enhance elements. *Immunol Rev* 89:49-68,
25 1986.

- Qiang BQ, McClelland M, Poddar S, Spokauskas A, Nelson M: The apparent
specificity of NotI (5'-GCGGCCGC-3') is enhanced by M.FnuDII or M.BepI
methyltransferases (5'-mCGCG-3'): cutting bacterial chromosomes into a few large
30 pieces. *Gene* 88(1):101-5, (Mar 30) 1990.

Raleigh EA, Wilson G: Escherichia coli K-12 restricts DNA containing 5-methylcytosine. *Proc Natl Acad Sci U S A* 83(23):9070-4, (Dec) 1986.

Reidhaar-Olson JF and Sauer RT: Combinatorial cassette mutagenesis as a probe of the
5 informational content of protein sequences. *Science* 241(4861):53-57, 1988.

Riechmann L and Weill M: Phage display and selection of a site-directed randomized single-chain antibody Fv fragment for its affinity improvement. *Biochemistry* 32(34):8848-8855, 1993.

10

Roberts RJ, Macelis D: REBASE--restriction enzymes and methylases. *Nucleic Acids Res* 24(1):223-35, (Jan 1) 1996.

Ryan AJ, Royal CL, Hutchinson J, Shaw CH: Genomic sequence of a 12S seed storage
15 protein from oilseed rape (Brassica napus c.v. jet neuf). *Nucl Acids Res* 17(9):3584, 1989.

Sambrook J, Fritsch EF, Maniatis T. Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, ©1982.

20

Sambrook J, Fritsch EF, Maniatis T. Molecular Cloning: A Laboratory Manual. Second Edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, ©1989.

25 Scopes RK. Protein Purification: Principles and Practice. Springer-Verlag, New York, NY, © 1982.

Silver SC and Hunt SW 3d: Techniques for cloning cDNAs encoding interactive transcriptional regulatory proteins. *Mol Biol Rep* 17(3):155-165, 1993.

30

Smith TF, Waterman MS. *Adv Appl Math* 2: 482-end of article, 1981.

- Smith TF, Waterman MS: Overlapping genes and information theory. *J Theor Biol* 91(2):379-80, (Jul 21) 1981.
- 5 Smith TF, Waterman MS: Identification of common molecular subsequences. *J Mol Biol* 147(1):195-7, (Mar 25) 1981.
- Smith TF, Waterman MS, Fitch WM: Comparative biosequence metrics. *J Mol Evol* S18(1):38-46, 1981.
- 10 Staudinger J, Perry M, Elledge SJ, Olson EN: Interactions among vertebrate helix-loop-helix proteins in yeast using the two-hybrid system. *J Biol Chem* 268(7):4608-4611, 1993.
- 15 Stemmer WP, Morris SK, Wilson BS: Selection of an active single chain Fv antibody from a protein linker library prepared by enzymatic inverse PCR. *Biotechniques* 14(2):256-265, 1993.
- Stemmer WP: DNA shuffling by random fragmentation and reassembly: in vitro
- 20 recombination for molecular evolution. *Proc Natl Acad Sci USA* 91(22):10747-10751, 1994.
- Sun D, Hurley LH: Effect of the (+)-CC-1065-(N3-adenine)DNA adduct on in vitro DNA synthesis mediated by Escherichia coli DNA polymerase. *Biochemistry* 31:10,
- 25 2822-9, (Mar 17) 1992,
- Tague BW, Dickinson CD, Chrispeels MJ: A short domain of the plant vacuolar protein phytohemagglutinin targets invertase to the yeast vacuole. *Plant Cell* 2(6):533-46, (June) 1990.
- 30 Takahashi N, Kobayashi I: Evidence for the double-strand break repair model of bacteriophage lambda recombination. *Proc Natl Acad Sci U S A* 87(7):2790-4, (Apr) 1990.

- Thiesen HJ and Bach C: Target Detection Assay (TDA): a versatile procedure to determine DNA binding sites as demonstrated on SP1 protein. *Nucleic Acids Res* 18(11):3203-3209, 1990.
- 5 Thomas M, Davis RW: Studies on the cleavage of bacteriophage lambda DNA with EcoRI Restriction endonuclease. *J Mol Biol* 91(3):315-28, (Jan 25) 1975.
- Tingey SV, Walker EL, Corruzzi GM: Glutamine synthetase genes of pea encode distinct polypeptides which are differentially expressed in leaves, roots and
10 nodules. *EMBO J* 6(1):1-9, 1987.
- Topal MD, Thresher RJ, Conrad M, Griffith J: NaeI endonuclease binding to pBR322 DNA induces looping. *Biochemistry* 30(7):2006-10, (Feb. 19) 1991.
- 15 Tramontano A, Chothia C, Lesk AM: Framework residue 71 is a major determinant of the position and conformation of the second hypervariable region in the VH domains of immunoglobulins. *J Mol Biol* 215(1):175-182, 1990.
- Tuerk C and Gold L: Systematic evolution of ligands by exponential enrichment: RNA
20 ligands to bacteriophage T4 DNA polymerase. *Science* 249(4968):505-510, 1990.
- van de Poll ML, Lafleur MV, van Gog F, Vrieling H, Meerman JH: N-acetylated and deacetylated 4'-fluoro-4-aminobiphenyl and 4-aminobiphenyl adducts differ in their ability to inhibit DNA replication of single-stranded M13 in vitro and of single-stranded
25 phi X174 in Escherichia coli. *Carcinogenesis* 13(5):751-8, (May) 1992.
- Vojtek AB, Hollenberg SM, Cooper JA: Mammalian Ras interacts directly with the serine/threonine kinase Raf. *Cell* 74(1):205-214, 1993.

Wenzler H, Mignery G, Fisher L, Park W: Sucrose-regulated expression of a chimeric potato tuber gene in leaves of transgenic tobacco plants. *Plant Mol Biol* 13(4):347-54, 1989.

- 5 Williams and Barclay, in Immunoglobulin Genes, The Immunoglobulin Gene Superfamily

Winnacker EL. From Genes to Clones: Introduction to Gene Technology. VCH Publishers, New York, NY, ©1987.

10

Winter G and Milstein C: Man-made antibodies. *Nature* 349(6307):293-299, 1991.

Yang X, Hubbard EJ, Carlson M: A protein kinase substrate identified by the two-hybrid system. *Science* 257(5070):680-2, (Jul 31) 1992.

15

USPN 4,683,195; Filed Feb. 7, 1986, Issued Jul 28, 1987. Mullis KB, Erlich HA, Arnheim N, Horn GT, Saiki RK, Scharf SJ: Process for Amplifying, Detecting, and/or Cloning Nucleic Acid Sequences.

20

USPN 4,683,202; Filed Oct. 25, 1985, Issued Jul. 28, 1987. Mullis KB: Process for Amplifying Nucleic Acid Sequences.

USPN 4,704,362; Filed Nov. 5, 1979, Issued Nov. 3, 1987. Itakura K, Riggs AD:

- 25 Recombinant Cloning Vehicle Microbial Polypeptide Expression.

WO 88/08453; Filed Apr. 14, 1988, Published Nov. 3, 1988. Alakhov JB, Baranov, VI, Ovodov SJ, Ryabova LA, Spirin AS: Method of Obtaining Polypeptides in Cell-Free Translation System.

30

WO 90/05785; Filed Nov. 15, 1989, Published May 31, 1990. Schultz P: Method for Site-Specifically Incorporating Unnatural Amino Acids into Proteins.

WO 90/07003; Filed Jan. 27, 1989, Published June 28, 1990. Baranov VI, Morozov IJ, Spirin AS: Method for Preparative Expression of Genes in a Cell-free System of Conjugated Transcription/translation.

5

WO 91/02076; Filed June 14, 1990, Published Feb. 21, 1991. Baranov VI, Ryabova LA, Yarchuk OB, Spirin AS: Method for Obtaining Polypeptides in a Cell-free System.

10 WO 91/05058; Filed Oct. 5, 1989, Published Apr. 18, 1991. Kawasaki G: Cell-free Synthesis and Isolation of Novel Genes and Polypeptides.

WO 91/17271; Filed May 1, 1990, Published Nov. 14, 1991. Dower WJ, Cwirla SE: Recombinant Library Screening Methods.

15 WO 91/18980; Filed May 13, 1991, Published Dec. 12, 1991. Devlin JJ: Compositions and Methods for Identifying Biologically Active Molecules.

WO 91/19818; Filed June 20, 1990, Published Dec. 26, 1991. Dower WJ, Cwirla SE, Barrett RW: Peptide Library and Screening Systems.

20

WO 92/02536; Filed Aug. 1, 1991, Published Feb. 20, 1992. Gold L, Tuerk C: Systematic Polypeptide Evolution by Reverse Translation.

25 WO 92/03918; Filed Aug. 28, 1991, Published Mar. 19, 1992. Lonberg N, Kay RM: Transgenic Non-human Animals Capable of Producing Heterologous Antibodies.

WO 92/03918; Filed Aug. 28, 1991, Published Mar. 19, 1992. Lonberg N, Kay RM: Transgenic Non-human Animals Capable of Producing Heterologous Antibodies.

30 WO 92/05258; Filed Sept. 17, 1991, Published Apr. 2, 1992. Fincher GB: Gene Encoding Barley Enzyme.

WO 92/14843; Filed Feb. 21, 1992, Published Sept. 3, 1992. Toole JJ, Griffin LC, Bock LC, Latham JA, Muenchau DD, Krawczyk S: Aptamers Specific for Biomolecules and Method of Making.

5

WO 93/08278; Filed Oct. 15, 1992, Published Apr. 29, 1993. Schatz PJ, Cull MG, Miller JF, Stemmer WP: Peptide Library and Screening Method.

WO 93/12227; Filed Dec 17, 1992, Published June 24, 1993. Lonberg, N; Kay RM:
10 Transgenic Non-human Animals Capable of Producing Heterologous Antibodies.

WO 93/12227; Filed Dec. 17, 1992, Published June 24, 1993. Lonberg N, Kay RM:
Transgenic Non-human Animals Capable of Producing Heterologous Antibodies.

15 WO 94/25585; Filed Apr. 25, 1994, Published Nov. 10, 1994. Lonberg, N, Kay RM:
Transgenic Non-human Animals Capable of Producing Heterologous Antibodies.

WO 94/25585; Filed Apr. 25, 1994, Published Nov. 10, 1994. Lonberg N, Kay RM:
Transgenic Non-human Animals Capable of Producing Heterologous Antibodies.

20

- Arslan T, Abraham AT, Hecht SM: Structurally altered substrates for DNA topoisomerase I. Effects of inclusion of a single 3'-deoxynucleotide within the scissile strand. *Nucleosides Nucleotides* 1998 Jan-Mar;17(1-3):515-30.
- Aupeix K, Toulme JJ: Binding of chemically-modified oligonucleotides to the double-stranded stem of an RNA hairpin. *Nucleosides Nucleotides* 1999 Jun-Jul;18(6-7):1647-50.
- Bazzanini R, Manfredini S, Durini E, Groschel B, Cinatl J, Balzarini J, De Clercq E, Imbach JL, Perigaud C, Gosselin G: Prodrugs of Ara-CMP and Ara-AMP with a S-acyl-2-thioethyl (SATE) biolabile phosphate protecting group: synthesis and biological evaluation. *Nucleosides Nucleotides* 1999 Apr-May;18(4-5):971-2.
- Blackburn GM, Liu X, Rosler A, Brenner C: Two hydrolase resistant analogues of diadenosine 5',5'''-P1,P3-triphosphate for studies with Fhit, the human fragile histidine triad protein. *Nucleosides Nucleotides* 1998 Jan-Mar;17(1-3):301-8.
- Bridson PK, Lin X, Melman N, Ji XD, Jacobson KA: Synthesis and adenosine receptor affinity of 7-beta-D-ribofuranosylxanthine. *Nucleosides Nucleotides* 1998 Apr;17(4):759-68.

- Brodin P, Gottikh M, Auclair C, Mouscadet JF: Inhibition of HIV-1 integration by mono- & bi-functionalized triple helix forming oligonucleotides. *Nucleosides Nucleotides* 1999 Jun-Jul;18(6-7):1717-8.
- Creighton TE: Proteins Structures and Molecular Principles. New York: W. H. Freeman and Co., 1984.
- De Clercq E: Carbocyclic adenosine analogues as S-adenosylhomocysteine hydrolase inhibitors and antiviral agents: recent advances. *Nucleosides Nucleotides* 1998 Jan-Mar;17(1-3):625-34.
- de Zwart M, Link R, von Frijtag Drabbe Kunzel JK, Cristalli G, Jacobson KA, Townsend-Nicholson A, IJzerman AP: A functional screening of adenosine analogues at the adenosine A2B receptor: a search for potent agonists. *Nucleosides Nucleotides* 1998 Jun;17(6):969-85.
- Egron D, Arzumanov AA, Dyatkina NB, Krayevsky A, Imbach JL, Aubertin AM, Gosselin G, Perigaud C: Synthesis, anti-HIV activity and stability studies of 3'-azido-2',3'-dideoxythymidine 5'-fluorophosphate. *Nucleosides Nucleotides* 1999 Apr-May;18(4-5):983-4
- Gianolio DA, McLaughlin LW: Synthesis and triplex forming properties of pyrimidine derivative containing extended functionality. *Nucleosides Nucleotides* 1999 Aug;18(8):1751-69.
- Gottikh MB, Volkov EM, Romanova EA, Oretskaya TS, Shabarova ZA: Synthesis of oligonucleotide-intercalator conjugates capable to inhibit HIV-1 DNA integration. *Nucleosides Nucleotides* 1999 Jun-Jul;18(6-7):1645-6.
- Hotoda H, Koizumi M, Ohmine T, Furukawa H, Nishigaki T, Abe K, Kosaka T, Tsutsumi S, Sone J, Kaneko M: Biologically active oligodeoxyribonucleotides. 10: anti-HIV-1 activity and stability of modified hexanucleotides containing glycerol-skeleton. *Nucleosides Nucleotides* 1998 Jan-Mar;17(1-3):243-52.
- JP10113194; Filed 19971022, Published 19980506. Donnelly, JJ ;Dwarki, VJ ;Liu, MA ;Montgomery, DL ;Parker, S ;Shiver, JW ;Ulmer JB: Nucleic Acid Preparation.

- Kang SH, Sinhababu AK, Cho MJ: Synthesis and biological activity of bis(pivaloyloxymethyl) ester of 2'-azido-2'-deoxyuridine 5'-monophosphate. *Nucleosides Nucleotides* 1998 Jun;17(6):1089-98.
- Krayevsky A, Arzumanov A, Shirokova E, Dyatkina N, Victorova L, Jasko M, Alexandrova L: dNTP modified at triphosphate residues: substrate properties towards DNA polymerases and stability in human serum. *Nucleosides Nucleotides* 1998 Jan-Mar;17(1-3):681-93.
- Krayevsky AA, Dyatkina NB, Semizarov DG, Victorova LS, Shirokova EA, Theil F, Von Janta Lipinski MJ, Gosselin G, Imbach JL: Reasons and limits of substrate activity of modified L-dNTP in DNA biosynthesis. *Nucleosides Nucleotides* 1999 Apr-May;18(4-5):863-4.
- Kvasnyuk EI, Mikhailopulo IA, Suhadolnik RJ, Henderson EE, Muto NF, Iacono KT, Homon J, Pfeleiderer W: Synthesis and biological activity of 2',5'-oligoadenylate trimers containing 5'-terminal 5'-amino-5'-deoxy- and 5'-amino-3',5'-dideoxyadenosine derivatives. *Nucleosides Nucleotides* 1999 Jun-Jul;18(6-7):1483-4.
- Liu J, Skradis A, Kolar C, Kolath J, Anderson J, Lawson T, Talmadge J, Gmeiner WH: Increased cytotoxicity and decreased in vivo toxicity of FdUMP[10] relative to 5-FU. *Nucleosides Nucleotides* 1999 Aug;18(8):1789-802.
- Lutz MJ, Will DW, Breipohl G, Benner SA, Uhlmann E: Synthesis of a monocharged peptide nucleic acid (PNA) analog and its recognition as substrate by DNA polymerases. *Nucleosides Nucleotides* 1999 Mar;18(3):393-401.
- Monaco V, van de Wetering KI, Meeuwenoord NJ, van den Elst HA, Stuivenberg HR, Visse R, van der Kaaden JC, Moolenaar GF, Verhoeven EE, Goosen N, van der Marel GA, van Boom JH: Synthesis and biological evaluation of modified DNA fragments for the study of nucleotide excision repair in *E. coli*. *Nucleosides Nucleotides* 1999 Jun-Jul;18(6-7):1339-41.
- Morozova OV, Kolpashchikov DM, Ivanova TM, Godovikova TS: Synthesis of new photocross-linking 5-C-base-substituted UTP analogs and their application in highly selective affinity labelling of the tick-borne encephalitis virus RNA replicase proteins. *Nucleosides Nucleotides* 1999 Jun-Jul;18(6-7):1513-4.

Nguyen-Ba N, Chan L, Quimpere M, Turcotte N, Lee N, Mitchell H, Bedard J:
Design and SAR study of a novel class of nucleotide analogues as potent anti-HCMV
agents. *Nucleosides Nucleotides* 1999 Apr-May;18(4-5):821-7.

Pandolfi D, Rauzi F, Capobianco ML: Evaluation of different types of end-capping modifications on the stability of oligonucleotides toward 3'- and 5'-exonucleases.

Nucleosides Nucleotides 1999 Sep;18(9):2051-69.

Pankiewicz KW, Lesiak-Watanabe K: Novel mycophenolic adenine bis(phosphonate)s as potent anticancer agents and inducers of cells differentiation.

Nucleosides Nucleotides 1999 Apr-May;18(4-5):927-32.

Perrin DM, Garestier T, Helene C: Expanding the catalytic repertoire of nucleic acid catalysts: simultaneous incorporation of two modified deoxyribonucleoside triphosphates bearing ammonium and imidazolyl functionalities. *Nucleosides Nucleotides* 1999 Mar;18(3):377-91.

Pfundheller HM, Koshkin AA, Olsen CE, Wengel J: Evaluation of oligonucleotides containing two novel 2'-O-methyl modified nucleotide monomers: a 3'-C-allyl and a 2'-O,3'-C-linked bicyclic derivative. *Nucleosides Nucleotides* 1999 Sep;18(9):2017-30.

Ramasamy KS, Stoisavljevic V: Synthesis and biophysical studies of modified oligonucleotides containing acyclic amino alcohol nucleoside analogs. *Nucleosides Nucleotides* 1999 Aug;18(8):1845-61.

Schinazi RF, Lesnikowski ZJ: Boron containing oligonucleotides. *Nucleosides Nucleotides* 1998 Jan-Mar;17(1-3):635-47.

Secrist JA 3rd, Parker WB, Allan PW, Bennett LL Jr, Waud WR, Truss JW, Fowler AT, Montgomery JA, Ealick SE, Wells AH, Gillespie GY, Gadi VK, Sorscher EJ: Gene therapy of cancer: activation of nucleoside prodrugs with E. coli purine nucleoside phosphorylase. *Nucleosides Nucleotides* 1999 Apr-May;18(4-5):745-57.

Shirokova EA, Shipitsin AV, Victorova LS, Dyatkina NB, Goryunova LE, Beabealashvilli RS, Hamilton CJ, Roberts SM, Krayevsky AA: Modified nucleoside 5'-triphosphonates as a new type of antiviral agents. *Nucleosides Nucleotides* 1999 Apr-May;18(4-5):1027-8.

Srivastava TK, Friedhoff P, Pingoud A, Katti SB: Application of oligonucleoside methylphosphonates in the studies on phosphodiester hydrolysis by Serratia endonuclease. *Nucleosides Nucleotides* 1999 Sep;18(9):1945-60.

Stattel JM, Yanachkov I, Wright GE: Synthesis and biochemical study of N2-(p-n-butylphenyl)-2'-deoxyguanosine 5'-(alpha,beta-imido)triphosphate (BuPdGMPNHPP): a non-substrate inhibitor of B family DNA polymerases. *Nucleosides Nucleotides* 1998 Aug;17(8):1505-13.

Terato H, Morita H, Ohyama Y, Ide H: Novel modification of 5-formyluracil by cysteine derivatives in aqueous solution. *Nucleosides Nucleotides* 1998 Jan-Mar;17(1-3):131-41.

Tomikawa A, Seno M, Sato-Kiyotaki K, Ohtsuki C, Hirai T, Yamaguchi T, Kawaguchi T, Yoshida S, Saneyoshi M: Synthetic nucleosides and nucleotides. 40. Selective inhibition of eukaryotic DNA polymerase alpha by 9-(beta-D-arabinofuranosyl)-2-(p-n-butylanilino) adenine 5'-triphosphate (BuAaraATP) and its 2'-up azido analog: synthesis and enzymatic evaluations. *Nucleosides Nucleotides* 1998 Jan-Mar;17(1-3):487-501.

US5580859; Filed 19940318, Issued 19961203. Felgner, PL. ;Wolff, JA. ;Rhodes, GH. ;Malone, RW. ;Carson, DA.: Delivery of exogenous DNA sequences in a mammal.

US5589466; Filed 19950126, Issued 19961231. Felgner, PL. ;Wolff, JA. ;Rhodes, GH. ;Malone, RW. ;Carson, DA.: Induction of a protective immune response in a mammal by injecting a DNA sequence.

US5641665; Filed 19941128, Issued 19970624. Hobart, PM. ;Margalith, M ;Parker, SE. ;Khatibi, S: Plasmids suitable for IL-2 expression.

US5693622; Filed 19950607, Issued 19971202. Wolff, JA. ;Duke, DJ. ;Felgner, PL.: Expression of exogenous polynucleotide sequences cardiac muscle of a mammal.

US5703055; Filed 19940126, Issued 19971230. Felgner, PL. ;Wolff, JA ;Rhodes, GH. ;Malone, RW ;Carson, DA.: Generation of antibodies through lipid mediated DNA delivery.

US5846946; Filed 19960614, Issued 19981208. Huebner, RC. ;Norman, JA. ;Liang, X ;Carner, KR. ;Barbour, AG. ;Luke, CJ.: Compositions and methods for administering *Borrelia* DNA.

US5910488; Filed 19951201, Issued 19990608. Nabel, GJ. ;Nabel, EG. ;Lew, D ;Marquet, M: Plasmids suitable for gene therapy.

Victorova LS, Semizarov DG, Shirokova EA, Alexandrova LA, Arzumanov AA, Jasko MV, Krayevsky AA: Human DNA polymerases and retroviral reverse transcriptases: selectivity in respect to dNTPs modified at triphosphate residues. *Nucleosides Nucleotides* 1999 Apr-May;18(4-5):1031-2.

von Janta-Lipinski M, Gaertner K, Lehmann C, Scheer H, Schildt J, Matthes E: Protein and RNA of human telomerase as targets for modified oligonucleotides. *Nucleosides Nucleotides* 1999 Jun-Jul;18(6-7):1719-20

WO9011092; Filed 19900321, A1 Published 19901004. Felgner, PL. ;Wolff, JA;Rhodes, GH. ;Malone, RW ;Carson, DA.: Expression Of Exogenous Polynucleotide Sequences In A Vertebrate.

WO9314778; Filed 19930121, A1 Published 19930805. Rhodes, GH. ;Dwarki, VJ.;Felgner, PL;Wang-Felgner, J;Manthorpe, M: Ex Vivo Gene Transfer.

WO9421797; Filed 19940314, A1 Published 19940929. Donnelly, JJ. ;Dwarki, VJ. ;Liu, MA. ;Montgomery, DL. ;Parker, SE. ;Shiver, JW. ;Ulmer, JB.: Nucleic Acid Pharmaceuticals.

WO9633736; Filed 19960426, A1 Published 19961031. Baruch DI; Pasloske BL; Howard, RJ: Malaria Peptides and Vaccines.

WO9735992; Filed 19970317, A1 Published 19971002. Hobart, PM. ;Liang, X: Tetracycline Inducible/Repressible Systems.

WO9926663; Filed 19981120, A2 Published 19990603. Horton, H ;Parker, S;Manthorpe, M; Felgner, P: Treatment Of Cancer Using Cytokine-Expressing Polynucleotides And Compositions Therefor.

WO9941368; Filed 19990210, A2 Published 19990819. Punnonen J, Stemmer WP, Whalen RG; Howard, R: Optimization of Immunomodulatory Properties of Genetic Vaccines.

WO9941369; Filed 19990210, A2 Published 19990819. Punnonen J, Stemmer WP, Whalen RG; Howard, R: Genetic Vaccine Vector Engineering.

WO9941383; Filed 19990210, A1 Published 19990819. Punnonen J, Bass, SH, Whalen, RG, Howard, R, Stemmer, WP: Antigen Library Immunization.

WO9941402; Filed 19990210, A2 Published 19990819. Punnonen J, Stemmer, WP, Howard R, Patten PA: Targeting of Genetic Vaccine Vectors.

CLAIMS

What is claimed is:

- 5 1. A method for producing and isolating a polypeptide having at least one desirable property comprised of the steps of:
- (a) subjecting a starting or parental polynucleotide set to a mutagenesis process so as to produce a progeny polynucleotide set; and
- 10 (b) subjecting the progeny polynucleotide set to an end selection-based screening and enrichment process, so as to select for a desirable subset of the progeny polynucleotide set;
- whereby the above steps can be performed iteratively and in any order and in combination,
- 15 whereby the end selection-based process creates ligation-compatible ends, whereby the creation of ligation-compatible ends is optionally used to facilitate one or more intermolecular ligations, that are preferably directional ligations, within members of the progeny polynucleotide set so as to achieve assembly &/or reassembly mutagenesis,
- 20 whereby the creation of ligation-compatible ends serves to facilitate ligation of the progeny polynucleotide set into an expression vector system and expression cloning,
- whereby the expression cloning of the progeny polynucleotide set serves to generate a polypeptide set,
- 25 whereby the generated polypeptide set can be subjected to an expression screening process, and
- whereby expression screening of the progeny polypeptide set provides a means to identify a desirable species, e.g. a mutant polypeptide or alternatively a polypeptide fragment, that has a desirable property, such as a specific
- 30 enzymatic activity.

2. A method according to claim 1, wherein the mutagenesis process of step (a) is comprised of a process, termed saturation mutagenesis, for generating, from a codon-containing parental polypeptide template, a progeny polypeptide set in which a full range of single amino acid substitutions is represented at each amino acid position, comprising the steps of:
- 5 (a) subjecting a working codon-containing template polynucleotide to polymerase-based amplification using a 32-fold degenerate oligonucleotide for each codon to mutagenized, where each of said 32-fold degenerate oligonucleotides is comprised of a first homologous sequence and a degenerate N,N,G/T triplet sequence, so as to generate a set of progeny polynucleotides; and
- 10 (b) subjecting said set of progeny polynucleotides to recombinant expression such that polypeptides encoded by the progeny polynucleotides are produced;
- 15 whereby the above steps can be performed iteratively and in any order and in combination,
- whereby, said method provides a means for generating all 20 amino acid changes at each amino acid site along a parental polypeptide template, because the degeneracy of the N,N,G/T sequence includes codons for all
- 20 20 amino acids,
- whereby said method also provides a means for generating all possible fragments of a parental polynucleotide template, which in turn can lead to recombinant expression of a set of encoded polypeptide fragments,
- 25 and
- whereby expression screening of the progeny polypeptide set provides a means to identify a desirable species, e.g. a mutant polypeptide or alternatively a polypeptide fragment, that has a desirable property, such as a specific enzymatic activity.
- 30

3. A method according to claim 2, where said 32-fold degenerate oligonucleotide is comprised of a first homologous sequence, a degenerate N,N,G/T triplet sequence, and a second homologous sequence.

5

4. A method according to claim 2, where said 32-fold degenerate oligonucleotide is comprised of a first homologous sequence, a plurality of degenerate N,N,G/T triplets sequences, and a second homologous sequence.

10

5. A method according to claim 2, that is further comprised of the step of:
(c) subjecting a working codon-containing template polynucleotide to polymerase-based amplification using a nondegenerate oligonucleotide for each codon to mutagenized,

15

whereby the use of each nondegenerate oligonucleotide provides a means to generate a single point mutation in a working polynucleotide, which in turn provides a means to recombinantly express an encoded polypeptide having a corresponding single amino acid change as long as the mutation at the polynucleotide level is not silent, and

20

whereby said method also provides a means for generating all possible fragments of a parental polynucleotide template, which in turn can lead to the recombinant expression of a set of encoded polypeptide fragments, and

25

whereby expression screening of the progeny polypeptide set provides a means to identify a desirable species, e.g. a mutant polypeptide or alternatively a polypeptide fragment, that has a desirable property, such as a specific enzymatic activity.

30

1 / 11




Figure 1.

Thermal stability of wild type (20F12) and mutant (Dhla 5) dehalogenase at 55°C. Purified protein in 50 mM Tris/SO₄²⁻ pH 7.8 was incubated at 55°C for variable times and aliquots were removed to assay for trichloropropane (TCP) hydrolysis. The assay was performed in 50 mM Tris/SO₄²⁻ pH 7.8, 30°C with 10 mM TCP. Product (Cl⁻) release was analyzed by a standard Cl⁻ assay. Initial rates (A450/min) were derived from the enzyme progress curves and plotted vs. incubation time. The data are fit to an exponential decay ($C \cdot \exp^{-kt}$) where $t_{1/2} = \ln 2/k$.




Figure 2.

Thermal stability of Dhla 5 dehalogenase at 55°C. Purified protein in 50 mM Tris/SO₄²⁻ pH 7.8 was incubated at 55°C for variable times and aliquots were removed to assay for trichloropropane (TCP) hydrolysis. The assay was performed in 50 mM Tris/SO₄²⁻ pH 7.8, 30°C with 10 mM TCP. Product (Cl⁻) release was analyzed by a standard Cl⁻ assay. Initial rates (A450/min) were derived from the enzyme progress curves and plotted vs. incubation time. The data are fit to an exponential decay ($C \cdot \exp^{-kt}$) where $t_{1/2} = \ln 2/k$.




Figure 3.

Enzyme progress curves for Dhla 5 and 20F12 at 30°C and 55°C. The reaction was performed in 50 mM Tris/ SO_4^{2-} , pH 7.8 with 10 mM TCP using purified protein. Equal amounts of protein were used in each assay. Product (Cl^-) concentration was determined at variable times using a standard Cl^- assay. Dhla 5 appears to have the same specific activity as 20F12 at both 30°C and 55°C but the enzyme continues to turn over after 20F12 is thermally inactivated.




Figure 4




Figure 4.




Figure 5

Thermal stability of Dhla 5 dehalogenase at 80°C. Purified protein in 50 mM Tris/SO₄²⁻ pH 7.8 was incubated at 80°C for variable times and aliquots were removed to assay for trichloropropane (TCP) hydrolysis. The assay was performed in 50 mM Tris/SO₄²⁻ pH 7.8, 30°C with 10 mM TCP. Product (Cl⁻) release was analyzed by a standard Cl⁻ assay. Initial rates (A450/min) were derived from the enzyme progress curves and plotted vs. incubation time. The data are fit to an exponential decay ($C \cdot \exp^{-kt}$) where $t_{1/2} = \ln 2/k$.




Figure 6.

Thermal stability of Dhla 5 and Dhla 8 dehalogenase at 80°C. Purified protein in 50 mM Tris/SO₄²⁻ pH 7.8 was incubated at 80°C for variable times and aliquots were removed to assay for trichloropropane (TCP) hydrolysis. The assay was performed in 50 mM Tris/SO₄²⁻ pH 7.8, 30°C with 10 mM TCP. Product (Cl⁻) release was analyzed by a standard Cl⁻ assay. Initial rates (A450/min) were derived from the enzyme progress curves and relative rates were plotted vs. incubation time. The data are fit to an exponential decay ($C \cdot \exp^{-kt}$) where $t_{1/2} = \ln 2/k$ giving values of 13 and 138 minutes for Dhla 5 and Dhla 8, respectively.




Figure 7.





Figure 8.



Figures 7, 8, & 9.

Differential scanning calorimetry of the three dehalogenase variants. The data were recorded using a Perkin Elmer Pyris 1 at a scan rate of 10°C/min. The protein concentration was 10 mg/ml in 40 μl (0.4 mg total protein). The buffer was 10 mM NaP_i pH 7.5. The melting temperatures derived from the data are: 68°C (20F12), 73°C (Dhla 5) and 73°C (Dhla 8). The data also indicate that 20F12 undergoes irreversible denaturation, Dhla 5 is partially reversible and Dhla 8 is fully reversible.

Figure 10A. End Selection Gene Assembly




Figure 10B. End Selection Gene Assembly

WO 00/53744

PCT/US00/06497

11 / 11

